

Interactive comment on “Autogenic versus allogenic controls on the evolution of a coupled fluvial megafan/mountainous catchment system: numerical modelling and comparison with the Lannemezan megafan system (Northern Pyrenees, France)” by Margaux Mouchené et al.

Anonymous Referee #1

Received and published: 9 September 2016

Referee report on

"Autogenic versus allogenic controls on the evolution of a coupled fluvial megafan/mountainous catchment system: numerical modelling and comparison with the Lannemezan megafan system" by

M. Mouchené et al.

While landscape evolution models have been quite successful in realistically reproduc-

Printer-friendly version

Discussion paper



ing erosional landforms the modeling of depositional environments like alluvial fans or river deltas has shown to be hard. This difficulty can be attributed to the fact that in alluvial fans and deltas erosion and deposition closely interact with each other on the same time and length scale.

In this article Mouchéné and coworkers apply the CIDRE model to study autogenic and allogenic controls and apply their simulation results to the Lannemezan megafan system in Southern France.

The paper is well written and definitely suitable for ESurf. However there are a few points, the authors should clarify before publication can be recommended.

Below my comments & suggestions for the authors:

Introduction:

As mentioned above alluvial fans are erosional/depositional environments which makes them difficult to characterize and model. The authors should emphasize this point more clearly in their introduction to distinct their work from classical landscape evolution model approaches.

Model setup:

Sentence: "We conducted a series of trial runs to adjust the relevant parameters in order to reproduce the first-order morphological traits of the northern Pyrenean foreland. In particular, the values for transport length (L), for the erodibilities of bedrock and sediments (respectively k_{br} and k_{all}) and for the critical shear stress (c) need to be established."

Please clarify how the simulation results were compared to the Pyrenean foreland and describe which measures were used to determine 'similarity'. This point is also related to the following:

Results:

Sentence: "We successfully reproduced the first-order morphology of a fluvial megafan constructed on a low-elevation, stable foreland, from the erosional products of a slowly uplifting mountain-range-like block"

The authors must clearly describe what they mean by "successful reproducing the first order morphology" actually means:

1. What is the "first order morphology the authors compare their results to?" 2. Which measure is used to determine "success"? 3. What was the initial condition which was put into the model?

See for example Topography of inland deltas, Seybold et al. GRL(2010) where the surface morphology of alluvial fans is characterized by means of simulations, lab experiments and Digital elevation model analysis and contrasted to another depositional environment specifically fluvial deltas.

Another good description of characteristics of alluvial fans can be found in: Alluvial Fans and their Natural Distinction from Rivers Based on Morphology, Hydraulic Processes, Sedimentary Processes, and Facies Assemblages, Blair et al. J. of Sedimentary Res. (1994),

p.6 L.31: What do the authors mean with "the flow is distributive" Do they mean that the system develops a diverging flow network.

Sec. 5.1 and 5.2 are rather descriptive. Do the authors have measures which characterize the formation of the fan and capture the switching of the channel network and the spatial distribution of erosion and deposition?

In summary the authors present a thorough description of their modeling effort but remain shallow in the quantitative interpretation of their results and a clear presentation of what they promised in the title: The contrasting of "autogenic versus allogenic controls" on the formation of depositional fans. I think the paper can make an interesting contribution if focused more clearly on the interpretation of the results using quantitative

Printer-friendly version

Discussion paper



tive measures.

Interactive comment on Earth Surf. Dynam. Discuss., doi:10.5194/esurf-2016-44, 2016.

ESurfD

Interactive
comment

Printer-friendly version

Discussion paper



Interactive comment on “Autogenic versus allogenic controls on the evolution of a coupled fluvial megafan/mountainous catchment system: numerical modelling and comparison with the Lannemezan megafan system (Northern Pyrenees, France)” by Margaux Mouchéné et al.

Margaux Mouchéné et al.

margaux.mouchene@univ-grenoble-alpes.fr

Received and published: 23 September 2016

We thank the reviewer for their encouraging comments, careful criticism of the text and insightful remarks that have helped us clarify and improve the manuscript. We give detailed responses to specific comments below.

The reviewer asked to "clarify how the simulation results were compared to the Pyrenean foreland and describe which measures were used to determine 'similarity' ". We

[Printer-friendly version](#)

[Discussion paper](#)



compared the first order morphology of the landscape produced by the model to that of the Northern Pyrenean landscape: in both the foreland and the mountain we assessed the mean elevation, highest elevation, river spacing, relief (as valley-to-ridge elevation difference); in the foreland we also assessed the length, width and northward slope of the megafan. All these parameters agree between the modeled landscape and DEM of the northern central Pyrenees within 30 % for similarity to be accepted. This will be specified in the corrected version of the manuscript.

The reviewer asked for clarification on the use of the term " distributive " to describe the flow dynamics over the megafan. We should indeed precise that the flow on the fan is alternatively, both in space and time, (1) channelized with multiple, rapid avulsions occurring, and (2) unchannelized, overflowing the riverbed and diverging over the foreland. This pattern can be seen in Figure 3.

The reviewer pointed out that sections 5.1 and 5.2 are " rather descriptive ". These sections describe the processes observed during the building and subsequent abandonment and incision of the megafan, Figures 3 and 4 are provided to support these descriptions. We do provide some quantitative data on the megafan evolution in sections 5.1 and 5.2 and figure captions (mean elevation change, aggradation rates, vertical amplitude of temporary and permanent entrenchment, river spacing, timeframe). However, we acknowledge the lack of quantitative data regarding water and sediment fluxes and the spatial distribution of erosion and deposition. We will provide these data in the corrected version of the manuscript (fluxes and maps of erosion/deposition).

Interactive comment on Earth Surf. Dynam. Discuss., doi:10.5194/esurf-2016-44, 2016.

Printer-friendly version

Discussion paper



Interactive comment on “Autogenic versus allogenic controls on the evolution of a coupled fluvial megafan/mountainous catchment system: numerical modelling and comparison with the Lannemezan megafan system (Northern Pyrenees, France)” by Margaux Mouchéné et al.

H. Sinclair (Referee)

hugh.sinclair@ed.ac.uk

Received and published: 17 November 2016

General comments: The Lannemezan Fan is a very interesting system to analyse, as it comprises a large radiating fan of incised channels, and forms the dominant morphological feature of the Aquitaine basin. This well presented study focuses on a numerical model using the code CIDRE in order to improve understanding of the controls on formation and subsequent incision of the fan. The model generates some interesting output which appears to approximately simulate the topography of the catchment and

[Printer-friendly version](#)

[Discussion paper](#)



a fan system that is broadly comparable to the Lannemezan system. I think there is an interesting story here in relation to controls on local base levels in continental foreland basins and the impact on growth, stability and incision of large alluvial fans; in this case, the analysis suggests autogenic control through capture of drainage networks has played a role in determining the form of the fan. As outlined in the specific comments below, there is a lot of scope for expanding the documentation of the Lannemezan fan. For example, it would be good to understand more on the timing of growth of the fan in relation to deformation of the north Pyrenean thrust wedge, and in terms of the sedimentological evolution of the system, and the evidence for tilting of its stratigraphy rather than its relatively modern terrace system. In these aspects, it is hard to judge whether the parameter combinations and boundary conditions used in the model are wholly suitable for the Lannemezan system. My recommendation is that the manuscript should be refocused primarily on the model as a series of generic experiments without aiming them at simulating one system. But that the aspects of the modelled fans be presented as examples of the autogenic controls likely to control a number of natural systems including the Lannemezan Fan. The developments of local base-levels generated by neighbouring river systems, and the implications for capture and incision of fan systems is very interesting and represents an important insight into the generic processes of alluvial fan growth in foreland basins.

P2 – L20-25 – The final paragraph of the Introduction is aimed at informing the reader of the aims and motivations for the study. The section starts by suggesting that the study will test hypotheses, but then doesn't define a hypothesis, and goes on to describe how the study will explore autogenic versus allogenic controls. This felt a bit vague, and so needs tightening up. I also think the title could reflect a more focused approach.

P3 – L3 – What is a well sorted 'sequence' in this context – are you referring to grain size distributions in the sand, or a clear distinction of sand and mudstone beds?

P3 – If the Lannemezan Fan is the system to which the subsequent modelling is to be compared, then we need to know a bit more about the geological context of the fan,

particularly its upstream catchment, as the title suggests this is an important component of the story. It needs to be stated that the growth of the fan is all post-orogenic (I assume?), and that there is no reason to expect that the feeder catchment has changed in size over the period of growth and erosion of the fan. It would also be important to describe the glacial impact on the catchment as this will have affected the sediment and water flux. Some of the brevity in this section points to the manuscript by Mouch  n   et al that has been submitted. Even if that paper is published by the time this comes out, I think there is a need to summarise in more detail some of the conclusions of that study, and give more background.

Bottom P3 – On initial reading, I was unclear how L is defined – it is stated that this is the transport length, but its not clear what transport this is referring to; L is often used as transport length in diffusion models referring to catchment length (e.. Allen et al., 2013). Now having read on, I realise we are talking about a sediment transport length per unit time – this needs clarification. The parameters for the model need summarising on a figure.

P6 – L5 – The model set-up involves a fault displacement of 0.3mm/yr at the mountain front allowing for the erosion of the range while depositing in the basin. It is suggested that this is reasonable for the Lannemezan Fan – in what way? When did the uplift of the northern Pyrenean thrust wedge relative to the Aquitaine Basin cease? The model reflects syn-orogenic rather than post-orogenic boundary conditions. Given many would argue that the Pyrenees ceased active crustal thickening at around 20 Ma, doesn't the fan represent a dominantly post-orogenic setting with some normal faulting in its recent past?

P4 and 6 – The final set-up is based around modifying key parameters in order to simulate a system that broadly looks like the Lannemezan Fan. I note the other reviewer asks about how this comparison is made, and this is an interesting point. With so many parameters it is impossible for the reader to assess the role of each one and the sensitivity of the system to changes in each of them. Consequently, as is often the case, we

Printer-friendly version

Discussion paper



put a degree of faith in the model output based on a recognition that the values used for each component seem reasonable within the bounds of experiment and empirical evidence. Based on this, I raise the question of the exponents (n and m) on q and S in the fluvial erosion model which are given as 0.6 and 0.7 (Table 1); doesn't this end up with unrealistic looking river profiles? (cf. point raised by other reviewer about comparisons). Similarly, I am curious to understand how the grainsize distributions are captured in the parameter L , and how sensitive the system might be to reasonable changes in grainsize as might be influenced by the onset of glacial erosion for example.

P8 – L25-30. The tilt experiments are interesting, but the tilt values used in the experiments are very likely to represent a minimum relative to the real values. The surface they choose to evaluate whether the fan is tilted is the relatively recent surface that caps the system (I think the author dates it in her thesis at around 300 Ka). However, if the Pyrenees has been in a post-orogenic state for ca. 20 Myr, then it is reasonable to expect that the fan has been tilted much more than this in response to post-orogenic erosion. Which then raises the question of how much of the current morphology of the fan is a record of post-orogenic erosion with channels that are unrelated to the Miocene depositional system. I believe the authors have evidence to indicate that this is not the case, and so I suggest this should be discussed/demonstrated in this manuscript.

P11 –L5 – “This required longer L , which may be interpreted as a smaller settling rate (Davy and Lague, 2009), is consistent with the downstream fining of sediment in the Lannemezan megafan.” Has downstream fining been demonstrated for the Lannemezan fan? This sedimentological evidence for both a radiating depositional system and documented fining rates needs to be summarised in a figure.

P11 – L20- The critical boundary condition effect here seems to be the open boundary at the sides of the basin. Firstly, there appears to be a rather unusual transverse (ie. E-W) set of channels that are regularly spaced, perpendicular to the main fan, and these seem to link to the open boundaries at the margins. What do these represent? Secondly, the lateral open boundary will always result in incision once a lateral channel

[Printer-friendly version](#)[Discussion paper](#)

taps into the main discharge, as it represents the shortest route, and hence steepest channel, to the zero elevation boundary. It suggests that there are no along-strike fans that maintain an elevation comparable to the modelled fan. So this modelled incision due to capture by the shortest route channel is not representative of the processes of base level change along the strike of a foreland basin, unless you're next to the coast.

P12 – The point raised above seems relevant to this discussion about the controls on incision into fans and the comparison to Pepin et al's study. The open boundary which seems to be maintained at zero elevation at the margin will result in steeper channels with higher bed shear stress than those that drain out at the northern margin of the system – therefore, the former will capture the drainage of the latter, and incise into the depositional base-level built by the longer channels. This seems an inevitable consequence of these boundary conditions. The subsequent comparison to the modern channel system, and the evolutionary model proposed in figure 12 requires that the Ariège and Upper Garonne systems are at lower elevations than the Neste, causing a steepening of the lateral channels that link the two, and incision into the previous fan. Is there evidence that these were at a lower elevation? Is the outlet of these rivers lower than the Neste? I can see this as an important mechanism for drainage capture, but I don't think the default model simulations presented here are a valuable guide to the processes of capture in the Aquitaine Basin. The model boundary conditions of a zero elevation margin to the basin are not the same as the capture by a neighbouring river along the strike of a foreland basin.

P14- L20-25. Is this 'uplift' referring to differential surface uplift between the Aquitaine Basin and the Pyrenees or is it a regional uplift relative to sea-level?

P15 – L9 – “Thus, tilting does not appear to play a major role in the evolution of the Lannemezan megafan”. I don't see how we can say this. The only record of tilting is from the very young terraces, whereas the fan itself is at least 20 Myr old. Is there any clear demonstration that the stratigraphy of the Lannemezan fan is not tilted?

Printer-friendly version

Discussion paper



Hugh Sinclair, Edinburgh

Interactive comment on Earth Surf. Dynam. Discuss., doi:10.5194/esurf-2016-44, 2016.

ESurfD

Interactive
comment

Printer-friendly version

Discussion paper



Interactive comment on “Autogenic versus allogenic controls on the evolution of a coupled fluvial megafan/mountainous catchment system: numerical modelling and comparison with the Lannemezan megafan system (Northern Pyrenees, France)” by Margaux Mouchéné et al.

H. Sinclair (Referee)

hugh.sinclair@ed.ac.uk

Received and published: 17 November 2016

General comments: The Lannemezan Fan is a very interesting system to analyse, as it comprises a large radiating fan of incised channels, and forms the dominant morphological feature of the Aquitaine basin. This well presented study focuses on a numerical model using the code CIDRE in order to improve understanding of the controls on formation and subsequent incision of the fan. The model generates some interesting output which appears to approximately simulate the topography of the catchment and a fan system that is broadly comparable to the Lannemezan system. I think there is an interesting story here in relation to controls on local base levels in continental foreland basins and the impact on growth, stability and incision of large alluvial fans; in this case, the analysis suggests autogenic control through capture of drainage networks has played a role in determining the form of the fan. As outlined in the specific comments below, there is a lot of scope for expanding the documentation of the Lannemezan fan. For example, it would be good to understand more on the timing of growth of the fan in relation to deformation of the north Pyrenean thrust wedge, and in terms of the sedimentological evolution of the system, and the evidence for tilting of its stratigraphy rather than its relatively modern terrace system. In these aspects, it is hard to judge whether the parameter combinations and boundary conditions used in the model are wholly suitable for the Lannemezan system. My recommendation is that the manuscript should be refocused primarily on the model as a series of generic experiments without aiming them at simulating one system. But that the aspects of the modelled fans be presented as examples of the autogenic controls likely to control a number of natural systems including the Lannemezan Fan. The developments of local base-levels generated by neighbouring river systems, and the implications for capture and incision of fan systems is very interesting and represents an important insight into the generic processes of alluvial fan growth in foreland basins.

We thank Pr. Sinclair for his encouraging comments, careful criticism of the text and insightful remarks that have helped us clarify and improve the manuscript. We agree that more background information on the Lannemezan system should have been included and have added this to the new version of the manuscript. However, our study was initially set up – in terms of temporal and spatial scaling – to study the Lannemezan megafan in particular; we therefore have not followed the suggestion to refocus this as a generic experiment, because we have only explored a small part of the vast parameter space in that case.

P2 – L20-25 – The final paragraph of the Introduction is aimed at informing the reader of the aims and motivations for the study. The section starts by suggesting that the study will test hypotheses, but then doesn't define a hypothesis, and goes on to describe how

the study will explore autogenic versus allogenic controls. This felt a bit vague, and so needs tightening up. I also think the title could reflect a more focused approach.

We agree with the reviewer that the final paragraph did not state clearly enough the aim of the study; this has been corrected in the new version of the manuscript.

P3 – L3 – What is a well sorted ‘sequence’ in this context – are you referring to grain size distributions in the sand, or a clear distinction of sand and mudstone beds?

We agree with the reviewer that the formulation was misleading, we are referring to a clear sorting of the grain size (upward fining, pebbles and boulders in a very fine matrix) and made this clearer in the revised version.

P3 – If the Lannemezan Fan is the system to which the subsequent modelling is to be compared, then we need to know a bit more about the geological context of the fan, particularly its upstream catchment, as the title suggests this is an important component of the story. It needs to be stated that the growth of the fan is all post-orogenic (I assume?), and that there is no reason to expect that the feeder catchment has changed in size over the period of growth and erosion of the fan. It would also be important to describe the glacial impact on the catchment as this will have affected the sediment and water flux. Some of the brevity in this section points to the manuscript by Mouchéné et al that has been submitted. Even if that paper is published by the time this comes out, I think there is a need to summarise in more detail some of the conclusions of that study, and give more background.

We agree with the reviewer and included more information on the catchment in the revised version.

Bottom P3 – On initial reading, I was unclear how L is defined – it is stated that this is the transport length, but its not clear what transport this is referring to; L is often used as transport length in diffusion models referring to catchment length (e.. Allen et al., 2013). Now having read on, I realise we are talking about a sediment transport length per unit time – this needs clarification. The parameters for the model need summarising on a figure.

As stated page 4 line 1, L is a coefficient that “determines the proportion of incoming sediment flux that is deposited in the cell”. Equation 2 shows that L has the dimension of a length ($D=qs/L$, with D the local deposition rate [LT^{-1}] and qs the incoming sediment flux per unit width [$L^2 T^{-1}$]). We refer readers to the paper by Carretier et al. (2015), which presents the model in much more detail.

P6 – L5 – The model set-up involves a fault displacement of 0.3mm/yr at the mountain front allowing for the erosion of the range while depositing in the basin. It is suggested that this is reasonable for the Lannemezan Fan – in what way? When did the uplift of the northern Pyrenean thrust wedge relative to the Aquitaine Basin cease?

The model reflects syn-orogenic rather than post-orogenic boundary conditions. Given many would argue that the Pyrenees ceased active crustal thickening at around 20 Ma, doesn't the fan represent a dominantly post-orogenic setting with some normal faulting in its recent past?

The reviewer is right in that uplift in the central Pyrenees is thought to have mostly ceased by the Miocene. The uplift is applied in the model to simulate the post-orogenic exhumation observed in the central part of the range (by thermochronological data, e.g. Jolivet et al., 2007) and the low uplift rates measured today (GPS data by Nguyen et al., 2016). It is also required to maintain sufficient relief (comparable to the observed relief in the Pyrenees) on long (10^7 y) timescales.

P4 and 6 – The final set-up is based around modifying key parameters in order to simulate a system that broadly looks like the Lannemezan Fan. I note the other reviewer asks about how this comparison is made, and this is an interesting point. With so many parameters it is impossible for the reader to assess the role of each one and the sensitivity of the system to changes in each of them. Consequently, as is often the case, we put a degree of faith in the model output based on a recognition that the values used for each component seem reasonable within the bounds of experiment and empirical evidence. Based on this, I raise the question of the exponents (n and m) on q and S in the fluvial erosion model which are given as 0.6 and 0.7 (Table 1); doesn't this end up with unrealistic looking river profiles? (cf. point raised by other reviewer about comparisons). Similarly, I am curious to understand how the grainsize distributions are captured in the parameter L, and how sensitive the system might be to reasonable changes in grainsize as might be influenced by the onset of glacial erosion for example.

The comparison between the model and the Lannemezan megafan is based on the first order morphology and timescale: we compare a number of physical (mean elevation, highest elevation, river spacing, relief in the range and foreland; in the foreland we also assessed the length, width and northward slope of the megafan) and temporal (duration of megafan building and incision phases) parameters from the model and the Lannemezan area DEM (ASTER GDEM2). Acceptable similarity is set at <30% difference between the two.

In the CIDRE model, the general detachment law, for both bedrock and sediments, is:

$$E = K(k_t q^m S^n - \tau_c)^p$$

The term $k_t q^m S^n - \tau_c$ is the shear stress, so that $m=0.6$ and $n=0.7$ (see for instance Tucker, 2004, ESPL). τ_c is the critical shear stress.

If $\tau_c=0$, the equation is simplified and the system described by a stream power law

$$E = K q^m S^n$$

where parameters K, m and n "incorporate" the other parameters k_t and p. In this case $m=0.5$ et $n=1$ (see Whipple and Tucker, 1999, JGR).

When $\tau_c > 0$, $k_t \approx 0.05$, p varies between 1 and 2 (depending on the abrasion process), and m and n are 0.6 and 0.7, respectively. We only evaluated cases where $\tau_c > 0$. With such values of m and n, it can be expected that the long river profiles are very concave ($\theta \sim 0.8$). For technical reasons, we did not extract long river profiles on the modeled megafan but the rivers coming from the mountain range are indeed rather concave ($\theta > 0.5$). On the Lannemezan megafan, all rivers exhibit high concavity ($\theta_{mean} = 0.7$).

Grainsize distributions can be monitored in the CIDRE model but we did not explore this parameter in this study.

P8 – L25-30. The tilt experiments are interesting, but the tilt values used in the

experiments are very likely to represent a minimum relative to the real values. The surface they choose to evaluate whether the fan is tilted is the relatively recent surface that caps the system (I think the author dates it in her thesis at around 300 Ka). However, if the Pyrenees has been in a post-orogenic state for ca. 20 Myr, then it is reasonable to expect that the fan has been tilted much more than this in response to post-orogenic erosion. Which then raises the question of how much of the current morphology of the fan is a record of post-orogenic erosion with channels that are unrelated to the Miocene depositional system. I believe the authors have evidence to indicate that this is not the case, and so I suggest this should be discussed/demonstrated in this manuscript.

To evaluate whether the fan is tilted, we use the basal surface of Lannemezan Formation, which caps the Miocene deposits in the apex area, as stated in section 5.3.4. In Mouchen 's thesis and Mouchen  et al (in press), we indeed estimated the abandonment of this surface at ≥ 300 ka. The surface we use for estimating tilting of the fan is older; the deposits at the base of the Lannemezan Formation contain fossils dated by Paris (1975) and Azambre et al. (1989) at the Miocene-Pliocene transition ("Pontien"). We use a scaling law to estimate the original depositional slope of the Pliocene deposits on top of this surface in order to retrieve the amount of tilting of this surface during the Quaternary (the tilting starts at the end of the building phase). The reviewer raises the question of how much of the morphology of the foreland is related to a Miocene depositional system as a megafan: our experiments suggest that the megafan hypothesis (megafan built in ~ 15 My, during the Miocene and subsequently rapidly dissected by river network following its abandonment) is plausible; we do not explore other scenarios to explain the foreland morphology (although this has been done by other authors; e.g. Bonnet et al, 2014, R union des Sciences de la Terre).

P11 –L5 – "This required longer L, which may be interpreted as a smaller settling rate (Davy and Lague, 2009), is consistent with the downstream fining of sediment in the Lannemezan megafan." Has downstream fining been demonstrated for the Lannemezan fan? This sedimentological evidence for both a radiating depositional system and documented fining rates needs to be summarised in a figure.

References to Crouzel (1957) and Azambre and Crouzel (1988), who documented downstream fining on the fan, have been added to the text.

P11 – L20- The critical boundary condition effect here seems to be the open boundary at the sides of the basin. Firstly, there appears to be a rather unusual transverse (ie. E-W) set of channels that are regularly spaced, perpendicular to the main fan, and these seem to link to the open boundaries at the margins. What do these represent? Secondly, the lateral open boundary will always result in incision once a lateral channel taps into the main discharge, as it represents the shortest route, and hence steepest channel, to the zero elevation boundary. It suggests that there are no along-strike fans that maintain an elevation comparable to the modelled fan. So this modelled incision due to capture by the shortest route channel is not representative of the processes of base level change along the strike of a foreland basin, unless you're next to the coast.

Our hypothesis is not base-level change. We propose that the size and shape of the fan is determined by the pre-existing drainage network: the spacing of major drainage axes

(Garonne-Ariège to the east and Adour to the west) controls the maximum size of the fan; they limit its extension and efficiently evacuate sediments to the distant basin. They correspond to the open boundary conditions imposed in the model.

P12 – The point raised above seems relevant to this discussion about the controls on incision into fans and the comparison to Pepin et al's study. The open boundary which seems to be maintained at zero elevation at the margin will result in steeper channels with higher bed shear stress than those that drain out at the northern margin of the system – therefore, the former will capture the drainage of the latter, and incise into the depositional base-level built by the longer channels. This seems an inevitable consequence of these boundary conditions. The subsequent comparison to the modern channel system, and the evolutionary model proposed in figure 12 requires that the Ariège and Upper Garonne systems are at lower elevations than the Neste, causing a steepening of the lateral channels that link the two, and incision into the previous fan. Is there evidence that these were at a lower elevation? Is the outlet of these rivers lower than the Neste? I can see this as an important mechanism for drainage capture, but I don't think the default model simulations presented here are a valuable guide to the processes of capture in the Aquitaine Basin. The model boundary conditions of a zero elevation margin to the basin are not the same as the capture by a neighbouring river along the strike of a foreland basin.

In the model, the elevation of the foreland boundaries is fixed at 0 m, this is indeed an approximation. However, the long river profiles of the Garonne and Adour in the foreland are very flat (<0.5%) and the difference in elevation between the megafan and these rivers in the model is comparable to that of the Lannemezan area. We now discuss this point in the manuscript.

P14- L20-25. Is this 'uplift' referring to differential surface uplift between the Aquitaine Basin and the Pyrenees or is it a regional uplift relative to sea-level?

This value is a GPS-derived vertical velocity and corresponds to the differential uplift of mountain belt with respect to the foreland (the Aquitaine basin is not uplifted).

P15 – L9 – “Thus, tilting does not appear to play a major role in the evolution of the Lannemezan megafan”. I don't see how we can say this. The only record of tilting is from the very young terraces, whereas the fan itself is at least 20 Myr old. Is there any clear demonstration that the stratigraphy of the Lannemezan fan is not tilted?

This sentence has been changed to say that the tilting does not seem to play a major role in the abandonment of the megafan, which is what our experiment was meant to test.

Hugh Sinclair, Edinburgh

Interactive comment on Earth Surf. Dynam. Discuss., doi:10.5194/esurf-2016-44, 2016.

Autogenic versus allogenic controls on the evolution of a coupled fluvial megafan/mountainous catchment system: numerical modelling and comparison with the Lannemezan megafan system (Northern Pyrenees, France)

5 Margaux Mouchéné^{1*}, Peter van der Beek¹, Sébastien Carretier^{2,3}, Frédéric Mouthereau²

¹Université Grenoble Alpes, CNRS, ISTerre, [CS-40700 F-38058](#), Grenoble, France

²GET, Observatoire Midi Pyrénées, Université de Toulouse, CNRS, IRD, 14 avenue E. Belin, F-31400 Toulouse, France

³Department of Geology, FCFM, Universidad de Chile, Santiago, Chile

**now at: Department of Earth and Environmental Sciences, Tulane University, New Orleans, LA 70118, USA*

10 Correspondence to: Margaux Mouchéné (margaux.mouchene@univ-grenoble-alpes.fr)

Abstract. Alluvial megafans are sensitive recorders of landscape evolution, controlled by **both** autogenic processes and allogenic forcing and influenced by the coupled dynamics of the fan with its mountainous catchment. The Lannemezan megafan in the northern Pyrenean foreland was abandoned by its mountainous feeder stream during the Quaternary and subsequently incised, leaving a flight of alluvial terraces along the stream network. We **use numerical models to** explore the relative roles of autogenic processes and external forcing in the building, abandonment and incision of a foreland megafan and compare the results with the inferred evolution of the Lannemezan megafan. Autogenic processes are sufficient to explain the building of a megafan and the long-term entrenchment of its feeding river at time and space scales that match the Lannemezan setting. Climate, through temporal variations in precipitation rate, may have played a role in the episodic pattern of incision at a shorter time-scale. In contrast, base-level changes, tectonic activity in the mountain range or tilting of the foreland through flexural isostatic rebound **do not appear to have played a role in the abandonment of the megafan.**

1 Introduction

Alluvial fans **and megafans** are prominent geomorphic objects of remarkably conical shape, constructed by the accumulation of sediments at the outlet of mountain valleys. They occupy a key position in the sediment routing system and, as such, have been widely used as recorders of external forcing on landscape evolution in a variety of settings. Controls on the building and incision of these deposits, through alternating phases of aggradation and erosion, have been shown to be related to climatic changes (Barnard et al., 2006; Arboleya et al., 2008; Assine et al., 2014), tectonic activity (DeCelles and Cavazza, 1999), base-level oscillations (Harvey, 2002) or to a combination of those factors (Abrams and Chadwick, 1994; Dade and Verdeyen, 2007; Schlunegger and Norton, 2014). Laboratory experiments reproducing alluvial fan dynamics have helped understanding the respective roles of these controls on **fan morphology, facies changes and cyclic erosion/deposition**

Margaux Mouchene 9/1/y 15:33

Supprimé: .

Margaux Mouchene 9/1/y 15:33

Supprimé: 00

Margaux Mouchene 9/1/y 15:35

Supprimé: using numerical modelling

Margaux Mouchene 9/1/y 15:36

Supprimé: unimportant

Margaux Mouchene 9/1/y 15:36

Supprimé: the

processes (Kim and Muto, 2007; Nicholas et al., 2009; Rohais et al., 2011; Guerit et al., 2014). Both analog and numerical modelling studies have shown evidence for autogenic processes that could be of critical importance in fan evolution (Humphrey and Heller, 1995; Coulthard et al., 2002; Nicholas and Quine, 2007). Temporary sediment storage on the fan results in cyclic behaviour, with alternating phases of deposition and incision in the absence of external forcing (eg. Coulthard et al., 2002). This behaviour is expressed in the thresholds (in runoff, slope or shear stress) defined and implemented in the models: a critical value must be reached and exceeded for transport to be effective; after some further time steps, this parameter value decreases below the threshold and deposition occurs again (Schumm, 1979; Roering et al., 1999; Whipple and Tucker, 1999; DiBiase and Whipple, 2011).

Another level of complexity, often overlooked in previous experiments of alluvial systems, comes from the strong coupling and feedbacks between the source catchment and the basin. The specific response time and amplitude of each part of the system to a given forcing may differ and this results in a complex, oscillating erosion signal (Densmore et al., 2007; Humphrey and Heller, 1995; Babault et al., 2005; Carretier and Lucazeau, 2005). Numerical modelling by Pepin et al. (2010) suggested that autogenic processes play a key role in the evolution of such a coupled system submitted to constant external forcing. For these authors, permanent autogenic entrenchment can occur in a coupled catchment-fan system without changes in boundary conditions and external forcing when (i) the transport threshold (critical shear stress) is significant and (ii) progradation is limited by an open boundary with fixed elevation (e.g. a large river system at the foot of the fan).

In the northern foreland of the Pyrenees (France), the Lannemezan megafan was built since the Miocene by the erosional products of the mountainous Neste River catchment, and was abandoned during the Quaternary (Mouchené et al., in press).

The respective roles of climate and tectonics in this evolution remain unresolved. In this study, we seek to test hypotheses on the mechanisms at play in the abandonment and incision of the Lannemezan megafan through numerical modelling of alluvial megafan construction and abandonment. Although the complexity of this natural case might not be fully reproduced by the numerical model, we run a series of model scenarios to explore the respective effect of potential forcing factors including autogenic dynamics, climate change, tectonic tilting, and base level change, on trends and patterns of incision (time and space scales, amplitudes) of the megafan. Disentangling the respective signals of autogenic processes and allogenic forcing requires understanding of (i) the wavelength and amplitude of each signal, (ii) the possible buffering effects of the response times of the fan and of the mountainous catchment, and (iii) the amplification/reduction factors introduced by the coupling of the system.

2 The Lannemezan megafan

Whereas the drainage network in the Pyrenean range is regularly spaced and mostly transverse to the structural trend, rivers of the northwestern foreland spread in a radial pattern over the convex topography of large Miocene alluvial fans (Fig. 1). The Lannemezan megafan is the most prominent geomorphic feature of the northern Pyrenean foreland, with a surface of 13,000 km² and a mean slope of 0.3°. Its characteristic semi-conical shape is outlined by the Garonne River (to the south,

Margaux Mouchene 9/1/y 15:37

Supprimé: i

Margaux Mouchene 9/1/y 15:37

Supprimé: s

Margaux Mouchene 9/1/y 15:37

Supprimé: and time

Margaux Mouchene 9/1/y 15:37

Supprimé: a

Margaux Mouchene 9/1/y 15:38

Supprimé: submitted

Margaux Mouchene 9/1/y 15:38

Supprimé: -

Margaux Mouchene 9/1/y 15:38

Supprimé: will

Margaux Mouchene 6/12/y 11:02

Supprimé: to understand the effect of different potential forcing factors (climate, tectonics, base level change, etc.) on

Margaux Mouchene 9/1/y 15:39

Supprimé: landscape

Margaux Mouchene 6/12/y 11:03

Supprimé: evolution

Margaux Mouchene 9/1/y 15:39

Supprimé: r

east and north) and by the radial river network on its surface. It was built since the middle Miocene (Biteau et al., 2006), while orogenic activity within the Pyrenean range had already waned (Vergès et al., 2002; Sinclair et al., 2005). Molasse-type deposits of middle- to late-Miocene age, with rounded pebbles and boulders in an abundant, clayey and sandy matrix, make up most of the megafan volume (Paris, 1975; Azambre et al., 1989). These are capped by the *Lannemezan Formation*, consisting of (i) an upward-fining, stratified clay and sand sequence, containing strongly weathered gravel and pebbles in a very fine matrix, dated by a Hipparion-bearing fauna at its base as latest Miocene to Pliocene (“Pontico-Pliocene”; Paris, 1975; Azambre et al., 1989), and (ii) a Quaternary sheet of very similar composition.

The Neste River exits the mountain range at the apex of the megafan and thus most probably provided the material building the megafan. Comparing sediment volumes and thermochronology-derived exhumation rates, Mouchéné (2016) showed that the relatively small Neste catchment could have exported a sufficiently large sediment flux to produce all of the megafan deposits. Contrary to the neighbouring Garonne valley, the Neste valley is believed to have mostly been exempt of glaciers during the Quaternary and no glacial features are known on the megafan itself (e.g. Delmas, 2015).

The Lannemezan megafan is currently disconnected from its source catchment and is being incised. The Neste turns sharply to the east at the megafan apex and incises the fan head ~100m vertically, before merging with the larger Garonne River at its mountainous outlet (Fig. 1): this drainage pattern suggests the capture of the Neste by the Garonne. During incision, the rivers of the northern foreland (including the Neste, Garonne and fan rivers) left a series of alluvial terraces. Mouchéné et al. (in press) dated the abandonment of the fan and the onset of incision at $\geq 300\text{ka}$, from ^{10}Be and ^{26}Al cosmogenic nuclide dating of the fan surface and terraces along the Neste valley. The episodic abandonment of these river terraces during incision may be related to changing fluvial dynamics during shifts between Quaternary cold and warm phases (Mouchéné et al., in press).

3 Model description

We use a recent version of the CIDRE code, which models landscape evolution in a continental setting (Carretier et al., 2015). We recall here the main characteristics of the code and refer the reader to Carretier et al. (2015) and references therein for further details.

At the beginning of each time step, a specified volume of water is distributed homogeneously over the cells making up the model surface. The propagation of water and sediment is performed in cascade, from the highest to the lowest cell and following decreasing elevation, to ensure mass conservation. A Multiple-Flow algorithm is used to propagate the water flux to downstream cells proportionally to the slope in each direction (Murray and Paola, 1997; Coulthard et al., 2002; Carretier et al., 2009), allowing a distributary drainage pattern to develop.

3.1 Mass balance

During a time step ∂t , the elevation z of a grid cell changes as follows:

Margaux Mouchene 5/1/y 12:37

Supprimé: well-sorted

Margaux Mouchene 7/1/y 10:15

Supprimé: However, this stream

Margaux Mouchene 9/1/y 15:42

Supprimé: now

Margaux Mouchene 7/1/y 10:14

Supprimé: this point

Margaux Mouchene 9/1/y 15:43

Supprimé: .

Margaux Mouchene 9/1/y 15:43

Supprimé: T

Margaux Mouchene 9/1/y 15:43

Supprimé: and related

Margaux Mouchene 9/1/y 15:44

Supprimé: was dated by Mouchéné et al. (submitted) at ~

Margaux Mouchene 9/1/y 15:45

Supprimé: -

... [1]

Margaux Mouchene 9/1/y 15:45

Supprimé: t

Margaux Mouchene 9/1/y 15:45

Supprimé: which was dated by these authors and

Margaux Mouchene 9/1/y 15:46

Supprimé: submitted

Margaux Mouchene 9/1/y 15:48

Supprimé:

Margaux Mouchene 9/1/y 15:47

Supprimé: . This

Margaux Mouchene 9/1/y 15:47

Supprimé: s

Margaux Mouchene 9/1/y 15:47

Supprimé: for

Margaux Mouchene 9/1/y 15:48

Supprimé: point of a

$$\frac{\partial z}{\partial t} = -\epsilon + D + U \quad (1)$$

where ϵ is a local erosion (detachment or entrainment) rate, D is a local deposition rate and U an uplift or subsidence rate. The local deposition rate D is defined as:

$$D = \frac{q_s}{L} \quad (2)$$

5 with q_s the incoming sediment flux per unit width and L the transport length. The transport length L determines the proportion of incoming sediment flux that is deposited in the cell: a large L results in little deposition, such as a steep slope or high water discharge would favour in natural settings. The cell outflux per unit width q_s is the sum of the sediment detached from a given cell plus the sediment eroded upstream and that crossed this cell without being deposited; it is thus non-local (e.g. Tucker and Bradley, 2010). This approach is generalised for both hillslope and fluvial processes by specifying ϵ and L in both cases.

3.2 Hillslope processes

The approach used by Carretier et al. (2015) is different from the “non-linear” diffusion model proposed by previous authors (Roering et al., 1999; Carretier et al., 2009; 2014). Instead, in this model, the elevation variation results from the difference between a local detachment rate and a deposition rate using Equations 1 and 2 where ϵ erosion rate and transport length L are defined as:

$$\epsilon = \kappa S \quad (3)$$

$$L = \frac{dx}{1-(S/S_c)^2} \quad (4)$$

15 Where κ is an erodibility coefficient, S is the steepest slope and S_c is a critical slope. If the slope is steeper than S_c , ϵ is set such that $S = S_c$. The detachment rate is proportional to the local gradient, but the deposition rate (q_s/L in Equation 2) depends on the slope and critical slope: when $S \ll S_c$, most of the sediment entering a cell is deposited there and when $S \sim S_c$, L becomes infinity and there is no deposition on the cell.

3.3 Fluvial Processes

For fluvial processes, a detachment algorithm including a threshold is used for sediment and bedrock:

$$25 \quad \epsilon = K(k_t q^m S^n - \tau_c)^p \quad (5)$$

$$L = \xi q \quad (6)$$

Margaux Mouchene 9/1/y 15:49

Supprimé: -

Margaux Mouchene 9/1/y 15:49

Supprimé: a

Margaux Mouchene 9/1/y 15:49

Supprimé: river

Margaux Mouchene 9/1/y 15:50

Supprimé: a

Margaux Mouchene 9/1/y 15:49

Supprimé:

Margaux Mouchene 9/1/y 15:50

Mis en forme: Police :Italique

where K is an erodibility coefficient, q is water discharge per unit flow width on the cell, S is slope and the exponents m, n and p are positive. k_t is the shear stress parameter so that $k_t q^m S^n = \tau$ (shear stress) and Equation 5 takes the classic form of the excess shear-stress formula (Tucker, 2004). τ_c is the critical shear stress required for clast detachment. p is set to 1 in our experiments (following Lavé and Avouac, 2001). The transport length L depends on particle size and density (included in the coefficient ξ). This law implies that the deposition rate decreases when the water discharge per unit width q increases.

For fluvial processes, the flow width w can be set to the cell width dx or to a river width such as:

$$w = k_w Q^{0.5} \quad (7)$$

where k_w is a coefficient depending on the lithology and Q is the total water discharge at a river section. Flow-width variation is critical in the modelling of alluvial fan evolution because it plays a role in avulsion processes, in the changing flow dynamics (a change in flux geometry may lead to overflowing and a shift to distributive flow), and in incision patterns (leaving alluvial terraces in some cases).

3.4 Cover effect

Erosion of sediment is different from that of bedrock (Equations 3 and 5) and, within the bedrock, different layers can be defined by their respective erodibility and detachment or slope thresholds (κ and S_c for hillslope processes and K for fluvial processes). During a time step dt , different layers can be eroded on a given cell: the erosion of each layer consumes part of dt so that less time remains to erode the underlying layer. This time reduction is taken into account by multiplying dt by $(1 - \frac{\text{volume layer}}{w dx \epsilon dt})$ between layers. In this way, the “cover effect” of a sediment layer covering the bedrock (e.g Whipple and Tucker, 2002; Lague, 2010) can be taken into account.

3.5 Lateral erosion

Flowing water can erode lateral cells, which are topographically above them and placed in a lateral direction perpendicular to each downstream direction. The lateral sediment flux Q_{sl} is defined as a fraction of the flux in the considered direction (e.g. Murray and Paola, 1997; Nicholas and Quine, 2007):

$$Q_{sl} = \alpha Q_s \quad (8)$$

where α is a bank erodibility coefficient; it is specified for sediment and implicitly determined for bedrock layers, proportionally to their erodibility (i.e. $\alpha_{\text{sediment}}/\alpha_{\text{bedrock}} = K_{\text{sediment}}/K_{\text{bedrock}}$ with K from Equations 5 and 7).

Margaux Mouchene 9/1/y 15:51
Mis en forme: Police :Italique

Margaux Mouchene 9/1/y 15:51
Supprimé: to be reached

Margaux Mouchene 9/1/y 15:52
Supprimé: a positive coefficient,

Margaux Mouchene 9/1/y 15:52
Mis en forme: Police :Italique

Margaux Mouchene 9/1/y 15:52
Supprimé:

Margaux Mouchene 9/1/y 15:52
Supprimé: ing

Margaux Mouchene 9/1/y 15:52
Supprimé: ,

4 Model setup

The model simulates the evolution of a 100×150 km region split into a foreland zone (100×100 km) and an uplifting mountain zone (100×50km; Fig. 2). The grid cell size is 500×500 m. The dimensions are chosen to allow megafan building on an area matching that of the Lannemezan megafan and to permit competing catchments to develop during the drainage network growth phase; they correspond to a compromise between computing time and spatial resolution. Our model has much larger dimensions than previous experiments on coupled catchment-foreland systems (Tucker, 2004; Nicholas et al., 2009; Pepin et al., 2010; Langston et al., 2015) and the foreland/mountain width ratio is much higher than in previous work (Pepin et al., 2010). The initial surface is a horizontal grid with a Gaussian elevation noise ($\sigma=0.5$ m) so we can study the system dynamics from the start of drainage network growth.

Although the convergence stopped in the early Miocene, the Pyrenees are still a high relief mountain range with long-term exhumation rates and present-day uplift rates both on the order of 0.1-0.3 mm y⁻¹ (e.g. Jolivet et al., 2007; Nguyen et al., 2016), possibly controlled by the ongoing isostatic response to post-orogenic erosion. As isostasy is currently not included in the model, we impose block uplift of the mountain part of the model at a constant rate of 0.3 mm y⁻¹ (except in experiments where this is explicitly modified, see below). Homogeneous precipitation is applied at a constant rate over the entire model ($P = 1$ m y⁻¹); this parameter is modified in some experiments (Exp. 2a, b, c).

The sides of the mountain block (southern border and southern third of the eastern and western borders) are closed; neither water nor sediment can exit the grid through these. The other boundaries are open, corresponding to transverse rivers of fixed elevation (0 m) and able to transport both sediment and water fluxes out of the grid. This is an approximation, as these transverse rivers will obviously have a downstream slope and therefore not be at a fixed constant elevation. However, the major rivers of the Pyrenean foreland have very low gradient (<0.5 %) and lie significantly lower than the streams incising the megafan.

We conducted a series of trial runs to adjust the relevant parameters in order to reproduce the first-order morphological traits of the northern Pyrenean foreland. In particular, the values for transport length (L), for the erodibilities of bedrock and sediments (respectively k_{br} and k_{all}) and for the critical shear stress (τ_c) need to be established. These parameters are critical in the relief evolution but are generally poorly constrained. Giachetta et al. (2015) provided a compilation of values for erodibilities of a set of lithologies in the context of the Iberian peninsula. However, these were used for models where the critical shear stress is zero and should be significantly different (approximately one to two orders of magnitude larger) when $\tau_c > 0$. The erodibility coefficient also depends on the value of m (e.g., Carretier et al., 2009), we therefore used a different value than that of Giachetta et al. (2015). In any case, the erodibility of sediment should be larger than that of the bedrock, the ratio between the two critically influencing the landscape morphology. We thus tested this ratio and the transport length L in order to reproduce the first-order characteristics of the northern Pyrenean landscape. Resemblance between the model and the ASTER DEM of the Lannemezan area was evaluated based on a number of morphological parameters (which need

Margaux Mouchene 9/1/y 15:53

Supprimé: -by-

Margaux Mouchene 9/1/y 15:53

Supprimé: x

Margaux Mouchene 9/1/y 15:53

Supprimé: x

Margaux Mouchene 9/1/y 15:53

Supprimé: x

Margaux Mouchene 7/1/y 11:01

Supprimé: The mountain part of the model is uplifted as a block at a constant rate of 0.3 mm y⁻¹

Margaux Mouchene 7/1/y 11:10

Supprimé: , which seems reasonable for the late stages of evolution of the central Pyrenees (Jolivet et al., 2007; Nguyen et al., in review)

Margaux Mouchene 9/1/y 15:58

Supprimé: Also,

Margaux Mouchene 9/1/y 15:58

Supprimé: t

Margaux Mouchene 9/1/y 15:58

Supprimé: and

Margaux Mouchene 9/1/y 15:58

Supprimé: sediment

to agree within 30% to be accepted); in the range and foreland we assessed maximum, minimum and mean elevations, river spacing and relief (i.e. valley-to-ridge elevation difference); in the foreland we also assessed the length, width and northward slope of the megafan. The duration of the megafan-building and -incision phases are also compared to the evolution of the Pyrenean foreland as described by previous authors (Crouzel 1957; Azambre et al., 1989; Mouchéné, et al., in press). The best-fit parameters used for the model runs are presented in Table 1. Pepin et al. (2010) suggested that the critical shear stress should be significant for permanent incision to occur. We thus fixed a positive value for τ_c (15 Pa) following Lavé and Avouac (2001) and Pepin et al. (2010).

5 Results

5.1 Megafan building

We successfully reproduced the first-order morphology of a fluvial megafan constructed on a low-elevation, stable foreland, from the erosional products of a slowly uplifting mountain-range-like block (Fig. 3). The drainage network initiates from the area of transition between the mountain and foreland blocks (Fig. 3 A). In the foreland, it propagates outward and fans aggrade. Evenly spaced rivers (every ~10 km) build small fans and progressively lengthen their watershed towards hinterland through headward incision. The fans quickly merge into a bajada, on top of which the flow is distributive (Fig. 3 B). At around 7.65 My, the mountain range becomes fully connected (i.e. all cells of the mountain block are connected to the base level through the river network) and the mountain outflux is dominated by a few large rivers (~5). In the meantime, aggradation continues in the foreland with a markedly conical pattern. The rivers situated at the easternmost and westernmost ends of the mountain range bend sharply to follow an along-strike course and quickly reach the open model boundaries, constrained by their short distance to a base-level outlet. In the following time steps, their watershed will increase in size by retreat of the drainage divide towards the middle of the range and the mountainous outlets of these streams will migrate towards the nearest border (Fig. 3 C). At this stage, the foreland deposits are mainly provided by a single central channel, the flow of which distributes sediments largely over the whole foreland, now clearly defining a megafan (the flow spanning 180° over the foreland).

Several episodes of temporary entrenchment (< 50m) occur during the building phase. They either concern the lower parts of the fan being incised by headward incision (Fig. 3 D) or the apex being incised by the main stream (Fig. 3 E). In both cases, within a few hundred thousand years the main stream has brought sufficient material to the entrenched zone to refill it and to overflow and become distributive again (Fig. 3 D and D'). This cyclic pattern is expected on megafans (Leier et al., 2005) and shows that the code mimics the natural fluvial dynamics of these settings.

On the long term, the mean elevation stabilizes in both the foreland and the mountain. From about 9 My, mean elevation stabilizes in the range (Fig. 3 B) but the elevation change remains slightly positive, which means that the relief is eroded at a slower rate than the applied uplift rate (i.e. true topographic steady state is not reached). Aggradation continues in the foreland, although at slow pace (0.015 to 0.02 mm y⁻¹ of mean elevation change), the timescale of aggradation is thus larger

Margaux Mouchene 7/1/y 14:11
Supprimé: minimum, maximum and mean elevations of the range and foreland megafan, relief and drainage network pattern.

Margaux Mouchene 9/1/y 16:00
Supprimé: -

Margaux Mouchene 9/1/y 16:01
Supprimé: -

Margaux Mouchene 9/1/y 16:01
Supprimé: towards the north

Margaux Mouchene 9/1/y 16:01
Supprimé: the south

Margaux Mouchene 9/1/y 16:01
Supprimé: a

Margaux Mouchene 9/1/y 16:02
Supprimé: probably

Margaux Mouchene 9/1/y 16:02
Supprimé: Meanwhile

Margaux Mouchene 9/1/y 16:02
Supprimé:

Margaux Mouchene 9/1/y 16:02
Supprimé: a

Margaux Mouchene 9/1/y 16:03
Supprimé:

Margaux Mouchene 9/1/y 16:03
Supprimé: higher

than that of the relief development. This is consistent with the observation of Babault et al. (2005) from an analog model. For them, aggradation in the foreland influences erosion of the mountain range by modifying the relative uplift rate (i.e. the difference between the uplift applied to the mountain block and the aggradation rate). Erosion of the range balances the continuously varying relative uplift rate, creating a “dynamic equilibrium” (Babault et al., 2005). A steady-state equilibrium (in which erosion rate equals uplift rate in the mountain) cannot be reached in such a landscape as long as aggradation occurs in the foreland.

5.2 Autogenic entrenchment

If the same conditions are maintained, natural entrenchment of the main stream occurs rapidly over a timescale that is two orders of magnitude smaller than the fan-building timescale (Fig. 4). Contrary to the building phase, during which episodes of temporary entrenchment occurred but were followed by refilling and overflow, the incision starting at 15.3 My near the apex is sufficient to constrain the main stream avulsions to the eastern half of the fan for the subsequent time steps (Fig. 4 G, H, I). A small stream that developed on the eastern foot of the fan episodically captures the main flow and is thus progressively deepened and incised through headward incision.

Still, the main flow remains highly distributive and overflows several times this path, before being finally permanently captured at around 15.57 My (Fig. 4 J). This event triggers a rapid incision phase, reaching nearly 150 m of incision close to the apex, and larger amounts further downstream. The mean elevation change in the foreland drops dramatically upon entrenchment and the fan is subsequently being eroded (mean elevation change remains negative for the rest of the experiment, Fig. 4). Erosion sharply increases in the mountain, especially in the watershed of the now connected main stream (Fig. 4). Figure 5 suggests that this incision leads to an increase of the relief due to rapid incision in the riverbed and little to no increased erosion on the hillslopes and ridges.

The main stream then erodes laterally its right bank in the foreland, tending towards an along-strike flow direction without further vertical incision (Fig. 4 K, L). Incision occurs in this bank and oblique to it, eventually capturing a secondary stream of the mountain range.

5.3 External forcing

Subsequent experiments start from the topography obtained at the end of the “building phase” at 15.3 My and aim at evaluating the respective roles of different external factors on the incision pattern of the megafan. We consecutively explore the influence of changing parameters related to climate (precipitation rate and frequency of precipitation events), base level change and tectonics (uplift rate and style). These models are run for 500 ky to evidence the effects of external factors at this specific timescale, which corresponds to the abandonment and incision timescale of both the model and the Lannemezan megafan. Parameters used for these experiments are summarized in Table 1.

Margaux Mouchene 9/1/y 16:03

Supprimé: the

Margaux Mouchene 9/1/y 16:03

Supprimé: this

Margaux Mouchene 9/1/y 16:04

Supprimé: ;

Margaux Mouchene 9/1/y 16:04

Supprimé: a

Margaux Mouchene 9/1/y 16:04

Supprimé: thalweg

Margaux Mouchene 9/1/y 16:04

Supprimé: a

Margaux Mouchene 9/1/y 16:04

Supprimé: ;

Margaux Mouchene 9/1/y 16:04

Supprimé: which

Margaux Mouchene 9/1/y 16:04

Supprimé: es

Margaux Mouchene 9/1/y 16:04

Supprimé: a

Margaux Mouchene 9/1/y 16:05

Supprimé: northern Pyrenean foreland

Margaux Mouchene 9/1/y 16:05

Supprimé: subsequently

Margaux Mouchene 9/1/y 16:05

Supprimé: the

5.3.1 Precipitation rate and style

Decreasing (Experiment 2a1) or increasing (Experiment 2a2) the precipitation rate only results in decelerating or accelerating the processes observed in the original experiment. The same evolution is observed in experiment 2a1 as using the default settings, but the evolution is slower and the model does not reach the permanent entrenchment stage after the 500 ky simulation. In the experiment with increased precipitation rate (2a2), erosion is enhanced and results in important widening of the valleys, but scattered deposition in the lower valleys create instabilities that perturb the model results.

In experiment 2b, we set the precipitation rate to follow a sinusoidal distribution with 100 ky cycles, to simulate the Quaternary climatic cycles. In this case, the trends of mean elevation change in the mountain and in the foreland are inversely correlated (Fig. 6). The mean elevation of the foreland slightly increases through the experiment but remains stable in periods of low runoff, as the sediment supply from the mountain is halted. The mountain is eroded in periods of maximum runoff, whereas the elevation increases (at the uplift rate) in periods of minimum runoff. There is a slight delay in the mountain response to the variations in precipitation: as the runoff starts to increase, the elevation in the mountain continues to rise at the uplift rate for another time step (10 ky) before it starts to decrease (Fig. 6). Similarly there is a small lag between maximum runoff and minimum mean elevation change (Fig. 6). This delay corresponds to the response time of the mountain to cyclic precipitation rate changes and is consistent with works by Carretier and Lucazeau (2005) and Braun et al. (2015), who suggested 1 to 30 ky offset between forcing and response to rainfall variability at orbital (Milankovitch) timescales. In our experiment, the same delay is observed in the foreland, although the signal is less clear for periods of high runoff (Fig. 6).

At the end of the 500-ky simulation, no permanent entrenchment is observed on the megafan. The small amounts of incision that occur on the fan when precipitation decreases are more than compensated by renewed sediment influx from the mountain as precipitations start to increase again. The incision of the riverbed in the mountain in periods of high runoff is more than compensated for by the uplift in periods of low runoff (dominated by the applied uplift; Fig. 6)

5.3.2 Base-level change

A 50-m drop in base level is applied at the beginning of Experiment 3a. This leads to erosion in the foreland through headward incision of a number of streams, developing mostly on the western and eastern borders and persisting until the end of the experiment. Connection between the main feeder channel and the largest incising stream on the eastern border happens earlier than in the default model (at around 250 ky) but the subsequent landscape evolution is very similar in both cases, although more incised streams remain at the end of this experiment (Fig. 7).

5.3.3 Uplift rate

Experiment 4a tests a scenario where uplift stops after the megafan building phase. In this experiment, the headward-incising stream connected to the eastern border captures a secondary river (at 300 ky) before connecting to the main central channel

Margaux Mouchene 9/1/y 16:06

Supprimé: w

Margaux Mouchene 9/1/y 16:06

Supprimé: experiment 2b

Margaux Mouchene 9/1/y 16:07

Supprimé:

Margaux Mouchene 9/1/y 16:08

Supprimé: periods

Margaux Mouchene 9/1/y 16:08

Supprimé: Overall, during more humid periods, the mountain is eroded and the sediments are transported to the foreland, which provides material for fan aggradation, whereas in periods of low precipitation, the material is not eroded and/or transported from the mountain to the piedmont (Fig. 6).

Margaux Mouchene 9/1/y 16:08

Supprimé: e

Margaux Mouchene 9/1/y 16:09

Supprimé: thalwegs

Margaux Mouchene 9/1/y 16:09

Supprimé: stream

Margaux Mouchene 9/1/y 16:09

Supprimé: thalweg

Margaux Mouchene 9/1/y 16:09

Supprimé: thalwegs

Margaux Mouchene 9/1/y 16:09

Supprimé: thalweg

Margaux Mouchene 9/1/y 16:10

Supprimé: is continuously incised and connects with the outlet of

at the end of the experiment (500 ky, Fig. 8). The mean erosion rate in the range decreases steadily down to 0.19 mm y⁻¹ (value for last 10 ky of the experiment).

Increasing the uplift rate to 1 mm y⁻¹ (Experiment 4c) quickly and permanently increases the elevation in the range without increasing the aggradation in the foreland significantly. This may be due to erosion (detachment and/or transport) not responding rapidly enough to catch up with this increase. Permanent entrenchment occurs at the end of the experiment (500 ky) through the same process as in the default experiment.

5.3.4 Tilting experiment

In Experiment 5, we seek to reproduce the effect of isostatic rebound on the erosional pattern of the range and its foreland. At the moment, the CIDRE model does not include flexure. We thus chose to simulate the first-order effect of the flexural response to erosional unloading of the range through simple linear tilting of the model. This corresponds to an uplift pattern that increases linearly from 0 at the north boundary to a maximum fixed value at the south boundary.

To scale the tilting to the observed geomorphic characteristics of the northern Pyrenean foreland, we estimate the potential tilting of the Lannemezan megafan since the onset of incision. We use a scaling law between fan area and fan slope to estimate the initial depositional slope of the Lannemezan Formation that caps the Miocene deposits. We use this formation because its base is the only mapped surface effectively preserved from erosion since deposition. We use a digitized geological map and an ASTER DEM (70 m resolution) to extrapolate the basal surface of the Lannemezan Formation using ArcGIS software and estimate its current slope at 0.5°. Figure 9A shows area and slope data for the Lannemezan megafan compared to data compilations from active and inactive alluvial fans and megafans from the Alps, the Andes and the Himalaya (Horton and DeCelles, 2002; Guzzetti et al., 1997). The discrepancy of the Lannemezan data with the scaling law suggests an estimated ~0.4° tilt, which will be simulated by uplift increasing linearly from 0 at the northern boundary to 2 mm y⁻¹ at the southern boundary of the model.

It should be noted that this scaling relationship suggests a depositional angle of ~0.1°, which is within the range of values for megafans (e.g. DeCelles and Cavazza, 1999) but is not consistent with the slope observed in the default experiment (~0.4-0.5°). We compare this result with the tilt estimated using another oft-used scaling relationship, between the catchment area and the fan slope (e.g. Champagnac et al., 2008). Figure 9B shows area and slope data for the Lannemezan megafan compared to data compilations from active and inactive alluvial fans from the Alps (Guzzetti et al., 1997; Crosta and Frattini, 2004; Champagnac et al., 2008). The discrepancy of the Lannemezan data with this scaling law only suggests an estimated 0.13° tilt, which will be simulated by uplift increasing linearly from 0 at the north boundary to 0.68 mm y⁻¹ at the southern boundary. We test both these minimum (0.13°, Experiment 5a) and maximum (0.4°, Experiment 5b) tilt scenarios.

With the linearly increasing uplift, the megafan continues to grow; the mean elevation change in the foreland is steady, positive and higher than in the default experiment (>0.2 mm y⁻¹, including uplift). In both experiments, connexion with the headward incising stream and entrenchment occur (at ~280ky in the lower tilt experiment, and at ~240ky in the higher tilt experiment) but only temporarily affects this trend because, as tilting continues, the river outflows from this path (at ~320 ky

Margaux Mouchene 9/1/y 16:10

Supprimé: much

Margaux Mouchene 9/1/y 16:11

Supprimé: this experiment (

Margaux Mouchene 9/1/y 16:11

Supprimé:)

Margaux Mouchene 9/1/y 16:11

Supprimé: a

Margaux Mouchene 9/1/y 16:11

Supprimé: e

Margaux Mouchene 9/1/y 16:12

Supprimé:

Margaux Mouchene 9/1/y 16:12

Supprimé: not surprising

Margaux Mouchene 9/1/y 16:12

Supprimé: a

Margaux Mouchene 9/1/y 16:12

Supprimé: -

Margaux Mouchene 9/1/y 16:12

Supprimé: estimations

Margaux Mouchene 9/1/y 16:12

Supprimé:

Margaux Mouchene 9/1/y 16:13

Supprimé: thalweg

Margaux Mouchene 9/1/y 16:13

Supprimé: that

Margaux Mouchene 9/1/y 16:13

Supprimé: the

and ~260 ky respectively; Fig. 10). Deposition in the main path causes the stream to overflow from this channel and resume distributive flow over the megafan, but instabilities in the models blur the results (Fig. 10). This suggests that, overall, tilting of the model prevents or limits permanent entrenchment. The mean elevation change in the mountain is steady and positive (~0.25 mm y⁻¹ and 1.2-1.3 mm y⁻¹ respectively), with peaks following the transient capture.

5 Figure 11 compares the evolution of a north-south topographic profile across Experiment 5a (0.13° tilt) and the default model. The megafan topography on this section is rather stable over the course of the default experiment. However, the megafan slope increases significantly in experiment 5a, showing that the tilt affects the megafan slope without being fully compensated by erosion.

6 Discussion

10 6.1 Megafan building

In the model, the main steps of the megafan building phases are: (i) foreland deposition starts with small fans that quickly merge into a *bajada*; mountain watersheds merge so only a few streams are left; (ii) the rivers situated near the boundary change their direction to reach the shortest flow path to the border and the central stream becomes the dominant provider for foreland sedimentation; and (iii) the megafan grows in response to cyclic flow dynamics (oscillating between channelized and distributive flow) and reaches dynamic equilibrium.

15 The timescale of the building phase of the megafan is long (>10 My) when compared to active megafans of similar volumes deposited during the Quaternary (e.g. in the Alps, Andes and Himalaya; Assine et al., 2014; Fontana et al., 2014; Abrahami, 2015) but compares well to older systems (e.g. Campanian-Maastrichtian Hams Fork formation in Utah; DeCelles and Cavazza, 1999) and is consistent with the Lannemezan megafan building phase encompassing the early-middle Miocene to Pliocene (i.e. ~15 My).

20 The long foreland (foreland length/mountain length = 2) allows for a large fan to develop but requires the model parameters to be set in a way that allows transportation over such great distance, in particular the parameter *L* must be large enough. This required longer *L*, which may be interpreted as a smaller settling rate (Davy and Lague, 2009), is consistent with the downstream fining of sediment in the Lannemezan megafan (Crouzel 1957; Azambre et al., 1989), even though sediment fining is not accounted for in CIDRE.

25 The boundary conditions, open in the foreland and closed in the mountain, play a key role in the development and evolution of the drainage network. In particular, open boundary conditions on all three sides of the foreland allow for (i) the central river to become dominant in the sediment flux deposited in the foreland (thus creating a megafan) as the more lateral rivers rapidly adapt their course to the shortest path reaching the base-level (along-strike to reach boundaries), and (ii) the conical shape to develop (contrary to Pepin et al., 2010, where cyclic boundary condition on lateral boundaries resulted in a more *bajada*-like landform). Open boundary conditions in the mountain would result in strike-parallel drainage, which shows that megafan building requires a relatively large range. Thus, in natural settings, transverse rivers with efficient fluvial transport

Margaux Mouchene 9/1/y 16:14

Déplacé (insertion) [1]

Margaux Mouchene 9/1/y 16:14

Déplacé vers le haut [1]: Deposition in the main path causes the stream to overflow from this channel and resume distributive flow over the megafan, but instabilities in the models blur the results (Fig. 10). This suggests that, overall, tilting of the model prevents or limits permanent entrenchment.

Margaux Mouchene 9/1/y 16:14

Supprimé: e

Margaux Mouchene 9/1/y 16:14

Supprimé: run

Margaux Mouchene 9/1/y 16:15

Supprimé:

Margaux Mouchene 9/1/y 16:15

Supprimé: thanks

Margaux Mouchene 9/1/y 16:15

Supprimé: (

Margaux Mouchene 9/1/y 16:15

Supprimé: E

Margaux Mouchene 9/1/y 16:15

Supprimé: M

Margaux Mouchene 9/1/y 16:15

Supprimé: Early

Margaux Mouchene 9/1/y 16:16

Mis en forme: Police :Italique

Margaux Mouchene 9/1/y 16:16

Supprimé: . S

Margaux Mouchene 9/1/y 16:16

Supprimé: Also, o

(to evacuate both water and sediments) appear necessary on all sides for a river/fan system to be singled out and grow into a megafan deposit. In the northern Pyrenean foreland, the Garonne/Ariège and Adour rivers could have played this role, which suggests that they might have existed prior to the Miocene onset of the megafan building.

In our model, the absence of subsidence in the foreland may have encouraged the development of a fan covering a large area, which imposes overfilled conditions in the foreland basin. High subsidence rate would have allowed thick accumulation close to the range and thus limited its northward extension (e.g. Allen et al., 2013). This hypothesis could be tested with the addition of an algorithm for flexure (Simpson, 2006; Naylor and Sinclair, 2008). Nevertheless, the overfill hypothesis may be justified by the deceleration of subsidence rates in the Pyrenean retro-foreland since the Eocene (Desegaulx and Brunet, 1990; Desegaulx et al., 1990). In any case, the impact of varying subsidence rate on megafan growth and abandonment remains to be evaluated (e.g. Dingle et al., 2016).

6.2 Autogenic incision

6.2.1 Time and space scales

The autogenic entrenchment happens around 15.57 My, which, within the framework of the Lannemezan megafan evolution, is consistent with the fan-building phase encompassing the early-middle Miocene to Pliocene and incision taking place during the Quaternary (since 300 ka; Mouchéné et al., in press).

At 15.3 My (end of the “building phase”), the mean elevation in the mountain range is about 1460 m, with a maximum elevation at 3160 m near the southern border, which is consistent with the northern flank of the Pyrenees. In the foreland, a maximum elevation of 950 m is reached at the fan apex in the model, which is higher than the current elevation of the Lannemezan megafan apex (~660 m), but the mean elevation of the foreland is around 270 m, comparable to that of the Lannemezan megafan. Megafan shape and dimensions (area) agree between our model and the Lannemezan megafan. At this point, the watershed of the main feeding river is about 1100 km², which is quite large when compared to the current Neste watershed (~750 km²). The scale of the vertical entrenchment of the river is similar in the model and in the Lannemezan case (~100-150 m near the apex)

6.2.2 Mechanism/necessary conditions

For Pepin et al. (2010), autogenic entrenchment of an alluvial fan occurs only if (i) progradation is limited by the open boundary with fixed elevation and (ii) the transport threshold (critical shear stress) is significant. Nicholas et al. (2009) also suggested that declining aggradation in the fan results from increasing fan area during progradation (building phase) and incision is triggered by the lack of accommodation space when boundary conditions are reached. In nature, some incised fans are linked to powerful transverse rivers (Milana and Ruzycki, 1999; Dühnforth et al., 2007; 2008) but the causality is not proven and external forcing is demonstrated in some cases (e.g. Dühnforth et al., 2008).

Margaux Mouchene 9/1/y 16:16

Supprimé: of

Margaux Mouchene 9/1/y 16:17

Supprimé: E

Margaux Mouchene 9/1/y 16:17

Supprimé: M

Margaux Mouchene 9/1/y 16:17

Supprimé: Early

Margaux Mouchene 9/1/y 16:17

Supprimé: submitted

Margaux Mouchene 9/1/y 16:18

Supprimé:

Margaux Mouchene 9/1/y 16:18

Supprimé:

Margaux Mouchene 9/1/y 16:18

Mis en forme: Exosant

Margaux Mouchene 9/1/y 16:18

Supprimé:

Margaux Mouchene 9/1/y 16:18

Supprimé:

Margaux Mouchene 9/1/y 16:18

Mis en forme: Exosant

Margaux Mouchene 9/1/y 16:19

Supprimé: (

Margaux Mouchene 9/1/y 16:19

Supprimé: ;

In our model, entrenchment naturally occurs, but it happens long after the moment where sediments reach the model boundaries (Fig. 3 and 4). This is different from Pepin et al (2010) since, in their experiments, autogenic entrenchment occurred precisely when the sediment reached the free border. We suspect that this difference comes from the lateral erosion included in our modelling and absent in the simulations of Pepin et al. (2010). Lateral erosion limits the incision by fostering lateral migration and channel widening. Although our modelling seems more realistic, the comparison of this prediction with natural settings is not straightforward because boundary conditions are likely to change over time (e.g. Harvey, 2002). Consistently with the findings of Pepin et al. (2010) and Nicholas and Quine (2007), all our models that predict autogenic entrenchment use a significant critical shear stress (entrainment threshold). We suspect this threshold to control part of the incision magnitude and the delay between the moment when sediment reach the free border and the moment when incision occurs. This aspect should be further evaluated by varying the critical shear stress in other experiments. Van Dijk et al. (2009) proposed that aggradation on fans allows a critical slope to be reached, triggering the incision. However, in the tilting experiments (Experiment 5), the fan slope reaches greater values than in the default experiment at the time of entrenchment, but permanent entrenchment does not occur. Therefore, attainment of a critical slope does not appear to constitute a threshold for entrenchment. One explanation may be that tilting fosters erosion in the mountain, with a larger incoming sediment discharge entering the foreland, which prevents incision from growing.

6.2.3 Incision pattern

In the model, permanent entrenchment results from (limited) incision near the apex by the feeding river and major headward incision of a stream from the foot of the fan until both ends meet to define a continuously entrenched pathway (Fig. 4). In the case of the Lannemezan megafan, we cannot provide evidence to support or disprove this mechanism but the drainage pattern at the apex of the Lannemezan megafan resembles the model (Fig. 12 A and B). We could therefore envisage the following scenario:

- A pre-existing river Ariège/lower Garonne River flowing through the foreland to the east of the megafan, while the Neste river feeds the Lannemezan megafan through a distributive deposition pattern (Fig. 12 C);
- A tributary of the ancestral Ariège/Garonne river retreats headward toward the apex of the megafan (Fig. 12 D);
- Headward incision of the tributary toward the west leads to sequential capture of (a) the upper Garonne and (b) the upper Neste, abandonment of the fan, rapid incision and terrace formation (preferably on the left bank of the now merged Neste/Garonne River due to southward river migration; Fig. 12 E).

The amount of incision is already very important in the first time step following the connection (~100 m near the apex) and will only be further increased by another 80 m near the apex. Downstream, the stream erodes its right bank towards a strike-parallel pathway but does not incise vertically (Fig. 4). This last characteristic resembles the lateral migration of the Neste and Garonne rivers during their incision, evidenced by the extensive alluvial terrace staircase left almost systematically on their left banks. In our model, the sediments of the channel bed could enhance the effect of lateral incision and inhibit further vertical incision through their cover effect (see also Hancock and Anderson, 2002; Brocard and van der Beek, 2006).

Margaux Mouchene 9/1/y 16:19
Supprimé:)... since, il... their experim... [2]

Margaux Mouchene 9/1/y 16:21
Supprimé: 's...or...nd Nicholas and Quin... [3]

Margaux Mouchene 9/1/y 16:23
Supprimé: for ... critical slope to be reach... [4]

Margaux Mouchene 9/1/y 16:24
Supprimé: thalweg

Margaux Mouchene 9/1/y 16:25
Supprimé: ...Neste river feedsing... [5]

Margaux Mouchene 9/1/y 16:26
Supprimé: s...toward the west (toward... [6]

Margaux Mouchene 9/1/y 16:27
Supprimé: migration...of the tributary t... [7]

Margaux Mouchene 9/1/y 16:28
Supprimé: x...on (~100 m near the apex) <... [8]

However, the terraces of the northern Pyrenean foreland prove that [the ~100-m incision of the Lannemezan megafan at its apex](#) was episodic, which contrasts with the pulse of incision predicted by the model. Cosmogenic nuclide surface exposure dating suggests that these incision episodes in the northern Pyrenean foreland are linked to cold-to-warm climatic transitions (Mouchené et al., [in press](#)).

6.3 Impact of climate change

In the model with sinusoidal precipitation rates (Experiment 2b), humid periods are characterized by erosion in the mountains and deposition in the foreland (with episodic incision); both decrease in drier periods because stream power decreases and less material is being transported from the mountains. The wet-to-dry transition corresponds to a decrease in sediment input but also to a decrease in fluvial efficiency as the runoff nears 0, which prevents incision.

In the northern Pyrenean foreland, incision and abandonment of alluvial terraces has been linked to cold-to-warm climatic transitions (Mouchené et al., [in press](#)) where the rapid decrease in sediment flux and gradual transitioning of the river to a single meandering thread, with a low width/depth ratio, would encourage vertical incision (e.g. Hancock and Anderson, 2002). Warm-to-cold transitions can also be associated with incision because of the increase in runoff variability and decline in vegetation that characterizes these periods, but in nature, they are usually more gradual than cold-to-warm transitions. During glacial (dry, cold) periods, regolith is actively produced on hillslopes by efficient frost cracking but it is mobilized only at the onset of the following interglacial (wetter) period, when rainfall increases (e.g. Carretier et al., 1998). To reproduce [and further explore this effect](#), we would need to include a climate (temperature)-dependant law for [sediment production in the model](#).

In nature, incision is not always related to the return of wetter conditions; Meyer et al. (1995) suggest that incision of the terraces in their study site in northwestern Yellowstone National Park happens during warmer, more drought-prone periods because of the infrequent floods scouring the channel bed. Langston et al. (2015) recently modelled a similar pattern of incision by applying more intense, longer duration precipitation events during interglacial periods, but without changing the average precipitation rate. Periglacial processes have also been suggested to be a key controlling factor for erosion (e.g. [Marshall et al., 2015](#); [Dosseto and Schaller, 2016](#)): erosion is enhanced during cold periods in regions where they occur; [whereas](#) it is enhanced during warmer periods in regions exempt of periglacial processes. Mass wasting processes could be the main driver for erosion increase during wet periods (e.g. [Bookhagen et al. 2005](#)) although their relationship to other environmental parameters, such as vegetation cover, remains disputed (e.g. [Istanbulluoglu and Bras, 2005](#); [Carretier et al., 2013](#); [Dosseto and Schaller, 2016](#)). Our current model does not take [such processes](#) into account. Aggradation and incision thus seem to be controlled by the variability in rainfall intensity and event duration but also by temperature-dependent hillslope processes, rather than by mean precipitation rate [alone](#).

[A number of studies have related](#) terrace incision with climate changes (e.g. [Barnard et al., 2006](#); [Bridgland and Westaway, 2008](#)). This also seems to be the case in the northern Pyrenean foreland, where terrace abandonment was related to

Margaux Mouchene 9/1/y 16:30

Supprimé: (100-m at the apex)

Margaux Mouchene 9/1/y 16:31

Supprimé:

Margaux Mouchene 9/1/y 16:30

Supprimé: submitted

Margaux Mouchene 9/1/y 16:31

Supprimé: ic

Margaux Mouchene 9/1/y 16:31

Supprimé: more

Margaux Mouchene 9/1/y 16:31

Supprimé: submitted

Margaux Mouchene 9/1/y 16:31

Supprimé: -

Margaux Mouchene 9/1/y 16:32

Supprimé: of sediment

Margaux Mouchene 9/1/y 16:32

Supprimé: Dühnforth et al. 2010;

Margaux Mouchene 9/1/y 16:39

Supprimé: b

Margaux Mouchene 9/1/y 16:32

Supprimé: such processes

Margaux Mouchene 9/1/y 16:33

Supprimé: In nature, a

Margaux Mouchene 9/1/y 16:33

Supprimé: ic

Quaternary climatic changes, although the model does not reproduce this pattern (it does not produce terraces at all). Several experiments suggest that the longer the foreland, the more it buffers the effects of short-period variations (Métivier and Gaudemer, 1999; Babault et al., 2005; Carretier and Lucazeau, 2005), so the effect of rapid climatic changes could be dampened by the large dimensions of the foreland in our model, preventing terrace formation. The lack of temperature-dependent processes in our experiments (glacial erosion, temperature-dependent regolith production) may also prevent terrace formation. Finally, the model resolution could be insufficient to resolve alluvial terraces.

6.4 Uplift rate

In the experiment where uplift stops after 15.3 My (Experiment 4a), the mountains erode at a rate of 0.19 mm y^{-1} , comparable to the highest values obtained through estimation of basin-averaged erosion rates using cosmogenic nuclides in river sands (0.01 to 0.16 mm y^{-1} ; Mouchen , 2016). Uplift is thought to have significantly decreased in the Pyrenees since the Miocene, with modern GPS-derived uplift rates being small ($0.1 \pm 0.2 \text{ mm y}^{-1}$ of differential uplift of the mountain belt with respect to a regional reference frame; Nguyen et al., 2016). Our results suggest that the Lannemezan megafan could have been built in a period of reduced tectonic uplift. The evolution of the piedmont is very similar to that of the default experiment (where uplift is maintained at 0.3 mm y^{-1}) except for the entrenchment that is refilled in experiment 4a. Thus, it appears that tectonic activity in the mountain belt does not strongly influence incision dynamics in the foreland.

6.5 Flexural isostatic rebound

We attempted to simulate the effect of flexural isostatic rebound on the incision pattern through tilting of the model. In the Alps, tilting of the foreland appears related to isostatic rebound in response to accelerated glacial erosion and possibly deep-seated geodynamic processes (Champagnac et al., 2008). This pattern has not been demonstrated for the Pyrenees. Although the simplistic approach we used does not reproduce the flexural response to erosional unloading of the range in detail, the slope of the fan topographic profile increases with time through this process, as suggested for alpine fans by Champagnac et al. (2008). Quantification of this increase in slope, although complicated by poor outcrop conditions, needs to be done in the northern Pyrenean piedmont to compare with the slope angles obtained in our model. In any case, in the experiment, tilting prevented permanent entrenchment so this mechanism cannot explain the abandonment of a foreland megafan.

In the model, the topographic profiles merge downstream as a consequence of tilting. The alluvial terraces along the northern Pyrenean rivers also merge downstream and this pattern is also observed in the Alpine foreland. However, this pattern does not necessarily relate to tilting of the megafan: in other settings, this characteristic has been interpreted as a climatic imprint on incision (Poisson and Avouac, 2004; Wobus et al., 2010; Pepin et al., 2013). Thus, tilting does not appear to play a major role in the abandonment of the Lannemezan megafan.

Margaux Mouchene 9/1/y 16:33

Supprim : the northern Pyrenees

Margaux Mouchene 9/1/y 16:34

Supprim : very

Margaux Mouchene 7/1/y 17:44

Supprim : ;

Margaux Mouchene 7/1/y 11:06

Supprim : in review

Margaux Mouchene 9/1/y 16:34

Supprim : is

Margaux Mouchene 9/1/y 16:35

Supprim : w

Margaux Mouchene 9/1/y 16:35

Supprim : (incision was only temporary)

Margaux Mouchene 7/1/y 17:46

Supprim : evolution

7. Conclusions

Numerical modelling of the evolution of a catchment/foreland system has provided (i) new insight in the building and incision of a foreland megafan and (ii) key elements to infer the driving forces in the natural evolution of the remarkable Lannemezan megafan and its mountainous catchment, in the northwestern Pyrenees.

5 For a megafan to develop, the foreland must be large enough to provide sufficient space for the fan to expand for a long period of time; a lack of subsidence may help this process. The role of pre-existing transverse rivers flowing across the foreland seems to be critical in the building and incision of the megafan. They rapidly capture the closest streams exiting the range, which allows for a central mountainous stream to be singled out and to provide for most of the foreland deposits stacked in the megafan. In the northern Pyrenean foreland, the through-going Adour and Garonne/Ariège Rivers may have helped shaping the Lannemezan megafan. the spacing of these pre-existing major drainage axes controls the size of the fan, they limit its extension and efficiently evacuate water and sediments out of the megafan. The megafan grows in response to the autogenic oscillations between sheet-flow and channelized flow. These oscillations trigger small incisions that are subsequently overfilled and rapid lateral movement of the flow over the whole fan surface.

15 Permanent entrenchment of the Lannemezan megafan could thus be the result of autogenic processes through (i) progressive headward incision of a stream from the foot of the fan (not too far from the apex) and (ii) final and rapid incision of the apex once this stream has captured the feeding river at its mountainous outlet. No external forcing is needed to induce long-term entrenchment on the order of magnitude observed in the field (100-m vertical incision near the apex) but external factors cannot be ruled out. In particular, on a shorter time-scale, incision may have been influenced by Quaternary climatic variations as suggested by the abandonment of terrace staircases along the foreland rivers incising the Lannemezan megafan.

20 Variations in precipitation rate alone do not appear to be sufficient to produce these episodic incision and alluviation phases and temperature-dependent hillslope processes may also be involved. In contrast, base-level changes, tectonic activity in the mountain range or tilting of the foreland through flexural isostatic rebound appear unimportant factors in the abandonment of the megafan.

25 Acknowledgments

This study was supported by French National Research Agency ANR (Project PYRAMID, ANR-11-BS56-0031) and forms part of MM's Ph.D. thesis funded by the French Ministry of Higher Education (MESR). ISTERre is part of Labex OSUG@2020 (ANR10 LABX56). We thank Hugh Sinclair and an anonymous referee for constructive comments that helped clarify the manuscript.

Margaux Mouchene 9/1/y 16:36

Supprimé: and

Margaux Mouchene 9/1/y 16:36

Supprimé: (

Margaux Mouchene 9/1/y 16:36

Supprimé: a

Margaux Mouchene 9/1/y 16:36

Supprimé:)

Margaux Mouchene 9/1/y 16:36

Supprimé: r

Margaux Mouchene 9/1/y 16:37

Supprimé: of

Margaux Mouchene 7/1/y 17:40

Supprimé: by clearing the sediment and water fluxes

Margaux Mouchene 9/1/y 16:37

Supprimé: thanks

Margaux Mouchene 9/1/y 16:37

Supprimé: thalweg

Margaux Mouchene 9/1/y 16:37

Supprimé: thalweg

Margaux Mouchene 9/1/y 16:37

Supprimé: seem

References

- Abrahami, R., 2015. Erosion et flux sédimentaires associés dans un bassin versant soumis à un régime de mousson : l'exemple de la Tista (Inde), du Sikkim au Mégafan. PhD Thesis, Université de Grenoble, 223pp.
- Abrams, M.J., Chadwick, O.H., 1994. Tectonic and climatic implications of alluvial fan sequences along the Batinah coast, Oman. *Journal of the Geological Society, London*, 151, 51–58.
- Allen, P.A., Armitage, J.J., Carter, A., Duller, R.A., Michael, N.A., Sinclair, H.D., Whitchurch, A.L., Whittaker, A.C., 2013. The Qs problem: Sediment volumetric balance of proximal foreland basin systems. *Sedimentology* 60, 102-130.
- Arboleya, M.-L., Babault, J., Owen, L.A., Teixell, A., Finkel, R.C., 2008. Timing and nature of Quaternary fluvial incision in the Ouarzazate foreland basin, Morocco. *Journal of the Geological Society, London*, 165, 1059–1073.
- Assine, M.L., Corradini, F.A., do Nascimento Pupim, F., McGlue, M.M., 2014. Channel arrangements and depositional styles in the São Lourenço fluvial megafan, Brazilian Pantanal wetland. *Sedimentary Geology* 301, 172–184.
- Azambre, B., Crouzel, F., Debroas, E.J., Soulé, J.C., Ternet, Y., 1989. Carte géologique de la France à 1/50000. Feuille 1053 Bagnères-de-Bigorre. Bureau des Recherches Géologiques et Minières, Orléans.
- Babault, J., Bonnet, S., Crave, A., Van Den Driessche, J., 2005. Influence of piedmont sedimentation on erosion dynamics of an uplifting landscape: An experimental approach. *Geology* 33, 301–304.
- Barnard, P.L., Owen, L.A., Finkel, R.C., 2006. Quaternary fans and terraces in the Khumbu Himal south of Mount Everest: their characteristics, age and formation. *Journal of the Geological Society* 163, 383–399.
- Biteau, J.J., Le Marrec, A., Le Vot, M., Masset, J.M., 2006. *The Aquitaine Basin. Petroleum Geoscience* 12, 247–273.
- Bookhagen, B., Thiede, R.C., Strecker, M.R., 2005. *Late Quaternary intensified monsoon phases control landscape evolution in the northwest Himalaya. Geology* 33, 149–152.
- Braun, J., Voisin, C., Gourlan, A.T., Chauvel, C., 2015. Erosional response of an actively uplifting mountain belt to cyclic rainfall variations. *Earth Surface Dynamics* 3, 1–14.
- Bridgland, D., Westaway, R., 2008. Climatically controlled river terrace staircases: A worldwide Quaternary phenomenon. *Geomorphology* 98, 285–315.
- Brocard, G.Y., van der Beek, P.A., 2006. Influence of incision rate, rock strength, and bedload supply on bedrock river gradients and valley-flat widths: Field-based evidence and calibrations from western Alpine rivers (southeast France). In: Willett, S.D., Hovius, N., Brandon, M.T., and Fisher, D.(Eds) *Tectonics, Climate, and Landscape Evolution*, Geological Society of America, pp.101-126, 2006, Geological Society of America Special Paper 398, doi: 10.1130/2006.2398(07)
- Carretier, S., Lucazeau, F., Ritz, J.-F., 1998. Approche numérique des interactions entre faille active, climat et érosion : exemple de la faille de Bogd, Gobi Altay, Mongolie. *Comptes Rendus de l'Académie des Sciences de Paris, Sciences de la terre et des planètes* 326, 391–397.
- Carretier, S., Lucazeau, F., 2005. How does alluvial sedimentation at range fronts modify the erosional dynamics of mountain catchments? *Basin Research* 17, 361–381.

Margaux Mouchene 9/1/y 16:38

Supprimé: (

Margaux Mouchene 9/1/y 16:38

Supprimé:)

- Carretier, S., Poisson, B., Vassallo, R., Pepin, E., Farias, M., 2009. Tectonic interpretation of transient stage erosion rates at different spatial scales in an uplifting block. *Journal of Geophysical Research* 114, F02003 doi: 10.1029/2008JF001080.
- Carretier, S., Regard, V., Vassallo, R., Aguilar, G., Martinod, J., Riquelme, R., Pepin, E., Charrier, R., Hérail, G., Farias, M., Guyot, J.-L., Vargas, G., Lagane, C., 2013. Slope and climate variability control of erosion in the Andes of central Chile. *Geology* 41 (2), 195–198.
- Carretier, S., Goddérés, Y., Delannoy, T., Rouby, D., 2014. Mean bedrock-to-saprolite conversion and erosion rates during mountain growth and decline. *Geomorphology* 209, 39–52.
- Carretier, S., Martinod, P., Reich, M., Goddérés, Y., 2015. Modelling sediment clasts transport during landscape evolution. *Earth Surface Dynamics* 3, 1221–1254.
- 10 Champagnac, J.-D., van der Beek, P., Diraison, G., Dauphin, S., 2008. Flexural isostatic response of the Alps to increased Quaternary erosion recorded by foreland basin remnants, SE France. *Terra Nova* 20, 213–220.
- Coulthard, T.J., Macklin, M.G., Kirkby, M.J., 2002. A cellular model of Holocene upland river basin and alluvial fan evolution. *Earth Surface Processes and Landforms* 27, 269–288.
- Crosta, G.B., Frattini, P., 2004. Controls on modern alluvial fan processes in the central Alps, northern Italy. *Earth Surface Processes and Landforms* 29, 267–293.
- 15 [Crouzel, F., 1957. Le Miocène continental du bassin d'Aquitaine. Bulletin du Service de la Carte Géologique de France 54, 264pp.](#)
- Dade, W.B., Verdeyen, M.E., 2007. Tectonic and climatic controls of alluvial-fan size and source-catchment relief. *Journal of the Geological Society, London* 164, 353–358.
- 20 Davy, P., Lague, D., 2009. The erosion/transport equation of landscape evolution models revisited. *Journal of Geophysical Research* 114, F03007, doi:10.1029/2008JF001146
- DeCelles, P.G., Cavazza, W., 1999. A comparison of fluvial megafans in the Cordilleran (Upper Cretaceous) and modern Himalayan foreland basin systems. *Geological Society of America Bulletin* 111, 1915–1334.
- 25 [Delmas, M., 2015. The last maximum ice extent and subsequent deglaciation of the Pyrenees: an overview of recent research, Cuadernos de Investigación Geográfica, 41 \(2\), 359-387.](#)
- Densmore, A.L., Allen, P.A., Simpson, G., 2007. Development and response of a coupled catchment fan system under changing tectonic and climatic forcing. *Journal of Geophysical Research*, 112, F01002, doi:10.1029/2006JF000474
- Desegaulx, P., Brunet, M.F., 1990. Tectonic subsidence of the Aquitaine Basin since Cretaceous times. *Bulletin de la Société Géologique de France* 8, (VI), 295–306.
- 30 Desegaulx, P., Roure, F., Villein, A., 1990. Structural evolution of the Pyrenees: tectonic inheritance and flexural behaviour in the continental crust. *Tectonophysics* 182, 211–225.
- DiBiase, R.A., Whipple, K.X., 2011. The influence of erosion thresholds and runoff variability on the relationships among topography, climate, and erosion rate. *Journal of Geophysical Research* 116, F04036 doi: 10.1029/2011JF002095.

Margaux Mouchene 9/1/y 16:40

Supprimé :

Margaux Mouchene 9/1/y 16:40

Supprimé :

Margaux Mouchene 9/1/y 16:40

Supprimé :

[Dingle, E.H., Sinclair, H.D., Attal, M., Milodowski, D.T., Singh, V., 2016. Subsidence control on river morphology and grain size in the Ganga Plain. *American Journal of Science* 316, 778–812.](#)

Dosseto, A., Schaller, M., 2016. The erosion response to Quaternary climate change quantified using uranium isotopes and in situ-produced cosmogenic nuclides. *Earth Science Reviews* 155, 60–81.

- 5 Dühnforth, M., Densmore, A.L., Ivy-Ochs, S., Allen, P.A., 2008. Controls on sediment evacuation from glacially modified and unmodified catchments in the eastern Sierra Nevada, California. *Earth Surface Processes and Landforms* 33, 1602–1613.
- Dühnforth, M., Densmore, A.L., Ivy-Ochs, S., Allen, P.A., Kubik, P.W., 2007. Timing and patterns of debris flow deposition on Shepherd and Symmes creek fans, Owens Valley, California, deduced from cosmogenic ¹⁰Be. *Journal of Geophysical Research* 112, F03S15 doi:10.1029–2006JF000562.
- 10 | Fontana, A., Mozzi, P., Marchetti, M., 2014. Alluvial fans and megafans along the southern side of the Alps. *Sedimentary Geology* 301, 150–171.
- Giachetta, E., Molin, P., Scotti, V.N., Faccenna, C., 2015. Plio-Quaternary uplift of the Iberian Chain (central–eastern Spain) from landscape evolution experiments and river profile modeling. *Geomorphology* 246, 48–67.
- Guerit, L., Métivier, F., Devauchelle, O., Lajeunesse, E., Barrier, L., 2014. Laboratory alluvial fans in one dimension. *Physical Review E* 90, 022203 doi: 10.1103–PhysRevE.90.022203.
- 15 Guzzetti, F., Marchetti, M., Reichenbach, P., 1997. Large alluvial fans in the north-central Po Plain (Northern Italy). *Geomorphology* 18, 119–136.
- Hancock, G.S., Anderson, R.S., 2002. Numerical modeling of fluvial strath-terrace formation in response to oscillating climate. *Geological Society of America Bulletin* 114, 1131–1142.
- 20 Harvey, A.M., 2002. The role of base-level change in the dissection of alluvial fans: case studies from southeast Spain and Nevada. *Geomorphology* 45, 67–87.
- Horton, B.K., DeCelles, P.G., 2002. Modern and ancient fluvial megafans in the foreland basin system of the central Andes, southern Bolivia: implications for drainage network evolution in fold-thrust belts. *Basin Research* 13, 43–63.
- Humphrey, N.F., Heller, P.L., 1995. Natural oscillations in coupled geomorphic systems: an alternative origin for cyclic sedimentation. *Geology* 23, 499–502.
- 25 Istanbuluoglu, E., Bras, R.L., 2005. Vegetation-modulated landscape evolution: effects of vegetation on landscape processes, drainage density, and topography. *Journal of Geophysical Research: Earth Surface*, 110 (F2), doi: 10.1029/2004JF000249.
- Jolivet, M., Labaume, P., Monié, P., Brunel, M., Arnaud, N., Campani, M., 2007. Thermochronology constraints for the propagation sequence of the south Pyrenean basement thrust system (France–Spain). *Tectonics* 26, TC5007 doi: 10.1029–2006TC002080.
- 30 Kim, W., Muto, T., 2007. Autogenic response of alluvial-bedrock transition to base-level variation: Experiment and theory. *Journal of Geophysical Research* 112, F03S14, doi: 10.1029/2006JF000561.

Margaux Mouchene 9/1/y 16:42

Supprimé: Dühnforth, M., Anderson, R.S., Ward, D., Stock, G.M., 2010. Bedrock fracture control of glacial erosion processes and rates. *Geology* 38 (5), 423–426 .

- Lague, D., 2010. Reduction of long-term bedrock incision efficiency by short-term alluvial cover intermittency. *Journal of Geophysical Research* 115, F02011 doi:10.1029–2008JF001210.
- Langston, A.L., Tucker, G.E., Anderson, R.S., 2015. Interpreting climate-modulated processes of terrace development along the Colorado Front Range using a landscape evolution model. *Journal of Geophysical Research: Earth Surface* 120, [2121-2138](#), doi: 10.1002–2014JF003403.
- Lavé, J., Avouac, J.-P., 2001. Fluvial incision and tectonic uplift across the Himalayas of central Nepal. *Journal of Geophysical Research* 106, 26561–26591.
- Leier, A.L., DeCelles, P.G., Pelletier, J.D., 2005. Mountains, monsoons, and megafans. *Geology* 33, 289–292.
- [Marshall, J.A., Roering, J.J., Bartlein, P.J., Gavin, D.G., Granger, D.E., Rempel, A.W., Praskievicz, S.J., Hales, T.C., 2015. Frost for the trees: Did climate increase erosion in unglaciated landscapes during the late Pleistocene? *Science Advances* 1, doi: 10.1126/sciadv.1500715.](#)
- Meyer, G.A., Wells, S.G., Jull, A.J.T., 1995. Fire and alluvial chronology in Yellowstone National Park: Climatic and intrinsic controls on Holocene geomorphic processes. *Geological Society of America Bulletin* 107, 1211–1230.
- Métivier, F., Gaudemer, Y., 1999. Stability of output fluxes of large rivers in South and East Asia during the last 2 million years: implications on floodplain processes. *Basin Research* 11, 293–303.
- Milana, J.P., Ruzycki, L., 1999. Alluvial-fan slope as a function of sediment transport efficiency. *Journal of Sedimentary Research* 69, 553–562.
- Mouchéné, M., van der Beek, P., Mouthereau, F., Carcaillet, J., [in press](#). Controls on Quaternary incision of the Northern Pyrenean foreland: chronological and geomorphological constraints from the Lannemezan megafan, SW France. *Geomorphology*.
- Mouchéné, M., 2016. Évolution post-orogénique du système couplé piémont/bassin versant : le méga-cône alluvial de Lannemezan et son bassin versant au Nord des Pyrénées. PhD thesis, Université Grenoble Alpes, 346pp.
- Murray, A.B., Paola, C., 1997. Properties of a cellular braided-stream model. *Earth Surface Processes and Landforms* 22, 1001–1025.
- Naylor, M., Sinclair, H.D., 2008. Pro-vs. retro-foreland basins. *Basin Research* 20, 285–303.
- [Nguyen, H. N., Vernant, P., Mazzotti, S., Khazaradze, G., and Asensio, E., 2016. 3-D GPS velocity field and its implications on the present-day post-orogenic deformation of the Western Alps and Pyrenees. *Solid Earth*, 7, 1349-1363, doi:10.5194/se-7-1349-2016.](#)
- Nicholas, A.P., Clarke, L., Quine, T.A., 2009. A numerical modelling and experimental study of flow width dynamics on alluvial fans. *Earth Surface Processes and Landforms* 34, 1985–1993.
- Nicholas, A.P., Quine, T.A., 2007. Modeling alluvial landform change in the absence of external environmental forcing. *Geology* 35, 527–530.
- Paris, J.-P., 1975. Carte géologique de la France à 1/50000. Feuille 1054 Montréjeau. Bureau des Recherches Géologiques et Minières, Orléans.


Margaux Mouchene 9/1/y 16:43

Supprimé: Submitted to


Margaux Mouchene 7/1/y 11:06

Supprimé: Nguyen, H. N., Vernant, P., Mazzotti, S., Khazaradze, G., and Asensio, E. 3D GPS velocity field and its implications on the present-day postorogenic deformation of the Western Alps and Pyrenees, *Solid Earth Discuss.*, doi:10.5194/se-2016-78, in review, 2016

- Pepin, E., Carretier, S., Herail, G., 2010. Erosion dynamics modelling in a coupled catchment–fan system with constant external forcing. *Geomorphology* 122, 78–90.
- Pepin, E., Carretier, S., Herail, G., Regard, V., Charrier, R., Farias, M., Garcia, V., Giambiagi, L., 2013. Pleistocene landscape entrenchment: a geomorphological mountain to foreland field case, the Las Tunas system, Argentina. *Basin Research* 25, 613–637.
- Poisson, B., Avouac, J.P., 2004. Holocene hydrological changes inferred from alluvial stream entrenchment in north Tian Shan (northwestern China). *The Journal of Geology* 112, 231–249.
- Roering, J.J., Kirchner, J.W., Dietrich, W.E., 1999. Evidence for nonlinear, diffusive sediment transport on hillslopes and implications for landscape morphology. *Water Resources Research* 35, 853–870.
- Rohais, S., Bonnet, S., Eschard, R., 2011. Sedimentary record of tectonic and climatic erosional perturbations in an experimental coupled catchment-fan system. *Basin Research* 24, 198–212.
- Schlunegger, F., Norton, K.P., 2014. Climate vs. tectonics: the competing roles of Late Oligocene warming and Alpine orogenesis in constructing alluvial megafan sequences in the North Alpine foreland basin. *Basin Research* 27, 230–245.
- Schumm, S.A., 1979. Geomorphic thresholds: the concept and its applications. *Transactions of the Institute of British Geographers* 485–515.
- Simpson, G.D.H., 2006. Modelling interactions between fold-thrust belt deformation, foreland flexure and surface mass transport. *Basin Research* 18, 125–143.
- [Sinclair, H.D., Gibson, M., Naylor, M., Morris, R.G., 2005. Asymmetric growth of the Pyrenees revealed through measurement and modeling of orogenic fluxes. *American Journal of Science* 305, 369–406.](#)
- Tucker, G.E., 2004. Drainage basin sensitivity to tectonic and climatic forcing: implications of a stochastic model for the role of entrainment and erosion thresholds. *Earth Surface Processes and Landforms* 29, 185–205.
- Tucker, G.E., Bradley, D.N., 2010. Trouble with diffusion: Reassessing hillslope erosion laws with a particle-based model. *Journal of Geophysical Research* 115, F00A10 doi: 10.1029–2009JF001264.
- van Dijk, M., Postma, G., Kleinhans, M.G., 2009. Autocyclic behaviour of fan deltas: an analogue experimental study. *Sedimentology* 56, 1569–1589.
- [Vergés, J., Fernández, M., Martínez, A., 2002. The Pyrenean orogen: pre-, syn-, and post-collisional evolution. In: Rosenbaum, G., Lister, G.S. \(Eds.\), *Reconstruction of the Evolution of the Alpine-Himalayan Orogen. Journal of the Virtual Explorer* 8, 57–74.](#)
- Whipple, K.X., Tucker, G.E., 1999. Dynamics of the stream-power river incision model: Implications for height limits of mountain ranges, landscape response timescales, and research needs. *Journal of Geophysical Research* 104, 17661–17674.
- Whipple, K.X., Tucker, G.E., 2002. Implications of sediment-flux-dependent river incision models for landscape evolution. *Journal of Geophysical Research* 107, doi: 10.1029–2000JB000044.
- Wobus, C.W., Tucker, G.E., Anderson, R.S., 2010. Does climate change create distinctive patterns of landscape incision? *Journal of Geophysical Research* 115, F04008.



5 **Figure 1:** The Lannemezan megafan and Neste catchment in the central northern Pyrenees (inset map shows location in southern France).



5 | Figure 2: The model grid consists of a mountain section, submitted to constant uplift U , and of a flat foreland section; both sections are submitted to precipitation. B.C. – Boundary Conditions; neither water nor sediment can cross a Closed B.C. while an Open B.C. corresponds to transverse rivers of fixed elevation capable of transporting both sediment and water fluxes out of the grid.

Margaux Mouchene 9/1/y 16:44
 Mis en forme: Police :italique





Figure 3: Building the megafan. Temporal evolution of the mean elevation change in the foreland and mountain and map views of the mountain and foreland landscapes (black lines are 100-m contour lines, water flux in blue shades) at the time steps marked with arrows, through the megafan building phase. (A) Drainage network initiates and propagates in the mountain block through headward incision while sediments are deposited along the front by regularly spaced streams. (B) Deposits merge in the foreland to form a *bajada* fed by a decreasing number of rivers as the mountain streams enlarge their basins. (C) Mean elevation in the range stabilizes and aggradation continues in the foreland, dominated by outflux of a central, main channel as the more lateral streams are drained directly toward the borders. (D) As aggradation continues, limited incision can occur along the borders of the fan (here on the western border) but (D') those thalwegs are quickly refilled. (E) Similarly, temporary incision can happen near the apex. (F) After 15.3 My, the megafan is built.




5 | Figure 4: Autogenic entrenchment. Temporal evolution of the mean elevation change in the foreland and mountain (black lines are 100-m contour lines, water flux in blue shades) at the time steps marked with arrows, through the autogenic incision phase (note change in time scale between this figure and Figure 3). Starting from the landscape obtained at 15.3 My (see F of Figure 3) the fan is incised when the main flow reaches the position of a small thalweg (e.g. G, I) but (H) continues to grow when the main flux overflows and migrates again on the fan. (J) After several of these cycles, the

Margaux Mouchene 9/1/y 16:45

Supprimé: a

5 | main flow is finally captured permanently in the thalweg. (K) As the main river now incises laterally towards the mountain front, a secondary stream is captured. (L) At the end of the experiment a large valley is incised along the front of the range. Topographic profiles across this valley (right panel) show that about 120m of incision occurs in the foreland at the time of capture (black profiles, inset J); subsequently the valley is mostly enlarged by lateral erosion in the foreland, deepened and enlarged near the apex, and markedly deepened in the mountain (red profiles, inset L; note that horizontal scale is different for each profile).

- Margaux Mouchene 9/1/y 16:46
- Supprimé: s
- Margaux Mouchene 9/1/y 16:45
- Supprimé: J
- Margaux Mouchene 9/1/y 16:46
- Supprimé: N



10 | Figure 5: Temporal elevation change of three locations in the mountain: bed of the main feeding river (red crosses), on nearby slope (black crosses) and ridge (blue crosses, see map view in right panel for locations). Following the entrenchment (marked with vertical grey line), the river rapidly incises, increasing (temporarily) the relief, as ridge elevations are not affected by the incision episode. The hillslope response is slow and lags behind that of the riverbed.




Figure 6: Experiment 2b (sinusoidal precipitation). Temporal evolution of (A) runoff, (B) mean elevation change in the foreland and (C) in the mountain. The trends of mean elevation change in the mountain and in the foreland are inversely correlated, and show a slight delay relative to the change in runoff, corresponding to the response time. (D) Temporal elevation change of the central valley floor in the mountain (location marked by red cross on inset map) during experiment 2b. The incision (blue arrows) related to humid periods (blue vertical lines are maximum runoff) is (over-) compensated in drier periods (grey vertical lines correspond to no runoff) by the uplift, so that the elevation generally increases through the experiment. The maximum incision is delayed from the maximum runoff (response time). Temporary deposits at the valley outlet around 230 ky dam the valley and trigger rapid backfilling of it, responsible for the high elevation between 230-310 ky; the following incision episode removes this dam.

Margaux Mouchene 9/1/y 16:47

Supprimé: trigger rapid backfilling (

Margaux Mouchene 9/1/y 16:47

Supprimé:) of




Figure 7: Model configuration at the end ($t=15.80\text{My}$) of experiment 3a (50-m drop in base-level at the onset of the incision phase). The megafan is incised by headward incision of a number of streams on its western and eastern borders (and marginally on the northern border).

5

Margaux Mouchene 9/1/y 16:47

Supprimé: initial

Margaux Mouchene 9/1/y 16:48

Supprimé: thalwegs




Figure 8: Experiment 4a, the incised stream first connects through headward incision to the secondary river (left, $t=300\text{ky}$) before being connected to the outlet of the central river at the end of the experiment (right, $t=500\text{ky}$).

10

Margaux Mouchene 9/1/y 16:48

Supprimé: thalweg




Figure 9: (A) Scaling relationship between fan area and fan slope for alluvial systems of the Alps, the Andes and the Himalaya; data from Guzzetti et al. (1997; crosses) and Horton and DeCelles (2002; open circles). Thick line is the best power-law fit to the combined data; $S_f = 2.42 A_f^{-0.30}$. Tilt (vertical arrow) for the Lannemezan megafan (red circle) is estimated as the difference between present-day slope and predicted slope from the power-law fit. (B) Scaling relationship between drainage area and fan slope for alluvial systems of the Alps, data from Guzzetti et al. (1997; crosses), Crosta and Frattini (2004; triangles) and Champagnac et al. (2008; black circle = Valensole, grey square = Chambaran). The Lannemezan megafan/Neste system (red circle) lies slightly out of the relation (fit; $S_f = 10.4 A_D^{-0.51 \pm 0.05}$). Quaternary tilt (vertical arrow) of the Lannemezan megafan surface is estimated at 0.13° ; the difference between present-day slope and slope predicted by the power-law fit. Modified after Champagnac et al. (2008).




Figure 10: Tilting experiments 5a (0.13° tilt) and 5b (0.4° tilt). Connexion with the headward incising stream occurs (at 280ky and 240 ky, respectively) but model instabilities in the channel, interpreted as deposition, induce overflowing (at 320ky and 260 ky, respectively). Distributive flow over the megafan resumes and lasts until the end of the experiment.

Margaux Mouchene 9/1/y 16:48
Supprimé:

Margaux Mouchene 9/1/y 16:48
Supprimé:

Margaux Mouchene 9/1/y 16:48
Supprimé: as

Margaux Mouchene 9/1/y 16:49
Supprimé: thalweg

Margaux Mouchene 9/1/y 16:49
Supprimé: ing




Figure 11: Evolution of a north-south topographic profile near the middle of the model during Experiment 5 (region-wide tilting). Inset shows that with default settings (Experiment 1), the slope of the megafan does not increase; regional tilting (Experiment 5) is needed to create an increasing northward slope through time.

5

- Margaux Mouchene 9/1/y 16:49
Supprimé: e
- Margaux Mouchene 9/1/y 16:49
Supprimé: e
- Margaux Mouchene 9/1/y 16:49
Supprimé: e
- Margaux Mouchene 9/1/y 16:49
Supprimé: ,
- Margaux Mouchene 9/1/y 16:49
Supprimé: e
- Margaux Mouchene 9/1/y 16:49
Supprimé: e





Figure 12: Comparison between (A) DEM of the Northern Pyrenees and foreland at the longitude of the Lannemezan megafan and (B) the final model (contour lines every 100 m) shown at the same scale. The proposed evolution for the Lannemezan megafan is schematized below: (C) A pre-existing river Ariège/lower Garonne River flows through the foreland while the Neste transports sediments deposited in the foreland through a distributive pattern to build a megafan; (D) a tributary of the Ariège/lower Garonne retreats headward and westward, towards the apex of the Lannemezan megafan while the megafan keeps growing; (E) the migration of the tributary leads to sequential capture of (1) the upper Garonne and (2) the upper Neste and thus the abandonment of the megafan; rivers incise together in the foreland deposit leaving a series of alluvial terraces (preferably on the left bank due to the direction of river migration indicated by white arrows).

- Margaux Mouchene 9/1/y 16:50
- Supprimé: an
- Margaux Mouchene 9/1/y 16:51
- Supprimé:
- Margaux Mouchene 9/1/y 16:51
- Supprimé: and range DEM and
- Margaux Mouchene 9/1/y 16:51
- Supprimé: right;
- Margaux Mouchene 9/1/y 16:51
- Supprimé: s
- Margaux Mouchene 9/1/y 16:52
- Supprimé: ,

τ_c	K_{br}	K_{all}	m	n	p	L	α
15 Pa	$0.5 \cdot 10^{-3}$	$4 \cdot 10^{-3}$	0.6	0.7	1	0.3	0.01

Settings						Results	
Experiment number	Precipitation rate	Precipitation occurrence	Base level	Uplift rate	Permanent entrenchment	Time of entrenchment after end of building at 15.3 My	
	(m y^{-1})	(Fraction of time step)	(m asl)	(mm y^{-1})		(ky)	
1-default	1	1	0	0.3	Yes	270	
Climate	2a1	0.5	1	0	No	-	
	2a2	2	1	0	Yes	180	
	2b	1	sinusoidal	0	No	-	
	2c	1	0.5	0	Yes	150	
Base Level	3a	1	1	-50	Yes	250	
Uplift	4a	1	1	0	Yes	500	
	4b	1	1	0	Yes	310	
	4c	1	1	0	Yes	500	
Tilting	5a	1	1	0	No	-	
	5b	1	1	0	No	-	

Table 1: **Parameter values used in the experiments.** Top: Fixed parameters for all model runs τ_c is the critical shear stress, K_{br} and K_{all} are the bedrock and sediment erodibility, respectively, m , n and p are coefficients for the fluvial erosion law, L is the transport length and α is the lateral erosion coefficient. Bottom: Model settings for the experimental runs and results.