



Glacial Isostatic Adjustment modelling: historical perspectives, recent advances, and future directions

Pippa L. Whitehouse¹

¹Department of Geography, Durham University, Durham, DH1 3LE, UK

Correspondence to: Pippa L. Whitehouse (pippa.whitehouse@durham.ac.uk)

5 **Abstract.** Glacial Isostatic Adjustment (GIA) describes the response of the solid Earth, the gravitational field, and
consequently the oceans to the growth and decay of the global ice sheets. It is a process that takes place relatively rapidly,
triggering 100 m-scale changes in sea level and solid Earth deformation over just a few tens of thousands of years. Indeed,
the first-order effects of GIA could already be quantified several hundred years ago without reliance on precise measurement
10 techniques and scientists have been developing a unifying theory for the observations for over 200 years. Progress towards
this goal required a number of significant breakthroughs to be made, including the recognition that ice sheets were once
more extensive, the solid Earth changes shape over time, and gravity plays a central role in determining the pattern of sea-
level change. This article describes in detail the historical development of the field of GIA and an overview of the processes
involved. Significant recent progress has been made as concepts associated with GIA have begun to be incorporated into
15 parallel fields of research; these advances are discussed, along with the role that GIA is likely to play in addressing
outstanding research questions within the field of Earth system modelling.

1 Introduction

The response of the solid Earth to the collapse of northern hemisphere ice sheets following the Last Glacial Maximum
(LGM) continues today at a rate so large (>10 mm/yr, e.g. Sella et al., 2007; Lidberg et al., 2010) that glacial isostatic
adjustment (GIA) is one of the few geophysical processes that can be readily observed on human timescales without recourse
20 to sophisticated scientific measurement techniques. For this reason, the more easily-observed impacts of GIA, such as
shoreline migration, played an important role in motivating the development of ideas associated with climate cycles, sea-
level change, geodesy and isostasy during the 19th and early 20th century (Jamieson, 1865; Croll, 1875; Woodward,
1888; Nansen, 1921; Daly, 1925). The field has a long history of combining observations with theory, and the exchange of
ideas between scientists working in a suite of different disciplines has repeatedly resulted in important scientific
25 breakthroughs.

The modern field of GIA addresses the classic geodynamics problem of determining the solid Earth response to surface load
changes by ice and ocean water, whilst at the same time solving for the gravitationally-consistent redistribution of meltwater
across the global ocean. Calculations are necessarily carried out on a global scale, and numerical models of GIA consider the
30 behaviour of three fundamental components of the Earth system: the solid Earth, the global ocean, and the ice sheets (see
Figure 1). Inputs to a GIA model typically include a priori estimates for the history of global ice-sheet change and the



35 rheology of the solid Earth, with changes to the shape of the ocean and the solid Earth being determined by solving the sea-
level equation (Farrell and Clark, 1976; see also Sect. 2.2). A wealth of data can be used to determine the details of GIA
model inputs, e.g. geological evidence can provide information on past ice extent (Bentley et al., 2014), while modelling of
mantle dynamics can be used to determine independent constraints on mantle viscosity (e.g. Mitrovica and Forte, 2004).
However, the reason GIA is of interest across so many disciplines is that the problem can be turned around, and observations
relating to past sea-level change or solid Earth deformation – the classical ‘outputs’ of a GIA model (Figure 1) – can be used
to infer information relating to the ‘inputs’, namely ice-sheet history and Earth rheology (e.g. Lambeck et al., 1998;Peltier,
2004). As in all disciplines where data play a crucial role in determining model parameters, uncertainties and
40 spatial/temporal gaps in the data leave room for non-uniqueness in the solutions invoked to explain the observations, but
intellectual input from a diverse range of sources over the last 200 years has helped to steer the field towards robust
explanations for the varied range of processes that are associated with GIA.

In this article the development of GIA modelling is traced from initial observations of rapid shoreline migration in 15th
45 century Sweden through to sophisticated approaches that incorporate feedbacks between ice, ocean and solid Earth
dynamics. The historical development of the field, underlying theory, important results, and data sets used to constrain GIA
modelling are discussed in Sect. 2, recent developments in the field are discussed in Sect. 3, and the article concludes with a
discussion of unresolved questions that warrant future attention (Sect. 4). We begin by motivating this review with a brief
summary of the fields that have been influenced by studies of GIA.

50 **1.1 Applications of GIA**

GIA plays a role in studies that span the fields of climate, cryosphere, geodesy, geodynamics, geomorphology, and natural
hazards. Some of the fundamental scientific questions that require consideration of GIA include: (i) linking ice-sheet
response to past climatic change; (ii) understanding the rheology of the interior of the Earth; (iii) determining present day ice
sheet mass balance and sea-level change; (iv) interpreting palaeo-sea-level records; (v) understanding ice-sheet dynamics;
55 (vi) quantifying tectonic hazard; (vii) reconstructing palaeo-drainage systems; (viii) interpreting the rotational state of the
Earth; (ix) understanding coastal change; and (x) understanding the causes of volcanism. Comparison with data is a central
component of GIA studies, and in many cases misfits between observations and predictions have led to new understanding
of factors that had not previously been considered, such as feedbacks between GIA and ice dynamics or the influence of
lateral variations in mantle rheology. The historical development of this multi-faceted subject is an interesting story.



60 2 Review of GIA modelling

2.1 Historical perspective

The people of Scandinavia must have been aware of the effects of GIA for many centuries. As an example, by 1491 AD the ancient port of Östhammar could no longer be reached by boat and the city had to be relocated closer to the sea (Ekman, 2009). However, it was not until 1731 AD that rigorous measurements of relative sea-level change – i.e. change in local
65 water depth – were initiated with the cutting of a ‘mean sea level mark’ into a coastal rock at Lövgrund, Sweden (Celsius, 1743). Using historical documents to extend the record of sea-level change back to 1563 AD, Celsius (1743) carried out the first calculations associated with GIA and determined that sea-level in the Gulf of Bothnia was falling at a rate of 1.4 cm/yr relative to the height of the land. He assumed that the change was due to a fall in sea level but at the time there was considerable disagreement, with others proposing that the cause was land uplift (see Ekman (2009) for a thorough review of
70 the subject).

The question was partly resolved by considering evidence for relative sea-level change in different locations around the world. Playfair (1802) noted that past sea levels had been higher in such diverse locations as Scotland, the Baltic, and the Pacific, but lower in the Mediterranean and southern England. Ideas associated with the concept of an equipotential surface
75 were yet to be put forward (e.g. Stokes, 1849), and therefore Playfair (1802) argued that since “the ocean... cannot rise in one place and fall in another” the differences must be associated with land level changes. Without any means to determine the timing of past relative sea-level change in different locations this argument is flawed, but more robust evidence was provided by Lyell (1835), who used an ingenious variety of observations to determine that the *rate* of relative sea-level change across Sweden varied from place to place. Following a similar argument to Playfair (1802), Lyell (1835) concluded
80 that his observations could only be explained by variations in the rate of land uplift, since (he assumed) sea-level fall would produce a spatially-uniform rate of change.

These early studies explored a number of explanations for the change in the height of the land, and the idea that an ice sheet could have depressed the land was first proposed by Jamieson (1865). He was familiar with the geomorphological evidence
85 for past ice cover across Scotland, but made the important observation that whilst marine deposits could be found well above current sea level near the coast, they were not present in lower areas in the interior. This led him to propose that the weight of the ice sheet must have depressed the coastal land below sea level, but that the presence of the ice prevented the interior from being flooded. This was the first suggestion that an ice sheet could deform the Earth, and whilst such ideas were relatively quickly taken up by field scientists (e.g. Chamberlin and Salisbury, 1885), they were not widely accepted by those
90 who took a more theoretical approach, largely due to ongoing disagreement regarding the structure of the interior of the Earth and the concept of isostasy (e.g. Barrell, 1919). Crucially, the idea that the surface of the Earth could deform in response to a change in surface load was neglected by those who first considered the effect of gravity on sea level.



95 An important contribution to the field came from Croll (1875), who proposed that there had been repeated glacial cycles and
hence periodic changes in the distribution of mass throughout the Earth system. Based on his assumption that the ocean takes
on a spherical form around the centre of mass of the Earth, he calculated the displacement of the centre of mass due to the
presence of an ice-cap, and found that the centre of mass of the system would be displaced towards the ice cap, thus
providing an explanation for the observation that sea levels were higher during glacial times in Scotland. However, Croll's
100 (1875) theory was flawed in two important ways: (i) he dismissed solid Earth deformation to be a local effect and did not
link it to ice loading, and (ii) he did not appreciate that the redistribution of mass throughout the Earth system would alter the
shape of the ocean surface. This second issue was addressed in detail by Woodward (1888) around a decade later.

Woodward's (1888) interest in the shape of the Earth's gravitational field was motivated by questions posed to him
regarding the differing elevations of contemporaneous palaeo-shorelines of the former Lake Bonneville (Gilbert, 1885b), and
105 the tilt of lake shorelines and glacial deposits associated with the past glaciation of North America (Chamberlin and
Salisbury, 1885; Gilbert, 1885a). The authors of these studies were supporters of the hypothesis that surface load changes, in
the form of water, ice or sediment, could deform the surface of the Earth, and they deduced that the solid Earth response to
surface loading could provide an explanation for their observations (Chamberlin and Salisbury, 1885; Gilbert, 1885b).
Indeed, Gilbert (1890) used palaeo-shoreline observations from Lake Bonneville to draw early conclusions on the
110 rheological properties of the Earth. However, building on the ideas of Croll (1875), this group of scientists also wondered
whether gravitational attraction, e.g. of an ice sheet, played a role in explaining their observations.

They turned to Woodward (1888) for the answer, and he carried out detailed calculations relating to the change in the shape
of the geoid that would arise due to the redistribution of surface mass associated with the appearance/disappearance of an ice
115 sheet. He used realistic estimates for the shape and size of an ice sheet, and took into account the self-attraction of the ocean
and the different densities of ice and water. He also appreciated the need to conserve mass when transferring water between
the ice sheets and the ocean, but addressed a simplified problem in which ice is transferred directly between the two polar
regions (cf. Croll, 1875), i.e. his calculations assumed no net change in ocean volume. The key result of this work was a
prediction of the perturbation to the height of the geoid at a series of radial distances from the centre of a growing ice sheet.
120 Although he did not formally account for changes in ocean volume, the magnitude of the geoid perturbation in the near-field
of the ice sheet led Woodward (1888) to hypothesize that as the ice sheet formed, water depths in the near-field would
increase, despite a net decrease in ocean volume – a result that still surprises many people today!

The one shortcoming of Woodward's (1888) analysis was his decision to neglect the deformation of the Earth in response to
125 surface loading. This led him to conclude that the volume of ice needed to explain the tilt of palaeo-shorelines in the Great
Lakes region (Gilbert, 1885a) was unfeasibly large. If he had been able to include an estimate of Earth deformation he may



well have realized that the palaeo-shorelines could be explained by a combination of postglacial rebound and tilting of the geoid surface due to the attraction of the former ice sheet (e.g. Figure 2).

130 Important advances towards understanding the role of Earth deformation were made by Nansen (1921, p.288), who used the
concepts of isostasy and mass balance to show that sea-level change could be explained by some combination of ice-sheet
melt and land deformation: “along great parts of the coasts of Fenno-Scandia, this rise of sea-level was more or less masked
by the still faster upheaval of the land, and it was only during certain periods when the temperature was much raised and the
melting of the ice caps much increased, that the rise of sea-level was sufficiently rapid to cause a pause in the negative shift
135 of the shoreline so considerable that conspicuous marine terraces, beaches, or shorelines could be developed.”

Nansen (1921) did not consider the gravitational effect of mass redistribution when discussing the causes of relative sea-
level change, but he did understand that the Earth would continue to deform viscously for a prolonged period following mass
redistribution, and hence that relative sea-level change would continue after the volume of the ocean stabilized. This
140 reasoning led him to suggest that ongoing subsidence of the seafloor due to the past addition of meltwater to the ocean could
explain observations of recent global sea-level fall (Daly, 1920) and that the growth and decay of peripheral bulges within
the ocean need to be accounted for when calculating the magnitude of sea-level change during a glacial cycle. This latter
point is an early description of the ‘ocean syphoning’ effect (see Sect. 2.2.4 and Figure 2b). Daly (1925) further explored the
implications of solid Earth deformation when seeking to interpret an impressive array of sea-level observations from around
145 the world, making use of the idea that the Earth would respond both elastically and viscously in response to surface loading.
He also highlighted the important, but little-known, work of Rudzki (1899), who took Woodward’s (1888) geoid calculations
and used them to re-calculate the spatially-variable sea-level change that would result from the melting of a circular ice cap,
but now accounting for the combined effects of (elastic) Earth deformation, the change in ocean volume, and the change in
the shape of the gravitational field due to the redistribution of both ice and solid Earth mass.

150 By the 1920’s a range of approaches had been used to estimate the magnitude of the sea-level lowstand at the peak of the last
glaciation (Daly, 1925). Nansen’s (1921) estimates for global ice volume were based on calculations linking the magnitude
of depression beneath the former ice sheets (evidenced by marine deposits, which are now located above present sea level) to
the relative density of ice and upper mantle material, accounting for the fact that some mantle material would be laterally
displaced to form peripheral bulges. Impressively, his estimate for the mean sea-level change associated with “the formation
155 of the Pleistocene ice-caps” was 130 m, a value that is almost identical to contemporary estimates (Lambeck et al., 2014).
Nansen’s (1921) calculations accounted for changes in the area of the ocean through time, and he identified several factors
that necessitate an iterative approach to calculating changes in sea level. In particular, he noted that since ice loading deforms
both continental and oceanic areas, the true change in water depth (which determines the deformation due to ocean loading)
160 will be different to the value that would be determined by considering a constant ocean area and a non-deforming Earth.



These feedbacks between ice and ocean loading are a fundamental feature of the sea-level equation (Farrell and Clark, 1976), which forms the basis of all contemporary GIA models.

165 These early scientists made impressive use of the data available to them, and the final piece of the puzzle came with the ability to determine the timing of past environmental change (e.g. De Geer, 1912), which allowed the first estimate for the viscosity of the mantle to be determined (Haskell, 1935). Studies into the viscosity of Earth's mantle developed rapidly from the 1960's onwards (e.g. McConnell, 1968; O'Connell, 1971; Cathles, 1975), and the stage was set for a global model of GIA that accounted for (i) ice-ocean mass conservation, (ii) viscoelastic deformation of the Earth, and (iii) gravitationally self-consistent perturbations to the shape of the geoid. The following section outlines the modelling approach that was developed
170 in the 1970's to address these issues. The models developed during this period underpin all contemporary studies of GIA, and provide confirmation of many of the fundamental ideas proposed during the 19th and early 20th century.

2.2 Development of numerical models of GIA

2.2.1 The sea-level equation

175 When modelling processes associated with GIA, water from a melting ice sheet is assumed to be instantaneously redistributed across the global ocean according to the shape of the geoid, where the geoid is an equipotential surface that defines the shape of the sea surface in the absence of dynamic forcing by atmospheric or oceanic circulation. The shape of the geoid depends on the distribution of mass throughout the Earth system. There are feedbacks to be considered because the change in the distribution of surface mass (e.g. the decrease in the mass of the ice sheet and the addition of mass to the ocean) must be taken into account when calculating the shape of the geoid as the meltwater is redistributed (Figure 2).
180 However, the situation is more complicated than this because the shrinking of the ice sheet and the transferral of water to the ocean causes the solid Earth to deform, and this redistribution of mass *inside* the Earth further alters the shape of the geoid. Calculations to determine the change in sea level due to the melting of an ice sheet must therefore be carried out iteratively. The problem is further complicated by the fact that the deformation of the Earth reflects both contemporary and past surface mass change due to the viscoelastic properties of the mantle (Cathles, 1975).

185 There are two fundamental unknowns within GIA: the history of the global ice sheets and the rheology of the solid Earth (Figure 1). These are traditionally determined via an iterative approach, using a range of data (see Sect. 2.3). Once they are known, or once a first estimate has been determined, then the spatially-varying history of sea-level change can be uniquely determined by solving the sea-level equation, which defines the gravitationally self-consistent redistribution of meltwater across the ocean. The theoretical development of the sea-level equation is covered extensively elsewhere (e.g. Farrell and Clark, 1976; Mitrovica and Milne, 2003; Spada and Stocchi, 2006), and hence we briefly restate the main form of the equation
190 here:



$$\Delta SL(\theta, \psi, t) = \frac{\rho_i}{\gamma} G_S \otimes_i I + \frac{\rho_w}{\gamma} G_S \otimes_o \Delta SL + C_{SL}(t) \quad (1)$$

195

$\Delta SL(\theta, \psi, t)$ is the change in sea level (or, equivalently, water depth) at co-latitude θ and longitude ψ , between time t and some reference time t_0 , I is the spatio-temporal evolution of global ice thickness change, ρ_i and ρ_w are ice and ocean water density, respectively, γ is the acceleration due to gravity at Earth's surface, G_S represents a Green's function that describes perturbations to the solid Earth displacement field and the gravitational potential due to surface loading, constructed by combining viscoelastic surface load Love numbers (Peltier, 1974; Spada and Stocchi, 2006), and \otimes_i and \otimes_o represent convolutions in space and time over the ice sheets and the ocean, respectively. Note the appearance of ΔSL on both sides of the equation, indicating that the sea-level equation is an integral equation and an iterative approach is required to solve it. The first two terms on the right-hand side of Eq. (1) are spatially-varying terms that describe the perturbation to sea level due to ice and ocean loading respectively, while $C_{SL}(t)$ is a time-dependent uniform shift in the geoid that is invoked to satisfy conservation of mass:

200

205

$$C_{SL}(t) = -\frac{m_i(t)}{\rho_w A_0(t)} - \frac{\rho_i}{\gamma} \overline{G_S \otimes_i I} - \frac{\rho_w}{\gamma} \overline{G_S \otimes_o \Delta SL} \quad (2)$$

The first term in Eq. (2) is often referred to as the 'eustatic' term; it describes the spatially uniform sea level change that takes place across ocean area $A_0(t)$ due to a change in ice mass of magnitude $m_i(t)$, in the absence of any solid Earth deformation. The final two terms are the spatial average over the ocean (indicated by the overbar) of the spatially-varying terms in Eq. (1). These final terms must be subtracted because although the mean of the spatially-varying perturbations will be zero when integrated over the whole of Earth's surface, the mean will not necessarily be zero when integrated over the ocean; hence a uniform shift is applied to conserve mass.

210

215

The following sections provide further insight into specific components of the sea-level equation, and a description of how it has been used to provide confirmation of several global-scale processes that were hypothesized during the 19th and early 20th centuries.

220

2.2.2 Representation of the solid Earth

In order to calculate the solid Earth response to surface load change over glacial timescales the Earth is commonly assumed to be a linear Maxwell viscoelastic solid (Peltier, 1974), although a number of studies alternatively adopt a power-law approach (Wu, 1998). The spatially-variable, time-dependent response of a Maxwell solid to surface load change can be calculated using viscoelastic Love numbers, which define the response of a spherically-symmetric, self-gravitating, viscoelastic sphere to an impulse point load (Peltier, 1974; Wu, 1978; Han and Wahr, 1995). The Love numbers reflect the assumed viscosity profile of the mantle, which must be defined a priori. Alternatively, if a power-law approach is used, the

225



effective viscosity of the mantle will depend on the stress field throughout the mantle, which depends on surface load change. The non-linear stress-strain relationships that form the basis of the power-law approach are based on the results of laboratory experiments that seek to understand the controls on deformation within the mantle (Hirth and Kohlstedt, 2003).
230 For both approaches the elastic and density structure throughout the Earth must be defined (e.g. Dziewonski and Anderson, 1981), and the deformation of the whole Earth must be considered (recall that the sea-level equation solves for global meltwater distribution).

In a GIA model the lithosphere is typically represented by an elastic layer at the surface of the Earth. The thickness of this
235 layer influences the wavelength of deformation, while the viscosity of the mantle controls the rate of deformation. It therefore follows that the rheological properties of the Earth may be inferred from observations (Figure 1) that reflect land uplift or subsidence in response to ice and ocean load change (e.g. Lambeck et al., 1998; Paulson et al., 2007a; Lambeck et al., 2014; Peltier et al., 2015; Lau et al., 2016; Nakada et al., 2016). However, in reality, poor data coverage, uncertainties associated with the ice load history, and spatial variations in Earth rheology make it difficult to uniquely determine an
240 optimal solution for Earth properties such as lithosphere thickness or mantle viscosity. To overcome this, some studies consider multiple geodynamic processes when seeking to constrain mantle rheology (e.g. Mitrovica and Forte, 2004), while others use independent data sets to define the rheological properties of the Earth. As an example, seismic wave speeds can be related to the temperature distribution in the mantle, which in turn may be related to mantle viscosity (Ivins and Sammis, 1995). This approach is discussed in more detail in Sect. 3.2.

245

2.2.3 Extensions to the sea-level equation

The original statement of the sea-level equation by Farrell and Clark (1976) did not account for temporal variations in ocean area, which can arise via two processes (Figure 3). First, since the ocean is not typically bounded by vertical cliffs, a rise or fall in sea level at a particular location will result in onlap or offlap, and hence an increase or decrease in the area of the
250 ocean, respectively. This issue was first addressed in a self-consistent manner by Johnston (1993). Secondly, during past glacial periods all the major ice sheets grew beyond the confines of the continent on which they were initially situated, expanding into the ocean and forming large areas of marine-grounded ice. Temporal variations in the extent of a marine-grounded ice sheet will alter the ocean area over which meltwater can be redistributed. The treatment of marine-grounded ice within the sea-level equation was first discussed by Milne (1998), and a detailed description of how to implement shoreline migration due to both processes is given in Mitrovica and Milne (2003).
255

An additional extension to the sea-level equation involves the treatment of rotational feedback (Figure 4). It is clear that since GIA alters the distribution of mass throughout the Earth system this will perturb the magnitude and direction of Earth's rotation vector (e.g. Nakiboglu and Lambeck, 1980; Sabadini et al., 1982; Wu and Peltier, 1984). These changes will, in turn,
260 affect a number of processes associated with GIA: changing the Earth's rotation vector will instantaneously alter the shape of



the sea surface, i.e. the shape of the geoid (Milne and Mitrovica, 1998), while over longer timescales it will excite solid Earth deformation (Han and Wahr, 1989), thus further altering the shape of the geoid (Figure 4a). Over both timescales these mechanisms result in a change in the distribution of water across the ocean and this will excite additional solid Earth deformation, thus further altering the rotational state of the Earth. These feedbacks were first implemented within the sea-level equation by Milne and Mitrovica (1998), and a number of important updates to the theory have been made in recent years (Mitrovica et al., 2005; Mitrovica and Wahr, 2011; Martinec and Hagedoorn, 2014).

2.2.4 Confirmation of early theories and implications for the interpretation of sea-level records

Solutions to the sea-level equation reflect processes that were originally described by Jamieson (1865), Croll (1875), Gilbert (1885b), Woodward (1888), and Nansen (1921). Temporal variations in water depth arise due to changes in the total mass of the ocean (as described by Croll (1875)) and the shape of its two bounding surfaces; the sea surface (as proposed by Woodward (1888)) and the solid Earth (as proposed by Jamieson (1865) and Gilbert (1885b)). Furthermore, as suggested by Nansen (1921), the shape of these bounding surfaces will continue to evolve even during periods of constant global ice mass due to the transient nature of the solid Earth response to surface loading. Observations of relative sea-level change therefore require careful interpretation, particularly if they are to be used to determine changes in global ice volume.

The magnitude of sea-level change in the far-field of the major ice sheets has long been used to constrain changes in global ice volume (e.g. Fairbanks, 1989; Fleming et al., 1998; Milne et al., 2002), but this approach is complicated by the fact that the location at which ‘eustatic’ or mean sea-level change is recorded will vary over time (Milne and Mitrovica, 2008). It is clear that sea-level change in the *near-field* of an ice sheet will reflect perturbations to the shape of the geoid and the solid Earth due to the presence, and loading effect, of the evolving ice sheet as well as changes in total ocean mass (e.g. Shennan et al., 2002), but *far-field* records of sea-level change will also be biased by long-wavelength, spatially-varying processes associated with GIA. The most important of these processes are outlined below:

- **Meltwater fingerprints (based on theory by Woodward, 1888):** Sea-level change associated with the addition of meltwater to the ocean will be spatially-variable (Milne et al., 2009); a decrease in water depth will be recorded in the near-field of a melting ice sheet due to solid Earth rebound and a fall in the height of the geoid in response to the decrease in ice mass (Figure 2a). Consequently, the increase in water depth far from the melting ice sheet will be greater than the global mean. Predictions of this ‘fingerprint’ of sea-level change associated with different ice-sheet melt scenarios have been used to distinguish between melt sources during past and present periods of rapid sea-level change (Mitrovica et al., 2001; Clark et al., 2002; Hay et al., 2015).
- **Ocean syphoning (originally hypothesized by Nansen, 1921):** In the same way that rebound in response to ice mass loss can persist for many thousands of years, subsidence of peripheral bulge regions also continues long after the ice



295 sheets have melted. These peripheral bulge regions surround the former ice sheets and are typically located offshore, and
hence their collapse acts to increase the capacity of the ocean basins (Figure 2b). In the absence of significant changes in
ocean volume, peripheral bulge collapse will result in a fall in absolute mean sea level (the height of the sea surface
relative to the centre of the Earth) even though global mean water depth will be unchanged. This ‘ocean syphoning’
effect explains why mid-Holocene sea-level highstands are observed across many equatorial regions (Mitrovica and
300 Peltier, 1991b; Mitrovica and Milne, 2002), and it must also be accounted for when interpreting contemporary
measurements of global sea-level change derived from satellite altimetry (Tamisiea, 2011). At sites located on a
subsiding peripheral bulge, relative sea-level rise will occur throughout an interglacial period, even if global ice volumes
remain roughly constant (Lambeck et al., 2012).

305 - **Continental levering:** During the LGM lowstand many continental shelves were sub-aerially exposed. Loading by the
ocean during the subsequent sea-level rise will have caused the newly-submerged continental shelves to be flexed
downwards and the margins of the continents to be flexed upwards (Walcott, 1972). This ‘continental levering’ effect
must be accounted for when interpreting sea-level records recovered from regions adjacent to extensive continental
shelves. In particular, it should be noted that coastlines orientated perpendicular to the continental shelf break will
310 experience differential amounts of uplift (e.g. Lambeck and Nakada, 1990; Clement et al., 2016).

2.2.5 Numerical methods to solve the sea-level equation

The sea-level equation is most commonly solved using a pseudo-spectral approach (e.g. Mitrovica and Peltier,
1991b; Mitrovica and Milne, 2003; Kendall et al., 2005; Spada and Stocchi, 2006; Adhikari et al., 2016). However, finite
315 element (e.g. Wu and van der Wal, 2003; Zhong et al., 2003; Paulson et al., 2005; Dal Forno et al., 2012), spectral-finite
element (e.g. Martinec, 2000; Tanaka et al., 2011), and finite volume (e.g. Latychev et al., 2005b) approaches have also been
used, while approaches that use the adjoint method are under development (Al-Attar and Tromp, 2014; Martinec et al., 2015).
The equations used to represent solid Earth deformation may differ between these approaches. A description of the different
methods used to determine the response of the solid Earth to surface loading is given in the GIA benchmarking study of
320 Spada et al. (2011). In all cases, an iterative approach is required to determine a self-consistent solution to the sea-level
equation due to its integral form.

2.3 Data

A fundamental component of GIA modelling is the use of data to constrain unknown factors associated with the ice history
and Earth rheology (Figure 1). Different data have different roles. For example, dated geomorphological evidence for past
325 ice extent can be used to define the surface load history, while observations relating to solid Earth rebound, such as relative
sea-level indicators or GPS data, can be used to tune the rheological model. There exist strong trade-offs between the timing
and the magnitude of past surface load change (Figure 5a), as well as between the load history and the assumed rheology



(Figure 5b). One way to address this non-uniqueness is to independently constrain ice history and Earth rheology outside the confines of the GIA model. Although, datasets that are sensitive to both factors, such as observations relating to past sea-level change, provide very powerful constraints on the coupled problem (e.g. Lambeck et al., 1998). In all cases where data are used to tune a GIA model, it is important to assess whether there are unmodelled processes reflected in the data that may bias the results, and care must be taken to assign realistic errors. The key datasets used in studies of GIA are briefly described below.

2.3.1 Relative sea-level data

A sea-level indicator is a piece of evidence that provides information on past sea level. In order to be compared with GIA model output, the age and current elevation of a sea-level indicator must be known (including associated uncertainties), as well as the relationship between the sea-level indicator and mean sea level. Past relative sea-level change will be preserved in the geological record as a change in the position of the shoreline or a change in water depth. Past shoreline change can be reconstructed by identifying the time at which a particular location was inundated by, or isolated from, the ocean (Figure 3). Such information can be derived from microfossil analysis of the sediment contained within isolation basins (lakes which were previously connected to the ocean, or former lakes that are now drowned) (e.g. Watcham et al., 2011), or by determining the age of an abandoned beach ridge, after accounting for the offset between the beach ridge and mean sea level in the modern setting. In some cases, sea-level indicators may only indicate whether a particular location was previously above or below sea level. For example, archaeological artefacts typically provide an upper bound on contemporaneous sea level, while the presence of any type of *in situ* marine material provides a lower bound on past sea level. More specifically, if a fossil shell or coral is found still in its growth position (either above or below present sea level), and its living-depth range is known, this can be used to determine past water depths (e.g. Deschamps et al., 2012). Although, note that temporal variations in local conditions, e.g. changes in water properties or tidal range, can alter the depth at which a particular species can survive (Hibbert et al., 2016). At higher latitudes, reconstructions of saltmarsh environments have proved very useful for determining not only past changes in water depth, but also more subtle information relating to whether sea level was rising or falling in the past (Barlow et al., 2013). Finally, if past shorelines can be continuously reconstructed over length scales of a few kilometres or more, then the subsequent warping of these contemporaneous surfaces provides a powerful constraint on GIA (McConnell, 1968).

2.3.2 Ice extent data

Data relating to past ice extent, thickness, and flow direction all contribute useful information to ice-sheet reconstructions, with the latter providing an indication of past ice-sheet dynamics, and hence the location of former ice domes (e.g. Margold et al., 2015). Terrestrial and marine geomorphological features that must have formed at the margin of a former ice sheet, such as moraines, grounding zone wedges, or deposits relating to ice-dammed lakes, can be used to build a picture of past ice extent if the age of the features can be precisely dated. Indeed, a series of snapshots of past ice-sheet extent have been



constructed from geomorphological data for the Laurentide, British-Irish, and Fennoscandian ice sheets (Dyke et al., 2003; Clark et al., 2012; Hughes et al., 2016). In contrast, determining past ice *thickness* over large spatial scales is more difficult. Field-based reconstructions of past ice thickness typically rely on cosmogenic exposure dating to determine when, and to what depth, mountain ranges in the interior of a former ice sheet were last covered by ice (Ballantyne, 2010). Care must be taken when interpreting such information because complex topography will perturb the local ice flow, with the result that local ice thickness fluctuations may not represent regional scale ice sheet thickness change. Another issue that must be taken into consideration is the fact that often only evidence relating to the last glacial advance will be preserved, with evidence relating to earlier fluctuations typically having been destroyed due to the erosive nature of ice.

The task of determining the history of an ice sheet that is still present is more difficult, since any evidence relating to a smaller-than-present ice sheet will be obscured. Such a configuration can be inferred if moraines are truncated by the current ice sheet or if contemporary ice-sheet retreat exposes organic material that has been preserved beneath the ice – such material can be dated to determine when it was overrun by ice (Miller et al., 2012). An alternative approach that should be pursued is the recovery of geological samples from beneath the current ice sheets; a number of techniques (e.g. cosmogenic exposure dating, optically stimulated luminescence dating) can be used to determine how long such samples have been covered by ice. Finally, sampling of ice cores extracted from the ice sheet can provide an indication of past ice thickness, via the analysis of either the gas bubbles preserved in the ice or the isotopic composition of the ice itself (Parrenin et al., 2007).

Due to the sparse nature of ice extent data, numerical ice-sheet models are often used to ‘fill the gaps’ between field constraints, drawing on the physics of ice flow to determine the likely configuration and thickness of past ice sheets (e.g. Simpson et al., 2009; Tarasov et al., 2012; Whitehouse et al., 2012a; Gomez et al., 2013; Briggs et al., 2014; Lecavalier et al., 2014; Gowan et al., 2016; Patton et al., 2017). See Sect. 3.1 and Sect. 4.2.1 for further discussion of the role of ice-sheet modelling within studies of GIA.

2.3.3 Surface deformation data

Global Positioning System (GPS) and Interferometric Synthetic Aperture Radar (InSAR) data provide a snapshot of present-day solid Earth deformation. The full potential of InSAR has yet to be realized in the field of GIA – current studies are limited to Iceland (Auriac et al., 2013) – but there is a long tradition of GPS data being used to constrain GIA models. These data must be corrected for signals associated with the global water cycle, atmospheric effects, and local processes associated with tectonics or sediment compaction (King et al., 2010). In areas where non-GIA signals are well constrained and there is a dense network of measurements, such as across North America (Sella et al., 2007) or Fennoscandia (Lidberg et al., 2007), GPS data have successfully been used to calibrate GIA models (e.g. Milne et al., 2001; Milne et al., 2004; Lidberg et al., 2010; Kierulf et al., 2014; Peltier et al., 2015). However, in regions where contemporary ice mass change also contributes to present-day solid Earth deformation, it becomes difficult to disentangle contributions from past and present ice-sheet change



(Thomas et al., 2011;Niield et al., 2014). Horizontal GPS rates are often more precise than vertical rates by an order of magnitude (King et al., 2010), but before they can be compared with GIA model output the velocity field due to plate motion must be removed. This is non-trivial, since neither plate motion nor GIA are perfectly known (King et al., 2016). Another barrier to the use of horizontal GPS rates in calibrating or validating GIA models is the issue that horizontal deformation in response to surface loading can be strongly perturbed by the presence of lateral variations in Earth rheology (Kaufmann et al., 2005), consequently, horizontal GPS rates have not yet been used to their full potential within the field of GIA.

Geodetic information is typically provided on a reference frame whose origin is located at the centre of mass of the entire Earth system (e.g. ITRF2008; Altamimi et al., 2012) while GIA model predictions are typically provided on a reference frame whose origin lies at the centre of mass of the solid Earth. Reference frame differences must therefore be accounted for when comparing model output with GPS data, along with uncertainties associated with realization of the origin of the reference frame (King et al., 2010).

2.3.4 Gravity data

Between 2002 and 2017, repeat measurements of the Earth's gravity field by the Gravity Recovery and Climate Experiment (GRACE) satellites allowed temporal variations in the distribution of mass throughout the cryosphere, the atmosphere, the oceans, and the solid Earth to be quantified (e.g. Wouters et al., 2014). One of the principle drivers of solid Earth deformation is GIA, and across regions that are currently ice-free, GRACE data (and measurements of the static gravity field by the 'Gravity field and steady-state Ocean Circulation Explorer', GOCE) have been used to quantify the magnitude and spatial pattern of the local GIA signal (e.g. Tamisiea et al., 2007; Hill et al., 2010; Metivier et al., 2016), past ice thickness (Root et al., 2015), and local viscosity structure (Paulson et al., 2007a). However, in areas where an ice sheet is still present variations in the local gravity field will reflect the solid Earth response to both past and present ice mass change, as well as contemporary changes to the mass of the ice sheet itself (Wahr et al., 2000). In this situation, a joint approach to solving for GIA and contemporary ice mass change is necessary, often via the combination of GRACE data with other data sets (see Sect. 3.4) (e.g. Sasgen et al., 2007; Riva et al., 2009; Ivins et al., 2011; Groh et al., 2012; Gunter et al., 2014; Martin-Espanol et al., 2016b).

On a more local scale, absolute gravity measurements have been used to study GIA (e.g. Peltier, 2004; Steffen et al., 2009; Mazzotti et al., 2011; Memin et al., 2011; Sato et al., 2012), while the relationship between surface gravity change and uplift rates can be employed to constrain GIA in regions where ice history and Earth structure are poorly known (e.g. Wahr et al., 1995; Purcell et al., 2011; van Dam et al., 2017).



2.3.5 Independent constraints on solid Earth properties

430 The rheology of the mantle and the thickness of the lithosphere are often inferred by comparing GIA model output with observations that reflect past and present rates of solid Earth deformation, such as GPS time series or records of past relative sea-level change (e.g. Lambeck et al., 1998; Whitehouse et al., 2012b; Argus et al., 2014). However, GIA model predictions will be sensitive to the assumed ice history, and therefore it can be useful to draw on independent information to constrain properties of the solid Earth.

435 For the purposes of GIA, the elastic and density structure of the Earth is assumed to follow that of the PREM (Dziewonski and Anderson, 1981), which is derived from seismic data. Lithosphere thicknesses can be inferred from inversions of gravity or seismic data, or via thermal modelling, although it should be noted that the response of the lithosphere will depend on the timescale of the loading (Watts et al., 2013), and hence values derived by considering e.g. the seismic thickness of the lithosphere, or its elastic thickness over geological timescales, may not be relevant for GIA. Finally, mantle viscosities can
440 be independently estimated via a number of approaches, including consideration of processes associated with mantle convection (e.g. Mitrovica and Forte, 2004), or via the conversion of seismic velocity perturbations into mantle viscosity variations (e.g. Ivins and Sammis, 1995; Wu and van der Wal, 2003; Latychev et al., 2005b; Paulson et al., 2005).

Although inferences of upper mantle viscosity derived from GIA modelling are typically dependent on the assumed ice
445 history, GIA-related processes that are sensitive to lower mantle viscosity are often not strongly dependent on the choice of ice history. The ongoing long wavelength response of the solid Earth to surface mass redistribution since the LGM principally depends on the viscosity of the lower mantle, and results in changes to the oblateness of the solid Earth (J_2), the position of the geocenter, and the orientation of the rotation pole (Figure 4). If these processes can be quantified (e.g. Gross and Vondrak, 1999; Cheng and Tapley, 2004) they can be used to place constraints on lower mantle viscosity (e.g. Paulson et al., 2007b; Mitrovica and Wahr, 2011; Mitrovica et al., 2015). However, it should be noted that these large-scale processes
450 will also be affected by contemporary surface mass redistribution, for example, associated with melting of the polar ice sheets (Adhikari and Ivins, 2016).

2.3.6 Stress field

455 Unloading of the solid Earth during deglaciation alters the regional stress field, and can trigger postglacial faulting (Stewart et al., 2000). However, it is not straightforward to infer past changes in surface loading from the regional faulting history because although deglaciation can trigger faulting, GIA-induced stress changes are probably only capable of triggering slip on pre-existing faults, with the fault expression reflecting the underlying tectonic stress field as well as the GIA-related stress field (Steffen et al., 2012; Craig et al., 2016). Finite-element modelling of glacially-induced faulting indicates that the
460 magnitude of fault slip is primarily governed by fault geometry rather than the magnitude of ice thickness change, although ice-sheet size has been shown to influence the timing of fault activation (Steffen et al., 2014).



2.4 Significant results

Over the past 40 years, GIA modelling has played a central role in advancing our understanding of the rheology of the Earth, the history of the global ice sheets, and the factors controlling spatially-variable sea-level change. Key results are briefly outlined below.

2.4.1 Mantle viscosity

GIA modelling is one of the principle approaches used to determine mantle viscosity. A range of global and regional studies indicate that the mean viscosity of the upper mantle lies in the range 10^{20} - 10^{21} Pa s, while the viscosity of the lower mantle is less tightly constrained to lie in the range 10^{21} - 10^{23} Pa s (e.g. Mitrovica, 1996; Lambeck et al., 1998; Milne et al., 2001; Peltier, 2004; Bradley et al., 2011; Steffen and Wu, 2011; Whitehouse et al., 2012b; Lambeck et al., 2014; Lecavalier et al., 2014; Peltier et al., 2015; Nakada et al., 2016). GIA modelling can be used to solve for the depth-dependent viscosity profile of the mantle, but it is important to assess the resolving power of the constraining data sets when considering the accuracy and uniqueness of the results (Mitrovica and Peltier, 1991a; Milne et al., 2004; Paulson et al., 2007b). Finally, GIA modelling has been used to identify a number of localized low viscosity regions ($<10^{20}$ Pa s), e.g. in Iceland (Pagli et al., 2007; Auriac et al., 2013), Alaska (Larsen et al., 2005; Sato et al., 2011), Patagonia (Ivins and James, 2004; Lange et al., 2014), and the Antarctic Peninsula (Simms et al., 2012; Niold et al., 2014).

2.4.2 Ice sheet change

The first-order time series of ice volume change since the LGM – thought to be equivalent to ~130 m sea-level rise - has been reconstructed via the comparison of global sea-level records with GIA model output (e.g. Lambeck et al., 2014), but small magnitude ice volume changes subsequent to the demise of the Laurentide and Fennoscandian ice sheets around 7 ka are less well constrained (Lambeck et al., 2014; Bradley et al., 2016). Combining GIA modelling with a probabilistic approach, Kopp et al. (2009) find that global ice volumes during the Last Interglacial (~125 ka) were at least 6.6 m smaller than present (95% probably; magnitude expressed as sea-level equivalent), but uncertainty associated with the interpretation and dating of sea-level indicators (Düsterhus et al., 2016b; Rovere et al., 2016) and neglect of non-GIA processes (Austermann et al., 2017) still hampers our ability to precisely reconstruct changes in global ice volume during this period. In order to reconstruct global ice volumes over longer timescales, the complicating effects of tectonics, dynamic topography, and sediment compaction must be accounted for (Rovere et al., 2014), although Raymo et al. (2011) have successfully demonstrated that the scatter in Pliocene shoreline elevations (typically found between 10 and 40 m above present sea level) can be explained once GIA processes are taken into account.

In addition to constraining global ice volumes, comparison of GIA model output with a range of data sets has been used to reconstruct the past configuration of individual ice sheets, including the Fennoscandian (Lambeck et al., 1998; Lambeck et



495 al., 2010), British-Irish (Peltier et al., 2002;Bradley et al., 2011), Laurentide (Tarasov et al., 2012;Simon et al.,
2016;Lambeck et al., 2017), Greenland (Tarasov and Peltier, 2002;Simpson et al., 2009;Lecavalier et al., 2014), and
Antarctic (Whitehouse et al., 2012a;Whitehouse et al., 2012b;Gomez et al., 2013;Ivins et al., 2013;Argus et al., 2014;Briggs
et al., 2014) ice sheets. Global reconstructions also exist (Peltier, 2004;Peltier et al., 2015), but the important question of
500 unresolved (Clark and Tarasov, 2014).

Finally, improved quantification of the geodetic signal associated with past ice-sheet change has led to recent improvements
in the accuracy of contemporary estimates of ice mass balance, as derived from GRACE or altimetry data (King et al.,
2012;Ivins et al., 2013). However, uncertainty associated with the ‘GIA correction’ that must be applied to such data sets
505 still poses a significant barrier to reaching a fully-reconciled estimate of contemporary ice mass balance (Shepherd et al.,
2012).

2.4.3 Sea-level change

Understanding the rate, magnitude, and spatial pattern of past, present and future sea-level change, and linking these changes
510 to climate forcing, is one of the most important questions facing modern society (IPCC, 2013). Important results that have
been derived using GIA modelling include:

- i) quantification of the maximum rate of global mean sea-level rise during the last deglaciation (e.g. Lambeck et al., 2014)
- ii) identification of the potential meltwater source(s) that contributed to rapid sea-level rise during the last deglaciation
(e.g. Clark et al., 2002;Gomez et al., 2015a;Liu et al., 2016)
- 515 iii) quantification of the maximum sea-level attained during past warm periods (e.g. Kopp et al., 2009;Dutton et al., 2015)
- iv) identification of the rate, pattern, and source of historical and contemporary sea-level change (e.g. Riva et al., 2010;Hay
et al., 2015;Rietbroek et al., 2016)
- v) quantification of the likely pattern of future sea-level change due to ice-sheet change (e.g. Mitrovica et al.,
2009;Slangen et al., 2014)

520 All of these results draw on the complex relationship between ice sheet-change and spatially-variable sea-level change, as
described by the sea-level equation. Significant advances in our understanding of sea-level and ice-sheet change have come
about due to improvements in data availability and GIA modelling capability during the last decade, but persistent
uncertainties associated with the GIA correction that must be applied when interpreting gravity, altimetry, tide gauge, or
GPS data (Tamisiea, 2011), and ongoing ambiguity associated with the interpretation of palaeo-data, mean that future
525 progress will require input from a diverse range of disciplines.



3 Recent Developments

Over the past decade there have been rapid advances in our understanding of how GIA processes can influence other dynamic systems, and an increased awareness of additional factors that must be considered when seeking to constrain or tune a GIA model using independent datasets. New approaches of isolating the GIA signal have also been devised. Four of the most important recent advances are briefly described in this section.

3.1 Ice dynamic feedbacks

Inferring the past evolution of the major ice sheets has been a central goal of GIA modelling since Jamieson (1865) first observed that the growth of an ice sheet will depress the land and affect the position of the ocean shoreline. However, it is only recently that glaciologically-consistent ice-sheet reconstructions, i.e. those developed using a numerical ice-sheet model, have begun to be produced for the purposes of GIA modelling (e.g. Tarasov and Peltier, 2002; Tarasov et al., 2012; Whitehouse et al., 2012a). A crucial boundary condition that must be defined when modelling the evolution of a marine-grounded ice sheet is the water depth of the surrounding ocean. This water depth determines where the ice sheet begins to float, a point known as the grounding line. More importantly, it determines the rate at which ice flows across the grounding line and into the ocean (because ice flux depends on ice thickness (Schoof, 2007)).

Numerical ice-sheet models are typically run assuming that sea-level change adjacent to an ice sheet will track global mean sea-level change. Far-field ice melt will indeed cause near-field sea-level rise (Figure 6a), but, due to the effects of GIA, water depth changes will not follow the global mean during near-field ice-sheet change (Figure 2). Nearly 40 years ago, Greischar and Bentley (1980) noted that solid Earth rebound triggered by ice loss from a marine-grounded ice sheet would reduce local water depths, and could promote grounding line advance (Figure 6b). The decreased gravitational attraction of the melting ice sheet also acts to reduce local water depths. Modelling both effects, Gomez et al. (2010) demonstrated that GIA has a stabilizing effect on the dynamics of a marine-grounded ice sheet, and can prevent or delay unstable grounding line retreat and ice loss.

Spatially-variable water depth boundary conditions were first used in conjunction with a numerical ice-sheet model, for the purposes of reconstructing past ice-sheet change, by Whitehouse et al. (2012a), who used a priori GIA model output to determine water depths around Antarctica, and hence ice flux across prescribed grounding line positions. Subsequently, Gomez et al. (2013) and de Boer et al. (2014) and have used fully-coupled ice sheet-GIA models to produce ice-sheet reconstructions that are consistent with spatially-variable sea-level change over time. It is interesting to note that LGM reconstructions for Antarctica generated using coupled models tend to contain 1-2 m less ice (expressed as sea-level equivalent) than reconstructions generated using uncoupled models (de Boer et al., 2017). If the coupled model results are robust this makes it difficult to account for the global mean sea-level lowstand during the LGM (Clark and Tarasov, 2014).



560 Considering future ice-sheet change, Adhikari et al. (2014) have used one-way coupling to quantify the impact of ongoing
GIA on Antarctic ice dynamics up to 2500 AD, while Gomez et al. (2015b) and Konrad et al. (2015) have used coupled
models to investigate the long-term evolution of the ice sheet, finding that GIA-related feedbacks have the potential to
significantly limit, or even halt, future ice loss if the upper mantle viscosity beneath West Antarctica is low enough for rapid
rebound to be triggered. A crucial factor in determining the stability of an ice sheet is the resistance provided by the
surrounding floating ice shelves. If rebound is fast enough for the ice shelves to re-ground on submerged topographic highs,
565 forming ice rises (Matsuoka et al., 2015), this significantly increases the chance of ice-sheet stabilization or even re-growth
(Kingslake et al., in review). Uncertainties associated with the bathymetry and the upper mantle viscosity beneath the
Antarctic and Greenland ice sheets currently present the greatest barriers to accurately quantifying the degree to which GIA-
related feedbacks have the potential to limit future ice loss from these regions.

3.2 Lateral variations in Earth rheology, and non-linear rheology

570 GIA models traditionally assume the Earth behaves as a linear Maxwell viscoelastic solid with a viscosity profile that varies
in the radial direction only and stays constant with time (e.g. Peltier et al., 2015). A number of studies have made use of a bi-
viscous Burgers rheology within a radially-varying framework, in which mantle deformation is dominated by the behaviour
of two viscosities linearly relaxing over different timescales (Yuen et al., 1986; Caron et al., 2017). However, an increasing
number of studies are making use of a framework that can accommodate three-dimensional variations in mantle viscosity
575 (e.g. Ivins and Sammis, 1995; Martinec, 2000; Kaufmann et al., 2005; Latychev et al., 2005b; Paulson et al., 2005; Steffen et
al., 2006; A. et al., 2013), possibly defined via use of a non-linear creep law – where the viscosity depends on the time-
varying stress field (Wu et al., 2005; van der Wal et al., 2013). The effect of including plate boundaries – which affect the
horizontal transmission of stress – and variations in the thickness and rheology of the lithosphere have also been explored
(Latychev et al., 2005a; Kuchar and Milne, 2015). The development of these ‘3D GIA models’ are motivated by (i)
580 convincing evidence for strong lateral variations in rheological properties beneath some regions, including Antarctica
(Heeszel et al., 2016); and (ii) the demonstration that consideration of lateral variations in rheology is required to correctly
model horizontal deformation (Kaufmann et al., 2005).

It is important to question whether such increased model complexity is necessary. Whitehouse et al. (2006) showed that
585 inclusion of 3D Earth structure perturbs uplift rate predictions across Fennoscandia by an amount greater than current GPS
accuracy, with significant implications for inferences of past ice-sheet history, while Kendall et al. (2006) demonstrated that
relative sea-level change predictions will be biased by >0.2 mm/yr at ~ 150 global tide gauge sites if 3D Earth structure is
neglected, with maximum differences exceeding several mm/yr. Since solid Earth deformation depends on both the surface
load history and Earth rheology, non-uniqueness is a problem when solving for these two unknowns. If a GIA model is tuned
590 to fit GIA-related observations (e.g. uplift rates, sea-level records) without accounting for lateral variations in rheology, the



resulting ice-sheet reconstruction is likely to be biased. For example, past ice thickness change is likely to be overestimated in regions where the local mantle viscosity is weaker than assumed by the model (Figure 5b). Similarly, global ice volumes may be incorrectly inferred if viscosity variations are ignored at far-field sea-level sites (Austermann et al., 2013). If the past ice history of a region has been independently determined then neglect of lateral variations in Earth structure will lead to bias in predictions of the GIA signal, and hence bias in estimates of contemporary ice sheet mass balance, potentially on the order of 10's Gt/yr (van der Wal et al., 2015). Furthermore, models that consider the coupled evolution of the ice sheet-solid Earth system (see Sect. 3.1) will be highly sensitive to the underlying viscosity field (Gomez et al., 2015b; Konrad et al., 2015; Pollard et al., 2017).

A range of approaches are used to define spatial variations in Earth rheology, but most rely on deriving a temperature field from a seismic velocity model (e.g. Ritsema et al., 2011), from which a viscosity field is derived (e.g. Ivins and Sammis, 1995; Latychev et al., 2005b). This derivation is not straightforward, and in particular, compositional effects must be accounted for when interpreting seismic velocity perturbations in terms of temperature perturbations (Wu et al., 2013). If a power-law approach is used, grain-scale deformation of mantle material is described by diffusion and dislocation creep through the use of a non-linear relationship where the strain rate depends on stress to some power. This power is thought to be 1 for diffusion creep – i.e. a linear response to forcing – but ~3.5 for dislocation creep (Hirth and Kohlstedt, 2003). This has implications for the inferred viscosity of the mantle: if dislocation creep is important, i.e. there is a non-linear relationship between stress and strain rate, then the effective viscosity of the mantle will depend on the von Mises stress. Since the von Mises stress depends on the evolution of the ice/ocean surface load it follows that the effective viscosity will be time dependent. Inputs to the power-law relationship include grain size, water content, and melt content, as well as temperature, although a lack of direct observational data for most of these parameters mean that values derived in laboratory experiments are often adopted (Hirth and Kohlstedt, 2003; Burgmann and Dresen, 2008; King, 2016). Significant further work is needed to better quantify the viscosity distribution throughout the mantle, and to determine the spatial resolution at which viscosity variations must be resolved to accurately reflect the global GIA process (Steffen and Wu, 2014).

3.3 Sedimentary isostasy

The isostatic response to sediment erosion and deposition, on glacial and longer timescales, has long been considered in studies of onshore and offshore crustal deformation, associated with both fluvial and glacial systems. However, the impact of sediment redistribution on Earth's gravitational and rotational fields, and the consequent effect on sea-level, has only recently been considered. Dalca et al. (2013) were the first to incorporate the gravitational, deformational, and rotational effects of sediment redistribution into a traditional GIA model (Figure 7a). The resulting theory has been used to demonstrate that the impact of sediment erosion and deposition, associated with both fluvial and glacial systems, can alter relative sea-level by several metres over the course of a glacial cycle and rates of present-day deformation by a few tenths of a mm/yr (Wolstencroft et al., 2014; Ferrier et al., 2015; Kuchar et al., 2017; van der Wal and IJpelaar, 2017). Although the



625 magnitude of the perturbation due to sediment loading is small, it is greater than the precision of modern geodetic methods,
and hence has the potential to bias contemporary estimates of sea-level change (Ferrier et al., 2015; van der Wal and IJpelaar,
2017). Perhaps the most important finding of these preliminary studies is the observation that in order for relative sea-level
indicators to be used to constrain past global ice volumes, they must first be corrected for the effects of both glacial and
sedimentary isostasy (Ferrier et al., 2015). As an example, if sediment loading has caused a sea-level indicator to subside
subsequent to its formation, this will lead to an overestimation of the magnitude of sea-level rise since the formation of the
630 sea-level indicator.

Sedimentary isostasy is not the only sediment-related process that affects sea level. Wolstencroft et al. (2014) and Kuchar et
al. (2017) both found that including the effects of sedimentary isostasy did not bring agreement between model predictions
and GPS-derived observations of contemporary land motion around the Mississippi Delta, and they concluded that sediment
635 compaction must play a significant role (Figure 7b). To address this Ferrier et al. (2017) have updated the theory developed
by Dalca et al. (2013) so that it accounts for all the competing processes associated with sediment redistribution, including
the decrease in water depth due to offshore deposition and the increase in water depth due to subsidence and compaction
(Figure 7). This state-of-the-art approach, which uses estimates of sediment porosity and saturation to determine the time-
evolving effects of compaction, has recently been used to provide a robust interpretation of sea-level indicators formed
640 during Marine Isotope Stage 3 (~50-37 ka) in the region of China's Yellow River Delta, and hence tighten constraints on
global ice volumes at this time (Pico et al., 2016).

In conclusion, both sediment loading and sediment compaction have been demonstrated to have a non-negligible effect on
relative sea-level change, and this has far-reaching implications for the interpretation of relative sea-level indicators within
645 the framework of a GIA model (see Sect. 2.3.1). Both effects should be accounted for in future GIA models, although better
constraints on the timing and distribution of erosion and deposition, over the last glacial cycle and beyond, are needed before
past global ice volumes can be robustly inferred from global sea-level records.

3.4 Inverse solutions

The primary motivation for the recent focus on developing GIA models for Greenland and Antarctica has been to permit
650 accurate interpretation of GRACE data, partitioning the mass change time series into contributions from GIA and
contemporary ice mass change (King et al., 2012; Ivins et al., 2013). However, GRACE data can also be used to determine an
estimate of the GIA signal by combining it with altimetry data. The basic premise of this method is that the GRACE
satellites detect total mass change, which is attributed to the redistribution of ice and solid Earth mass (after accounting for
atmospheric and oceanic mass change), whereas altimetry satellites such as ICESat measure surface elevation change, which
655 is attributed to a combination of solid Earth deformation and ice volume change. A density model is used to convert the
ICESat height measurements to mass, and the two datasets can be used to solve for GIA-related deformation, as shown in the



following equation, where \dot{h}_{GIA} is the rate of surface elevation change due to GIA, \dot{h}_{ICESat} is the rate of surface elevation change measured by ICESat, \dot{m}_{GRACE} is the rate of total mass change measured by GRACE, ρ_{rock} is the mean density of the upper ~100km of the solid Earth, and ρ_{surf} is the mean density of the ice volume change:

660

$$\dot{h}_{GIA} = \frac{\dot{m}_{GRACE} - \rho_{surf} \cdot \dot{h}_{ICESat}}{\rho_{rock} - \rho_{surf}} \quad (3)$$

665

This method of simultaneously solving for ice mass change and GIA was originally suggested by Wahr et al. (2000). Riva et al. (2009) first implemented it for Antarctica using 5 years of data, and Gunter et al. (2014) improved the method by using a firn densification model instead of assuming a constant density of ice. Further advances were made by Martin-Espanol et al. (2016b), who estimated the GIA signal across Antarctica using a statistical inversion of a suite of geodetic observations, including GRACE, altimetry, InSAR, and GPS data, and a priori information about the wavelength of the GIA signal, while Sasgen et al. (2017) extended the method to account for lateral variations in Earth structure. The advantage of using an inverse approach over forward models of GIA is that the solution is not dependent on the ice loading history or the Earth structure (other than the assumed density of the Earth). Using the GRACE data in a slightly different way, Sasgen et al. (2013) developed a method that systematically searches the forward model parameter space to determine an optimal fit to the GRACE data, using GPS data to provide a further constraint on deformation rates. These data-driven GIA solutions currently exhibit significant variability (Martin-Espanol et al., 2016a), partly due to the large number of assumptions that must be made during processing of the raw data, but they have formal uncertainties attached to them, and hence provide a useful comparison for output derived from process-based forward models of GIA.

670

675

4 Future directions and challenges

In this final section a number of emerging areas of research are outlined. Some of the most exciting future developments will come as GIA modelling becomes even better integrated with a range of disciplines.

4.1 Data interpretation and assimilation

680

A central component of GIA modelling has always been the use of independent datasets to either prescribe model inputs, tune model parameters, or test model outputs. As the accuracy and coverage of these datasets improves, along with our understanding of the processes they record, it is important to be aware that it will not be possible to honour all of the constraints provided by the data. A strategy is therefore needed in which the available data are used to develop the most accurate GIA model possible and assign a level of uncertainty to the resulting model output.

685



690 Many of the datasets that record GIA processes also record competing processes. Around 50% of contemporary global mean sea-level rise is due to thermosteric effects, reflecting a change in the density of the ocean rather than its mass (IPCC, 2013); the thermosteric contribution to post-LGM global mean sea-level rise has not been quantified. Changes in oceanic or atmospheric circulation will alter the dynamic topography of the ocean surface, while changes in the tidal regime or ocean conditions (acidity, temperature, opacity) can alter the habitat distribution of species that are used to infer the position of the shoreline. When seeking to reconstruct past ice extent it is important to acknowledge that a change in local ice flow or wind direction can alter the height of the ice-rock contact, while changes in ice surface elevation, as recorded by ice cores, will reflect changes in both ice thickness and the height of the underlying bed (Bradley et al., 2012; Lecavalier et al., 2013). Finally, land motion, as recorded by GPS, will reflect the solid Earth response to contemporary surface mass redistribution (ice, water, sediment) as well as tectonics and sediment compaction. Over longer timescales (≥ 100 kyr) land motion also reflects the dynamic response of the lithosphere to mantle convection (Rowley et al., 2013; Austermann et al., 2017). As far as is possible, these competing factors must be accounted for, or appropriate error bars should be attached to the data in a format that can be easily incorporated into a GIA modelling framework (Düsterhus et al., 2016a).

700 Once a suitable data set has been identified, it can be used to determine an optimal GIA model. Due to data gaps and uncertainties there are often trade-offs between the magnitude and timing of the surface loading history and Earth rheology that can explain the observations (Figure 5). Traditionally, an iterative approach has been used to determine a single, optimal ice history/Earth model combination that provides the best fit to a range of data types (Lambeck et al., 1998; Peltier et al., 2015). Using data assimilation or statistical emulation, in combination with large ensemble modelling, it should be possible to determine a suite of GIA solutions that provide a reasonable fit to all available constraints. Such an approach has already been pioneered for ice-sheet modelling (Briggs et al., 2013; Pollard et al., 2016), but it requires careful consideration of the probability distribution associated with each piece of constraining data and the method used to score each model run (Briggs and Tarasov, 2013). When applied to GIA modelling, decisions will have to be made on: how to weight different data types, the length scale over which each is relevant, how to treat uncertainties in age and elevation, whether to use raw data or statistical reconstructions (e.g. Khan et al., 2015), and how to account for additional metadata, such as information on whether sea-level was rising or falling at a particular location. The uncertainty on the resulting GIA estimate should directly reflect uncertainties in the constraining data.

715 In order to optimize the search through the parameter space associated with ice history and Earth rheology a number of approaches could be taken. When determining the past evolution of the global ice sheets, a previously-used method involves taking an existing global reconstruction and scaling the thickness of each ice sheet in turn (e.g. Caron et al., 2017). However, building on new understanding of the feedbacks between ice dynamics and GIA (Sect. 3.1), completely new glacial scenarios should also be explored, guided by the output from coupled models. When determining the optimal Earth model, it will be



720 important to allow for the possibility that different regions are characterized by different viscosity profiles, guided by independent constraints on mantle viscosity variations (Sect. 2.3.5).

725 There will always be some parts of the parameter space that cannot be constrained. A useful exercise is to determine the locations in which it would be most beneficial to have new data constraints (e.g. Wu et al., 2010), but there are often logistical barriers to collecting the most useful data. For example, it would be very useful to know the rate of solid Earth rebound beneath the present-day ice sheets and across the ocean floor; it would be useful to know the past thickening history of the ice sheets as well as their thinning history; and it would be useful to know how sea level has changed at locations distal from current and past shorelines. In the near future, improvements in data coverage will come through the application of novel analytical techniques in regions where sea-level reconstruction have so far proved challenging, e.g. mangroves, and the more widespread use of gravity data to constrain the GIA signal where sea-level and GPS data are lacking (Root et al., 730 2015).

4.2 Coupled modelling

735 Significant advances in understanding GIA have often stemmed from a cross-disciplinary approach. The gravitational theory developed by the mathematician Woodward in the late 19th century came about as the result of a question posed by geologists Gilbert and Chamberlin (Woodward, 1888). More recently, observations of a mid-Holocene highstand throughout the low latitudes led to recognition of the ocean syphoning process (Mitrovica and Milne, 2002), while disparate observations relating to the magnitude of the Pliocene highstand led to advances in the modelling of GIA over multiple glacial cycles (Raymo et al., 2011). Sea-level observations, and more recently GPS observations, have often been the motivation for developing new hypotheses relating to the history of the major ice sheets (Bradley et al., 2015) and the processes governing ice dynamics (Pollard et al., 2015). Future progress is likely to be made by fully integrating a number of 740 different disciplines, i.e. via coupled modelling.

4.2.1 Coupled ice sheet-3D GIA modelling

745 Coupled ice sheet-GIA modelling has already been discussed in Sect. 3.1, as has 3D GIA modelling (Sect. 3.2), but since the impact of GIA on ice dynamics has been shown to be strongest in locations where mantle viscosity is unusually low (Pollard et al., 2017) it will be crucial to incorporate lateral variations in Earth structure into future coupled ice sheet-GIA models in order to accurately predict the evolution of the many ice sheets that are located above low viscosity regions (e.g. West Antarctica, Patagonia, Alaska, Iceland; see Sect. 4.3). This will be computationally challenging and careful experiment design will be needed to ensure an efficient, yet sufficient, search of the parameter space.



750 4.2.2 Coupled GIA-ocean/atmosphere modelling

The impacts of GIA, i.e. spatially variable perturbations to the sea surface and solid Earth, are not currently considered within atmosphere or ocean circulation models, but are likely to have a non-negligible effect on these systems. Perturbations to the seafloor due to the growth and decay of submerged peripheral bulges (Figure 2b) have been shown to be sufficient to perturb ocean circulation (Rugenstein et al., 2014), while the spatially-variable sea-level change that is predicted to accompany ice melt will impact the global tidal regime (Wilmes et al., 2017). Similarly, changes in surface topography due to solid Earth deformation will affect atmospheric circulation patterns and consequently precipitation patterns. This latter factor has clear implications for the mass balance of an ice sheet, and has been proposed as an explanation for hysteresis within glacial cycles (Abe-Ouchi et al., 2013). During the LGM the areal extent of the continents will have been greater as a result of GIA. Additional land bridges will have existed, with direct implications for migration pathways and ocean circulation, and in the northern hemisphere the additional land mass may have facilitated ice-sheet expansion or inception. GIA is also able to provide insight into the timing, magnitude and source of freshwater inputs to the ocean during deglaciation. In order to accurately model the palaeo-circulation of the ocean and the atmosphere, some factors associated with GIA should be incorporated into future modelling efforts.

765 4.2.3 Coupled GIA-surface process modelling

GIA-related solid Earth deformation during the last glacial cycle will have affected the routing of palaeo-drainage systems (Wickert et al., 2013), it will have governed the location and extent of ice-dammed lakes (Lambeck et al., 2010; Lambeck et al., 2017; Patton et al., 2017), and it continues to influence coastal evolution (Whitehouse et al., 2007). Accounting for GIA-related factors such as changes to the shape of the land surface and changes to the position of the base level within landscape evolution models will lead to a more complete understanding of Earth surface processes. The full impacts of GIA, including spatially-variable sea-level change and changes to the shape of Earth's gravitational field, should be accounted for in future studies that seek to understand the isostatic response to glaciation, erosion, and sedimentation (e.g. Mey et al., 2016; Moucha and Ruetenik, 2017).

775 4.2.4 GIA-Climate feedbacks

GIA potentially plays a role in modulating climate cycles. Postglacial rebound acts to reduce the pressure in the mantle, and this has been implicated in promoting terrestrial volcanism (Sigmundsson et al., 2010; Praetorius et al., 2016). Huybers and Langmuir (2009) argue that the CO₂ release associated with increased volcanism during the last deglaciation may have been sufficient to promote further ice melt, raising the possibility that glacial rebound, CO₂ release, and ice dynamics are part of a positive feedback loop. But postglacial rebound is not the only GIA-related process that affects the rate of CO₂ release: as the ice sheets wax and wane this alters global mean sea level, and the resulting pressure changes at the seafloor are thought to be sufficient to perturb rates of decompression melting and submarine volcanism at mid-ocean ridges. It was originally assumed that mid-ocean ridge volcanism would be suppressed during periods of sea-level rise (i.e. during deglaciation) and this would



act to counter the contemporaneous climatic effects of increased terrestrial volcanism (Huybers and Langmuir, 2009).
785 However, the evidence suggests that there is actually a significant delay in the response of the mid-ocean ridge system to a
change in seafloor pressure. This can be deduced from records of hydrothermal and magmatic activity along mid-ocean
ridges (Crowley et al., 2015;Lund et al., 2016), while models of melt transport through the mantle predict a lag of at least
60 kyr between a fall in sea-level and an increase in CO₂ emissions (Burley and Katz, 2015). The magnitude of the lag
790 depends on a number of factors, including the plate spreading rate and the rate of sea-level change, so it is not a simple task
to quantify the time-dependent net effect of terrestrial and marine volcanic processes on atmospheric CO₂. But tantalizingly,
Huybers and Langmuir (2017) have been able to reproduce ~100 kyr glacial cycles by assuming a lag of 50 kyr and
considering feedbacks between deglaciation, sea-level rise, volcanism, and CO₂ emissions, allowing for the temperature
dependence of CO₂ degassing from the ocean. The CO₂ perturbations described here are triggered by GIA processes, and
there is clear scope for further exploration of the feedbacks between the spatially-variable isostatic response to surface load
795 changes, including changes in sediment loading (Sternai et al., 2016), volcanism, climate change, and ice dynamics.

4.3 Low viscosity regions

A number of glaciated regions - Alaska, Iceland, the northern Antarctic Peninsula, and Patagonia – are situated on active or
recently-active plate boundaries. The high mantle temperatures associated with such tectonic settings mean that the viscosity
of the upper mantle is likely to be very low in these regions, typically $\leq 10^{19}$ Pa s (e.g. Sato et al., 2011;Auriac et al.,
800 2013;Lange et al., 2014;Niield et al., 2014), and hence the relaxation time of the mantle will be short, on the order of years to
decades. This has two important implications: (i) the viscous response to any surface mass change prior to a few thousand
years ago will have decayed, and instead (ii) viscous deformation will be dominated by the response to recent, or even
contemporary, ice mass change.

805 Glaciated low viscosity regions are exciting places to study GIA, but the issues listed above complicate attempts to separate
the geodetic signal into a response to past and contemporary change. It is typically assumed that GIA is a linear background
signal that reflects the viscous response to long-past ice-sheet change and any departure from the linear trend reflects the
elastic response to contemporary ice-sheet change. However, in a low viscosity region the short relaxation time of the mantle
means that the solid Earth response to historical ice mass change, i.e. the GIA signal, may not be linear over the epoch of
810 geodetic measurements, making it difficult to isolate. Furthermore, the geodetic response to contemporary ice mass change
may contain both an elastic and a viscous signal (Niield et al., 2014). The problem becomes more tractable if the viscosity of
the mantle can be absolutely determined, for example, via careful analysis of GPS time series in response to known surface
mass change, permitting more reliable predictions of the time-varying elastic and viscous components of deformation.

815 The low viscosity values inferred from GIA for regions such as Iceland and Alaska are similar to the values determined in
studies of post-seismic deformation (Arnadottir et al., 2005;Johnson et al., 2009), and it has been suggested that power-law



820 flow may sufficiently describe both processes due to the similarity of the timescales over which they take place (James et al.,
2009). There is the additional complication that afterslip must be accounted for in post-seismic studies (Ingleby and Wright,
2017), but in general, the changing deformation rates observed during an earthquake cycle suggest that the Earth either
825 follows a power-law rheology (Freed and Burgmann, 2004; Freed et al., 2006), or a rheology comprising several different
relaxation times (Pollitz, 2005; Hetland and Hager, 2006). Is there a single rheological law that can explain GIA, post-seismic
deformation, intra-plate deformation, and deformation in response to sediment or lake loading (Gilbert, 1890; Dickinson et
al., 2016)? It is clear that the Earth behaves differently over different timescales (Burgmann and Dresen, 2008; Watts et al.,
2013), and it will be interesting to see how parallel fields of research progress towards quantifying the rheological structure
of the solid Earth, and the degree to which the rheology reflects the forcing that is applied.

5 Conclusions

830 The ideas developed by Jamieson (1865), Croll (1875), Woodward (1888), Nansen (1921), and (Daly, 1925) laid the
foundations for the development of the sea-level equation (Farrell and Clark, 1976), which provides a robust description of
the spatially-variable sea-level change and solid Earth deformation that takes place over the course of a glacial cycle, and
835 forms the central component of any GIA model. Over the past 40 years, GIA modelling has been instrumental in providing a
joined-up framework for reconstructing the response of global ice sheets to past climatic forcing. With the rapid
development of cross-disciplinary science in the last two decades the field has expanded beyond geodynamics, incorporating
important developments from geodesy, glaciology, and seismology, whilst embracing new results from the fields of geology
and geomorphology. Current predictions for the global GIA signal differ, and future research will revolve around identifying
the reasons for these differences, and incorporating new data constraints and dynamic processes into the modelling.

Author Contribution

PLW wrote the manuscript. Grace Nield is acknowledged below for her assistance in drafting the figures.

Competing interests

The author declares that they have no conflict of interest.

840 Acknowledgements

Grace Nield is warmly acknowledged for her help in producing the figures and for providing invaluable feedback during the
writing of the manuscript. PLW is supported by a UK Natural Environment Research Council (NERC) Independent
Research Fellowship (NE/K009958/1). This article is a contribution to the SCAR SERCE program.



References

- 845 A., G., Wahr, J., and Zhong, S. J.: Computations of the viscoelastic response of a 3-D compressible Earth to surface loading: an application to Glacial Isostatic Adjustment in Antarctica and Canada, *Geophys J Int*, 192, 557-572, Doi 10.1093/Gji/Ggs030, 2013.
- Abe-Ouchi, A., Saito, F., Kawamura, K., Raymo, M. E., Okuno, J., Takahashi, K., and Blatter, H.: Insolation-driven 100,000-year glacial cycles and hysteresis of ice-sheet volume, *Nature*, 500, 190-193, 10.1038/nature12374, 2013.
- 850 Adhikari, S., Ivins, E. R., Larour, E., Seroussi, H., Morlighem, M., and Nowicki, S.: Future Antarctic bed topography and its implications for ice sheet dynamics, *Solid Earth*, 5, 569-584, 10.5194/se-5-569-2014, 2014.
- Adhikari, S., and Ivins, E. R.: Climate-driven polar motion: 2003-2015, *Science Advances*, 2, e1501693, 10.1126/sciadv.1501693, 2016.
- Adhikari, S., Ivins, E. R., and Larour, E.: ISSM-SESAWv1.0: mesh-based computation of gravitationally consistent sea-
855 level and geodetic signatures caused by cryosphere and climate driven mass change, *Geosci Model Dev*, 9, 1087-1109, 10.5194/gmd-9-1087-2016, 2016.
- Al-Attar, D., and Tromp, J.: Sensitivity kernels for viscoelastic loading based on adjoint methods, *Geophys J Int*, 196, 34-77, 10.1093/gji/ggt395, 2014.
- Altamimi, Z., Metivier, L., and Collilieux, X.: ITRF2008 plate motion model, *Journal of Geophysical Research: Solid Earth*,
860 117, B07402, 10.1029/2011jb008930, 2012.
- Argus, D. F., Peltier, W. R., Drummond, R., and Moore, A. W.: The Antarctica component of postglacial rebound model ICE-6G_C (VM5a) based on GPS positioning, exposure age dating of ice thicknesses, and relative sea level histories, *Geophys J Int*, 198, 537-563, 10.1093/Gji/Ggu140, 2014.
- Arnadottir, T., Jonsson, S., Pollitz, F. F., Jiang, W. P., and Feigl, K. L.: Postseismic deformation following the June 2000
865 earthquake sequence in the south Iceland seismic zone, *Journal of Geophysical Research: Solid Earth*, 110, B12308, 10.1029/2005jb003701, 2005.
- Auriac, A., Spaans, K. H., Sigmundsson, F., Hooper, A., Schmidt, P., and Lund, B.: Iceland rising: Solid Earth response to ice retreat inferred from satellite radar interferometry and visocelastic modeling, *Journal of Geophysical Research: Solid Earth*, 118, 1331-1344, 10.1002/jgrb.50082, 2013.
- 870 Austermann, J., Mitrovica, J. X., Latychev, K., and Milne, G. A.: Barbados-based estimate of ice volume at Last Glacial Maximum affected by subducted plate, *Nat Geosci*, 6, 553-557, Doi 10.1038/Ngeo1859, 2013.
- Austermann, J., Mitrovica, J. X., Huybers, P., and Rovere, A.: Detection of a dynamic topography signal in last interglacial sea-level records, *Science Advances*, 3, e1700457, 10.1126/sciadv.1700457, 2017.
- Ballantyne, C. K.: Extent and deglacial chronology of the last British-Irish Ice Sheet: implications of exposure dating using
875 cosmogenic isotopes, *J. Quat. Sci.*, 25, 515-534, 10.1002/jqs.1310, 2010.



- Barlow, N. L. M., Shennan, I., Long, A. I., Gehrels, W. R., Saher, M. H., Woodroffe, S. A., and Hillier, C.: Salt marshes as late Holocene tide gauges, *Global and Planetary Change*, 106, 90-110, [10.1016/j.gloplacha.2013.03.003](https://doi.org/10.1016/j.gloplacha.2013.03.003), 2013.
- Barrell, J.: The status of the theory of isostasy, *American Journal of Science*, 48, 291-338, 1919.
- Bentley, M. J., Cofaigh, C. O., Anderson, J. B., Conway, H., Davies, B., Graham, A. G. C., Hillenbrand, C. D., Hodgson, D. A., Jamieson, S. S. R., Larter, R. D., Mackintosh, A., Smith, J. A., Verleyen, E., Ackert, R. P., Bart, P. J., Berg, S., Brunstein, D., Canals, M., Colhoun, E. A., Crosta, X., Dickens, W. A., Domack, E., Dowdeswell, J. A., Dunbar, R., Ehrmann, W., Evans, J., Favier, V., Fink, D., Fogwill, C. J., Glasser, N. F., Gohl, K., Gollidge, N. R., Goodwin, I., Gore, D. B., Greenwood, S. L., Hall, B. L., Hall, K., Hedding, D. W., Hein, A. S., Hocking, E. P., Jakobsson, M., Johnson, J. S., Jomelli, V., Jones, R. S., Klages, J. P., Kristoffersen, Y., Kuhn, G., Leventer, A., Licht, K., Lilly, K., Lindow, J., Livingstone, S. J., Masse, G., McGlone, M. S., McKay, R. M., Melles, M., Miura, H., Mulvaney, R., Nel, W., Nitsche, F. O., O'Brien, P. E., Post, A. L., Roberts, S. J., Saunders, K. M., Selkirk, P. M., Simms, A. R., Spiegel, C., Stollendorf, T. D., Sugden, D. E., van der Putten, N., van Ommen, T., Verfaillie, D., Vyverman, W., Wagner, B., White, D. A., Witus, A. E., Zwartz, D., and Consortium, R.: A community-based geological reconstruction of Antarctic Ice Sheet deglaciation since the Last Glacial Maximum, *Quat. Sci. Rev.*, 100, 1-9, [10.1016/j.quascirev.2014.06.025](https://doi.org/10.1016/j.quascirev.2014.06.025), 2014.
- Bradley, S. L., Milne, G. A., Shennan, I., and Edwards, R.: An improved Glacial Isostatic Adjustment model for the British Isles, *J. Quat. Sci.*, 26, 541-552, [Doi 10.1002/Jqs.1481](https://doi.org/10.1002/Jqs.1481), 2011.
- Bradley, S. L., Siddall, M., Milne, G. A., Masson-Delmotte, V., and Wolff, E.: Where might we find evidence of a Last Interglacial West Antarctic Ice Sheet collapse in Antarctic ice core records?, *Global and Planetary Change*, 88-89, 64-75, [10.1016/j.gloplacha.2012.03.004](https://doi.org/10.1016/j.gloplacha.2012.03.004), 2012.
- Bradley, S. L., Hindmarsh, R. C. A., Whitehouse, P. L., Bentley, M. J., and King, M. A.: Low post-glacial rebound rates in the Weddell Sea due to Late Holocene ice-sheet readvance, *Earth Planet. Sci. Lett.*, 413, 79-89, [10.1016/j.epsl.2014.12.039](https://doi.org/10.1016/j.epsl.2014.12.039), 2015.
- Bradley, S. L., Milne, G. A., Horton, B. P., and Zong, Y. Q.: Modelling sea level data from China and Malay-Thailand to estimate Holocene ice-volume equivalent sea level change, *Quat. Sci. Rev.*, 137, 54-68, [10.1016/j.quascirev.2016.02.002](https://doi.org/10.1016/j.quascirev.2016.02.002), 2016.
- Briggs, R., Pollard, D., and Tarasov, L.: A glacial systems model configured for large ensemble analysis of Antarctic deglaciation, *Cryosphere*, 7, 1949-1970, [DOI 10.5194/tc-7-1949-2013](https://doi.org/10.5194/tc-7-1949-2013), 2013.
- Briggs, R. D., and Tarasov, L.: How to evaluate model-derived deglaciation chronologies: a case study using Antarctica, *Quat. Sci. Rev.*, 63, 109-127, [DOI 10.1016/j.quascirev.2012.11.021](https://doi.org/10.1016/j.quascirev.2012.11.021), 2013.
- Briggs, R. D., Pollard, D., and Tarasov, L.: A data-constrained large ensemble analysis of Antarctic evolution since the Eemian, *Quat. Sci. Rev.*, 103, 91-115, [10.1016/j.quascirev.2014.09.003](https://doi.org/10.1016/j.quascirev.2014.09.003), 2014.
- Burgmann, R., and Dresen, G.: Rheology of the lower crust and upper mantle: Evidence from rock mechanics, geodesy, and field observations, *Annu. Rev. Earth Planet. Sci.*, 36, 531-567, [10.1146/annurev.earth.36.031207.124326](https://doi.org/10.1146/annurev.earth.36.031207.124326), 2008.



- 910 Burley, J. M. A., and Katz, R. F.: Variations in mid-ocean ridge CO₂ emissions driven by glacial cycles, *Earth Planet. Sci. Lett.*, 426, 246-258, [10.1016/j.epsl.2015.06.031](https://doi.org/10.1016/j.epsl.2015.06.031), 2015.
- Caron, L., Metivier, L., Greff-Lefftz, M., Fleitout, L., and Rouby, H.: Inverting Glacial Isostatic Adjustment signal using Bayesian framework and two linearly relaxing rheologies, *Geophys J Int*, 209, 1126-1147, [10.1093/gji/ggx083](https://doi.org/10.1093/gji/ggx083), 2017.
- Cathles, L. M.: *The Viscosity of Earth's Mantle*, Princeton Univ. Press, Princeton, N.J., 1975.
- 915 Celsius, A.: Anmärkning om vatnets förminskande så i Östersjön som Vesterhafvet, *Kongl. Svenska Wetenskaps Academiens Handlingar*, 4, 33-50, 1743.
- Chamberlin, T. C., and Salisbury, R. D.: *The Driftless Area of the Upper Mississippi Valley*, 6th Annual Report of the U.S. Geological Survey, 1885.
- Cheng, M. K., and Tapley, B. D.: Variations in the Earth's oblateness during the past 28 years, *J Geophys Res-Sol Ea*, 109, B09402, [10.1029/2004jb003028](https://doi.org/10.1029/2004jb003028), 2004.
- 920 Clark, C. D., Hughes, A. L. C., Greenwood, S. L., Jordan, C., and Sejrup, H. P.: Pattern and timing of retreat of the last British-Irish Ice Sheet, *Quat. Sci. Rev.*, 44, 112-146, [10.1016/j.quascirev.2010.07.019](https://doi.org/10.1016/j.quascirev.2010.07.019), 2012.
- Clark, P. U., Mitrovica, J. X., Milne, G. A., and Tamisiea, M. E.: Sea-level fingerprinting as a direct test for the source of global meltwater pulse IA, *Science*, 295, 2438-2441, 2002.
- 925 Clark, P. U., and Tarasov, L.: Closing the sea level budget at the Last Glacial Maximum, *Proc. Natl. Acad. Sci. U.S.A.*, 111, 15861-15862, [10.1073/pnas.1418970111](https://doi.org/10.1073/pnas.1418970111), 2014.
- Clement, A. J. H., Whitehouse, P. L., and Sloss, C. R.: An examination of spatial variability in the timing and magnitude of Holocene relative sea-level changes in the New Zealand archipelago, *Quat. Sci. Rev.*, 131, 73-101, [10.1016/j.quascirev.2015.09.025](https://doi.org/10.1016/j.quascirev.2015.09.025), 2016.
- 930 Conrad, C. P.: The solid Earth's influence on sea level, *Geol Soc Am Bull*, 125, 1027-1052, [10.1130/B30764.1](https://doi.org/10.1130/B30764.1), 2013.
- Craig, T. J., Calais, E., Fleitout, L., Bollinger, L., and Scotti, O.: Evidence for the release of long-term tectonic strain stored in continental interiors through intraplate earthquakes, *Geophys Res Lett*, 43, 6826-6836, [doi:10.1002/2016GL069359](https://doi.org/10.1002/2016GL069359), 2016.
- Croll, J.: *Climate and time in their geological relations: A theory of secular changes of the Earth's climate*, London, 1875.
- 935 Crowley, J. W., Katz, R. F., Huybers, P., Langmuir, C. H., and Park, S. H.: Glacial cycles drive variations in the production of oceanic crust, *Science*, 347, 1237-1240, [10.1126/science.1261508](https://doi.org/10.1126/science.1261508), 2015.
- Dal Forno, G., Gasperini, P., and Spada, G.: Implementation of the Complete Sea Level Equation in a 3D Finite Elements Scheme: A Validation Study, in *Proceedings of the VII Hotine-Marussi Symposium*, Rome, July 6–10, 2012,
- Dalca, A. V., Ferrier, K. L., Mitrovica, J. X., Perron, J. T., Milne, G. A., and Creveling, J. R.: On postglacial sea level-III. Incorporating sediment redistribution, *Geophys J Int*, 194, 45-60, [10.1093/gji/ggt089](https://doi.org/10.1093/gji/ggt089), 2013.
- 940 Daly, R. A.: A general sinking of sea-level in recent time, *Proc. Natl. Acad. Sci. U.S.A.*, 6, 246-250, DOI [10.1073/pnas.6.5.246](https://doi.org/10.1073/pnas.6.5.246), 1920.
- Daly, R. A.: Pleistocene changes of level., *American Journal of Science*, 10, 281-313, 1925.



- 945 de Boer, B., Stocchi, P., and van de Wal, R. S. W.: A fully coupled 3-D ice-sheet-sea-level model: algorithm and applications, *Geosci Model Dev*, 7, 2141-2156, 10.5194/gmd-7-2141-2014, 2014.
- de Boer, B., Stocchi, P., Whitehouse, P. L., and van de Wal, R. S. W.: Current state and future perspectives on coupled ice-sheet - sea-level modelling, *Quat. Sci. Rev.*, 169, 13-28, 10.1016/j.quascirev.2017.05.013, 2017.
- De Geer, G.: A Geochronology of the past 12,000 years, *Congres de Geologie International, Comptes Rendues*, 11, 241-253, 1912.
- 950 Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A. L., Henderson, G. M., Okuno, J., and Yokoyama, Y.: Ice-sheet collapse and sea-level rise at the Bolling warming 14,600 years ago, *Nature*, 483, 559-564, 10.1038/Nature10902, 2012.
- Dickinson, H., Freed, A. M., and Andronicos, C.: Inference of the viscosity structure and mantle conditions beneath the Central Nevada Seismic Belt from combined postseismic and lake unloading studies, *Geochem Geophys Geosy*, 17, 1740-1757, 10.1002/2015gc006207, 2016.
- 955 Düsterhus, A., Rovere, A., Carlson, A. E., Horton, B. P., Klemann, V., Tarasov, L., Barlow, N. L. M., Bradwell, T., Clark, J., Dutton, A., Gehrels, W. R., Hibbert, F. D., Hijma, M. P., Khan, N., Kopp, R. E., Sivan, D., and Tornqvist, T. E.: Palaeo-sea-level and palaeo-ice-sheet databases: problems, strategies, and perspectives, *Clim Past*, 12, 911-921, 10.5194/cp-12-911-2016, 2016a.
- 960 Düsterhus, A., Tamisiea, M. E., and Jevrejeva, S.: Estimating the sea level highstand during the last interglacial: a probabilistic massive ensemble approach, *Geophys J Int*, 206, 900-920, 2016b.
- Dutton, A., Carlson, A. E., Long, A. J., Milne, G. A., Clark, P. U., DeConto, R., Horton, B. P., Rahmstorf, S., and Raymo, M. E.: Sea-level rise due to polar ice-sheet mass loss during past warm periods, *Science*, 349, aaa4019, 10.1126/science.aaa4019, 2015.
- 965 Dyke, A. S., Moore, A., and Robertson, L.: Deglaciation of North America, Geological Survey of Canada, Ottawa, 2003.
- Dziewonski, A. M., and Anderson, D. L.: Preliminary Reference Earth Model, *Phys Earth Planet In*, 25, 297-356, 1981.
- Ekman, M.: The Changing Level of the Baltic Sea during 300 Years: A Clue to Understanding the Earth, Summer Institute for Historical Geophysics, 2009.
- Fairbanks, R. G.: A 17,000-year glacio-eustatic sea level record: influence of glacial melting rates on the Younger Dryas 970 event and deep-ocean circulation, *Nature*, 342, 637-642, 1989.
- Farrell, W. E., and Clark, J. A.: On postglacial sea level, *Geophysical Journal of the Royal Astronomical Society*, 46, 647-667, 1976.
- Ferrier, K. L., Mitrovica, J. X., Giosan, L., and Clift, P. D.: Sea-level responses to erosion and deposition of sediment in the Indus River basin and the Arabian Sea, *Earth Planet. Sci. Lett.*, 416, 12-20, 10.1016/j.epsl.2015.01.026, 2015.
- 975 Ferrier, K. L., Austermann, J., Mitrovica, J. X., and Pico, T.: Incorporating sediment compaction into a gravitationally self-consistent model for ice age sea-level change, *Geophys J Int*, 211, 663-672, 10.1093/gji/ggx293, 2017.



- Fleming, K., Johnston, P., Zwartz, D., Yokoyama, Y., Lambeck, K., and Chappell, J.: Refining the eustatic sea-level curve since the Last Glacial Maximum using far- and intermediate-field sites, *Earth Planet. Sci. Lett.*, 163, 327-342, 1998.
- 980 Freed, A. M., and Burgmann, R.: Evidence of power-law flow in the Mojave desert mantle, *Nature*, 430, 548-551, 10.1038/nature02784, 2004.
- Freed, A. M., Burgmann, R., Calais, E., and Freymueller, J.: Stress-dependent power-law flow in the upper mantle following the 2002 Denali, Alaska, earthquake, *Earth Planet. Sci. Lett.*, 252, 481-489, 10.1016/j.epsl.2006.10.011, 2006.
- Gilbert, G. K.: Proceedings of the Section of Geology and Geography, *Science*, 6, 222, 1885a.
- Gilbert, G. K.: The Inculcation of Scientific Method by Example, with an illustration drawn from the Quaternary Geology of
985 Utah, *American Journal of Science, Series 3*, vol. 31, 1885b.
- Gilbert, G. K.: Lake Bonneville, U.S. Geological Survey, Monograph 1, 1890.
- Gomez, N., Mitrovica, J. X., Huybers, P., and Clark, P. U.: Sea level as a stabilizing factor for marine-ice-sheet grounding lines, *Nat Geosci*, 3, 850-853, 10.1038/Ngeo1012, 2010.
- Gomez, N., Pollard, D., and Mitrovica, J. X.: A 3-D coupled ice sheet-sea level model applied to Antarctica through the last
990 40 ky, *Earth Planet. Sci. Lett.*, 384, 88-99, 2013.
- Gomez, N., Gregoire, L. J., Mitrovica, J. X., and Payne, A. J.: Laurentide-Cordilleran Ice Sheet saddle collapse as a contribution to meltwater pulse 1A, *Geophys Res Lett*, 42, 3954-3962, 10.1002/2015gl063960, 2015a.
- Gomez, N., Pollard, D., and Holland, D.: Sea-level feedback lowers projections of future Antarctic Ice-Sheet mass loss, *Nat Commun*, 6, 8798, 10.1038/ncomms9798, 2015b.
- 995 Gowan, E. J., Tregoning, P., Purcell, A., Montillet, J. P., and McClusky, S.: A model of the western Laurentide Ice Sheet, using observations of glacial isostatic adjustment, *Quat. Sci. Rev.*, 139, 1-16, 10.1016/j.quascirev.2016.03.003, 2016.
- Greischar, L. L., and Bentley, C. R.: Isostatic Equilibrium Grounding Line between the West Antarctic Inland Ice-Sheet and the Ross Ice Shelf, *Nature*, 283, 651-654, 1980.
- Groh, A., Ewert, H., Scheinert, M., Fritsche, M., Rulke, A., Richter, A., Rosenau, R., and Dietrich, R.: An investigation of
1000 Glacial Isostatic Adjustment over the Aundsen Sea sector, West Antarctica, *Global and Planetary Change*, 98-99, 45-53, 2012.
- Gross, R. S., and Vondrak, J.: Astrometric and space-geodetic observations of polar wander, *Geophys Res Lett*, 26, 2085-2088, Doi 10.1029/1999gl1900422, 1999.
- Gunter, B. C., Didova, O., Riva, R. E. M., Ligtenberg, S. R. M., Lenaerts, J. T. M., King, M. A., van den Broeke, M. R., and
1005 Urban, T.: Empirical estimation of present-day Antarctic glacial isostatic adjustment and ice mass change, *Cryosphere*, 8, 743-760, 10.5194/tc-8-743-2014, 2014.
- Han, D. Z., and Wahr, J.: Postglacial Rebound Analysis for a Rotating Earth, *Geoph Monog Series*, 49, 1-6, 1989.
- Han, D. Z., and Wahr, J.: The Viscoelastic Relaxation of a Realistically Stratified Earth, and a Further Analysis of Postglacial Rebound, *Geophys J Int*, 120, 287-311, DOI 10.1111/j.1365-246X.1995.tb01819.x, 1995.



- 1010 Haskell, N. A.: The motion of a viscous fluid under a surface load, *Physics-J Gen Appl P*, 6, 265-269, Doi 10.1063/1.1745329, 1935.
- Hay, C. C., Morrow, E., Kopp, R. E., and Mitrovica, J. X.: Probabilistic reanalysis of twentieth-century sea-level rise, *Nature*, 517, 481-484, 10.1038/nature14093, 2015.
- Heeszel, D. S., Wiens, D. A., Anandakrishnan, S., Aster, R. C., Dalziel, I. W. D., Huerta, A. D., Nyblade, A. A., Wilson, T. J., and Winberry, J. P.: Upper mantle structure of central and West Antarctica from array analysis of Rayleigh wave phase velocities, *Journal of Geophysical Research: Solid Earth*, 121, 10.1002/2015JB012616, 2016.
- 1015 Hetland, E. A., and Hager, B. H.: The effects of rheological layering on post-seismic deformation, *Geophys J Int*, 166, 277-292, 10.1111/j.1365-246X.2006.02974.x, 2006.
- Hibbert, F. D., Rohling, E. J., Dutton, A., Williams, F. H., Chutcharavan, P. M., Zhao, C., and Tamisiea, M. E.: Coral indicators of past sea-level change: A global repository of U-series dated benchmarks, *Quat. Sci. Rev.*, 145, 1-56, 2016.
- 1020 Hill, E. M., Davis, J. L., Tamisiea, M. E., and Lidberg, M.: Combination of geodetic observations and models for glacial isostatic adjustment fields in Fennoscandia, *Journal of Geophysical Research: Solid Earth*, 115, B07403, 10.1029/2009jb006967, 2010.
- Hirth, G., and Kohlstedt, D.: Rheology of the upper mantle and the mantle wedge: A view from the experimentalists, in: *Inside the Subduction Factory*, edited by: Eiler, J., *Geophysical Monograph Series*, AGU, 83-105, 2003.
- 1025 Hughes, A. L. C., Gyllencreutz, R., Lohne, O. S., Mangerud, J., and Svendsen, J. I.: The last Eurasian ice sheets - a chronological database and time-slice reconstruction, *DATED-1, Boreas*, 45, 1-45, 10.1111/bor.12142, 2016.
- Huybers, P., and Langmuir, C.: Feedback between deglaciation, volcanism, and atmospheric CO₂, *Earth Planet. Sci. Lett.*, 286, 479-491, 10.1016/j.epsl.2009.07.014, 2009.
- 1030 Huybers, P., and Langmuir, C. H.: Delayed CO₂ emissions from mid-ocean ridge volcanism as a possible cause of late-Pleistocene glacial cycles, *Earth Planet. Sci. Lett.*, 457, 238-249, 10.1016/j.epsl.2016.09.021, 2017.
- Ingleby, T., and Wright, T. J.: Omori-like decay of postseismic velocities following continental earthquakes, *Geophys Res Lett*, 44, 3119-3130, 10.1002/2017gl072865, 2017.
- IPCC: Climate change 2013: the physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change, 1535, 2013.
- 1035 Ivins, E. R., and Sammis, C. G.: On Lateral Viscosity Contrast in the Mantle and the Rheology of Low-Frequency Geodynamics, *Geophys J Int*, 123, 305-322, DOI 10.1111/j.1365-246X.1995.tb06856.x, 1995.
- Ivins, E. R., and James, T. S.: Bedrock response to Llanquihue Holocene and present-day glaciation in southernmost South America, *Geophys Res Lett*, 31, L24613, 2004.
- 1040 Ivins, E. R., Watkins, M. M., Yuan, D.-N., Dietrich, R., Casassa, G., and Rülke, A.: On-land ice loss and glacial isostatic adjustment at the Drake Passage: 2003–2009, *Journal of Geophysical Research*, 116, B02403, doi:10.1029/2010JB007607, 2011.



- Ivins, E. R., James, T. S., Wahr, J., Schrama, E. J. O., Landerer, F. W., and Simon, K. M.: Antarctic Contribution to Sea-Level Rise Observed by GRACE with Improved GIA Correction, *Journal of Geophysical Research: Solid Earth*, 118, 3126-3141, 10.1002/jgrb.50208, 2013.
- 1045
- James, T. S., Gowan, E. J., Wada, I., and Wang, K. L.: Viscosity of the asthenosphere from glacial isostatic adjustment and subduction dynamics at the northern Cascadia subduction zone, British Columbia, Canada, *Journal of Geophysical Research: Solid Earth*, 114, B04405, 10.1029/2008jb006077, 2009.
- Jamieson, T. F.: On the History of the Last Geological Changes in Scotland, *Quarterly Journal of the Geological Society of London*, 21, 161-203, 1865.
- 1050
- Johnson, K. M., Burgmann, R., and Freymueller, J. T.: Coupled afterslip and viscoelastic flow following the 2002 Denali Fault, Alaska earthquake, *Geophys J Int*, 176, 670-682, 10.1111/j.1365-246X.2008.04029.x, 2009.
- Johnston, P.: The Effect of Spatially Nonuniform Water Loads on Prediction of Sea-Level Change, *Geophys J Int*, 114, 615-634, 1993.
- 1055
- Kaufmann, G., Wu, P., and Ivins, E. R.: Lateral viscosity variations beneath Antarctica and their implications on regional rebound motions and seismotectonics, *J. Geodyn.*, 39, 165-181, 2005.
- Kendall, R. A., Mitrovica, J. X., and Milne, G. A.: On post-glacial sea level - II. Numerical formulation and comparative results on spherically symmetric models, *Geophys J Int*, 161, 679-706, 10.1111/j.1365-246X.2005.02553.x, 2005.
- Kendall, R. A., Latychev, K., Mitrovica, J. X., Davis, J. E., and Tamisiea, M. E.: Decontaminating tide gauge records for the influence of glacial isostatic adjustment: The potential impact of 3-D Earth structure, *Geophys Res Lett*, 33, L24318, 10.1029/2006gl028448, 2006.
- 1060
- Khan, N. S., Ashe, E., Shaw, T. A., Vacchi, M., Walker, J., Peltier, W. R., Kopp, R. E., and Horton, B. P.: Holocene Relative Sea-Level Changes from Near-, Intermediate-, and Far-Field Locations, *Current Climate Change Reports*, 1, 247-262, 2015.
- 1065
- Kierulf, H. P., Steffen, H., Simpson, M. J. R., Lidberg, M., Wu, P., and Wang, H. S.: A GPS velocity field for Fennoscandia and a consistent comparison to glacial isostatic adjustment models, *Journal of Geophysical Research: Solid Earth*, 119, 6613-6629, 10.1002/2013jb010889, 2014.
- King, M. A., Altamimi, Z., Boehm, J., Bos, M., Dach, R., Elosegui, P., Fund, F., Hernandez-Pajares, M., Lavallee, D., Cerveira, P. J. M., Penna, N., Riva, R. E. M., Steigenberger, P., van Dam, T., Vittuari, L., Williams, S., and Willis, P.: Improved Constraints on Models of Glacial Isostatic Adjustment: A Review of the Contribution of Ground-Based Geodetic Observations, *Surv Geophys*, 31, 465-507, DOI 10.1007/s10712-010-9100-4, 2010.
- 1070
- King, M. A., Bingham, R. J., Moore, P., Whitehouse, P. L., Bentley, M. J., and Milne, G. A.: Lower satellite-gravimetry estimates of Antarctic sea-level contribution, *Nature*, 491, 586-589, 10.1038/nature11621, 2012.
- King, M. A., Whitehouse, P. L., and van der Wal, W.: Incomplete separability of Antarctic plate rotation from glacial isostatic adjustment deformation within geodetic observations, *Geophys J Int*, 204, 324-330, 2016.
- 1075



- King, S. D.: Reconciling laboratory and observational models of mantle rheology in geodynamic modelling, *J. Geodyn.*, 100, 33-50, [10.1016/j.jog.2016.03.005](https://doi.org/10.1016/j.jog.2016.03.005), 2016.
- 1080 Kingslake, J., Scherer, R., Albrecht, T., Coenen, J., Powell, R., Reese, R., Stansell, N., Tulaczyk, S., Wearing, M. G., and Whitehouse, P. L.: Extensive Holocene ice sheet grounding line retreat and uplift-driven re-advance in West Antarctica, in review.
- Konrad, H., Sasgen, I., Pollard, D., and Klemann, V.: Potential of the solid-Earth response for limiting long-term West Antarctic Ice Sheet retreat in a warming climate, *Earth Planet. Sci. Lett.*, 432, 254-264, [10.1016/j.epsl.2015.10.008](https://doi.org/10.1016/j.epsl.2015.10.008), 2015.
- 1085 Kopp, R. E., Simons, F. J., Mitrovica, J. X., Maloof, A. C., and Oppenheimer, M.: Probabilistic assessment of sea level during the last interglacial stage, *Nature*, 462, 863–867, [10.1038/nature08686](https://doi.org/10.1038/nature08686), 2009.
- Kuchar, J., and Milne, G. A.: The influence of viscosity structure in the lithosphere on predictions from models of glacial isostatic adjustment, *J. Geodyn.*, 86, 1-9, [10.1016/j.jog.2015.01.002](https://doi.org/10.1016/j.jog.2015.01.002), 2015.
- Kuchar, J., Milne, G., Wolstencroft, M., Love, R., Tarasov, L., and Hijma, M.: The Influence of Sediment Isostatic Adjustment on Sea-Level Change and Land Motion along the US Gulf Coast, *Journal of Geophysical Research: Solid Earth*, 122, <https://doi.org/10.1002/2017JB014695>, 2017.
- 1090 Lambeck, K., and Nakada, M.: Late Pleistocene and Holocene sea-level change along the Australian coast, *Global and Planetary Change*, 89, 143-176, 1990.
- Lambeck, K., Smither, C., and Johnston, P.: Sea-level change, glacial rebound and mantle viscosity for northern Europe, *Geophys J Int*, 134, 102-144, 1998.
- 1095 Lambeck, K., Purcell, A., Zhao, J., and Svensson, N. O.: The Scandinavian Ice Sheet: from MIS 4 to the end of the Last Glacial Maximum, *Boreas*, 39, 410-435, DOI [10.1111/j.1502-3885.2010.00140.x](https://doi.org/10.1111/j.1502-3885.2010.00140.x), 2010.
- Lambeck, K., Purcell, A., and Dutton, A.: The anatomy of interglacial sea levels: The relationship between sea levels and ice volumes during the Last Interglacial, *Earth Planet. Sci. Lett.*, 315, 4-11, [10.1016/j.epsl.2011.08.026](https://doi.org/10.1016/j.epsl.2011.08.026), 2012.
- Lambeck, K., Rouby, H., Purcell, A., Sun, Y. Y., and Sambridge, M.: Sea level and global ice volumes from the Last Glacial Maximum to the Holocene, *Proc. Natl. Acad. Sci. U.S.A.*, 111, 15296-15303, [10.1073/pnas.1411762111](https://doi.org/10.1073/pnas.1411762111), 2014.
- 1100 Lambeck, K., Purcell, A., and Zhao, S.: The North American Late Wisconsin ice sheet and mantle viscosity from glacial rebound analyses, *Quat. Sci. Rev.*, 158, 172-210, [10.1016/j.quascirev.2016.11.033](https://doi.org/10.1016/j.quascirev.2016.11.033), 2017.
- Lange, H., Casassa, G., Ivins, E. R., Schroder, L., Fritsche, M., Richter, A., Groh, A., and Dietrich, R.: Observed crustal uplift near the Southern Patagonian Icefield constrains improved viscoelastic Earth models, *Geophys Res Lett*, 41, 805-812, [10.1002/2013gl058419](https://doi.org/10.1002/2013gl058419), 2014.
- 1105 Larsen, C. F., Motyka, R. J., Freymueller, J. T., Echelmeyer, K. A., and Ivins, E. R.: Rapid viscoelastic uplift in southeast Alaska caused by post-Little Ice Age glacial retreat, *Earth Planet. Sci. Lett.*, 237, 548-560, [10.1016/j.epsl.2005.06.032](https://doi.org/10.1016/j.epsl.2005.06.032), 2005.



- 1110 Latychev, K., Mitrovica, J. X., Tamisiea, M. E., Tromp, J., and Moucha, R.: Influence of lithospheric thickness variations on
3-D crustal velocities due to glacial isostatic adjustment, *Geophys Res Lett*, 32, L01304, 2005a.
- Latychev, K., Mitrovica, J. X., Tromp, J., Tamisiea, M. E., Komatitsch, D., and Christara, C. C.: Glacial isostatic adjustment
on 3-D Earth models: a finite-volume formulation, *Geophys J Int*, 161, 421-444, 2005b.
- Lau, H. C. P., Mitrovica, J. X., Austermann, J., Crawford, O., Al-Attar, D., and Latychev, K.: Inferences of mantle viscosity
based on ice age data sets: Radial structure, *Journal of Geophysical Research: Solid Earth*, 121, 6991-7012,
1115 10.1002/2016jb013043, 2016.
- Lecavalier, B. S., Milne, G. A., Vinther, B. M., Fisher, D. A., Dyke, A. S., and Simpson, M. J. R.: Revised estimates of
Greenland ice sheet thinning histories based on ice-core records, *Quat. Sci. Rev.*, 63, 73-82,
10.1016/j.quascirev.2012.11.030, 2013.
- Lecavalier, B. S., Milne, G. A., Simpson, M. J. R., Wake, L., Huybrechts, P., Tarasov, L., Kjeldsen, K. K., Funder, S., Long,
1120 A. J., Woodroffe, S., Dyke, A. S., and Larsen, N. K.: A model of Greenland ice sheet deglaciation constrained by
observations of relative sea level and ice extent, *Quat. Sci. Rev.*, 102, 54-84, 10.1016/j.quascirev.2014.07.018, 2014.
- Lidberg, M., Johansson, J. M., Scherneck, H. G., and Davis, J. L.: An improved and extended GPS-derived 3D velocity field
of the glacial isostatic adjustment (GIA) in Fennoscandia, *J Geodesy*, 81, 213-230, 10.1007/s00190-006-0102-4, 2007.
- Lidberg, M., Johansson, J. M., Scherneck, H. G., and Milne, G. A.: Recent results based on continuous GPS observations of
1125 the GIA process in Fennoscandia from BIFROST, *J. Geodyn.*, 50, 8-18, DOI 10.1016/j.jog.2009.11.010, 2010.
- Liu, J., Milne, G. A., Kopp, R. E., Clark, P. U., and Shennan, I.: Sea-level constraints on the amplitude and source
distribution of Meltwater Pulse 1A, *Nat Geosci*, 9, 130-134, 10.1038/Ngeo2616, 2016.
- Lund, D. C., Asimow, P. D., Farley, K. A., Rooney, T. O., Seeley, E., Jackson, E. W., and Durham, Z. M.: Enhanced East
Pacific Rise hydrothermal activity during the last two glacial terminations, *Science*, 351, 478-482,
1130 10.1126/science.aad4296, 2016.
- Lyell, C.: The Bakerian Lecture: On the Proofs of a Gradual Rising of the Land in Certain Parts of Sweden, *Philosophical
Transactions of the Royal Society of London*, 125, 1-38, 1835.
- Margold, M., Stokes, C. R., and Clark, C. D.: Ice streams in the Laurentide Ice Sheet: Identification, characteristics and
comparison to modern ice sheets, *Earth-Sci Rev*, 143, 117-146, 10.1016/j.earscirev.2015.01.011, 2015.
- 1135 Martin-Espanol, A., King, M. A., Mangion, A. Z., Andrews, S., Moore, P., and Bamber, J.: An assessment of forward and
inverse GIA solutions for Antarctica, *Journal of Geophysical Research*, 121, 6947-6965, 2016a.
- Martin-Espanol, A., Zammit-Mangion, A., Clarke, P. J., Flament, T., Helm, V., King, M. A., Luthcke, S. B., Petrie, E.,
Remy, F., Schon, N., Wouters, B., and Bamber, J. L.: Spatial and temporal Antarctic Ice Sheet mass trends, glacio-
isostatic adjustment, and surface processes from a joint inversion of satellite altimeter, gravity, and GPS data, *Journal
of Geophysical Research: Earth Surface*, 121, 182-200, 10.1002/2015jf003550, 2016b.
- 1140 Martinec, Z.: Spectral-finite element approach to three-dimensional viscoelastic relaxation in a spherical earth, *Geophys J
Int*, 142, 117-141, DOI 10.1046/j.1365-246x.2000.00138.x, 2000.



- Martinec, Z., and Hagedoorn, J.: The rotational feedback on linear-momentum balance in glacial isostatic adjustment, *Geophys J Int*, 199, 1823-1846, 10.1093/gji/ggu369, 2014.
- 1145 Martinec, Z., Sasgen, I., and Velinsky, J.: The forward sensitivity and adjoint-state methods of glacial isostatic adjustment, *Geophys J Int*, 200, 77-105, 10.1093/gji/ggu378, 2015.
- Matsuoka, K., Hindmarsh, R. C. A., Moholdt, G., Bentley, M. J., Pritchard, H. D., Brown, J., Conway, H., Drews, R., Durand, G., Goldberg, D., Hattermann, T., Kingslake, J., Lenaerts, J. T. M., Martín, C., Mulvaney, R., Nicholls, K. W., Pattyn, F., Ross, N., Scambos, T., and Whitehouse, P. L.: Antarctic ice rises and rumpled: Their properties and
1150 significance for ice-sheet dynamics and evolution, *Earth-Sci Rev*, 150, 724-745, 2015.
- Mazzotti, S., Lambert, A., Henton, J., James, T. S., and Courtier, N.: Absolute gravity calibration of GPS velocities and glacial isostatic adjustment in mid-continent North America, *Geophys Res Lett*, 38, L24311, 10.1029/2011gl049846, 2011.
- 1155 McConnell, R. K.: Viscosity of Mantle from Relaxation Time Spectra of Isostatic Adjustment, *Journal of Geophysical Research*, 73, 7089–7105, 10.1029/JB073i022p07089, 1968.
- Memin, A., Rogister, Y., Hinderer, J., Omang, O. C., and Luck, B.: Secular gravity variation at Svalbard (Norway) from ground observations and GRACE satellite data, *Geophys J Int*, 184, 1119-1130, 10.1111/j.1365-246X.2010.04922.x, 2011.
- 1160 Metivier, L., Caron, L., Greff-Lefftz, M., Pajot-Metivier, G., Fleitout, L., and Rouby, H.: Evidence for postglacial signatures in gravity gradients: A clue in lower mantle viscosity, *Earth Planet. Sci. Lett.*, 452, 146-156, 10.1016/j.epsl.2016.07.034, 2016.
- Mey, J., Scherler, D., Wickert, A. D., Egholm, D. L., Tesauero, M., Schildgen, T. F., and Strecker, M. R.: Glacial isostatic uplift of the European Alps, *Nat Commun*, 7, 13382, 10.1038/ncomms13382, 2016.
- 1165 Miller, G. H., Geirsdottir, A., Zhong, Y. F., Larsen, D. J., Otto-Bliesner, B. L., Holland, M. M., Bailey, D. A., Refsnider, K. A., Lehman, S. J., Southon, J. R., Anderson, C., Bjornsson, H., and Thordarson, T.: Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean feedbacks, *Geophys Res Lett*, 39, L02708, 10.1029/2011gl050168, 2012.
- Milne, G. A.: Refining models of the glacial isostatic adjustment process, PhD Thesis, University of Toronto, Toronto, 1998.
- Milne, G. A., and Mitrovica, J. X.: Postglacial sea-level change on a rotating Earth, *Geophys J Int*, 133, 1-19, 1998.
- 1170 Milne, G. A., Davis, J. L., Mitrovica, J. X., Scherneck, H. G., Johansson, J. M., Vermeer, M., and Koivula, H.: Space-geodetic constraints on glacial isostatic adjustment in Fennoscandia, *Science*, 291, 2381-2385, 2001.
- Milne, G. A., Mitrovica, J. X., and Schrag, D. P.: Estimating past continental ice volume from sea-level data, *Quat. Sci. Rev.*, 21, 361-376, 2002.
- 1175 Milne, G. A., Mitrovica, J. X., Scherneck, H. G., Davis, J. L., Johansson, J. M., Koivula, H., and Vermeer, M.: Continuous GPS measurements of postglacial adjustment in Fennoscandia: 2. Modeling results, *Journal of Geophysical Research: Solid Earth*, 109, B02412, 10.1029/2003jb002619, 2004.



- Milne, G. A., and Mitrovica, J. X.: Searching for eustasy in deglacial sea-level histories, *Quat. Sci. Rev.*, 27, 2292-2302, 10.1016/j.quascirev.2008.08.018, 2008.
- Milne, G. A., Gehrels, W. R., Hughes, C. W., and Tamisiea, M. E.: Identifying the causes of sea-level change, *Nat Geosci*, 2, 471-478, Doi 10.1038/Ngeo544, 2009.
- 1180 Mitrovica, J. X., and Peltier, W. R.: A Complete Formalism for the Inversion of Postglacial Rebound Data - Resolving Power Analysis, *Geophys J Int*, 104, 267-288, DOI 10.1111/j.1365-246X.1991.tb02511.x, 1991a.
- Mitrovica, J. X., and Peltier, W. R.: On postglacial geoid subsidence over the equatorial oceans, *J Geophys Res-Sol Ea*, 96, 20053-20071, 1991b.
- 1185 Mitrovica, J. X.: Haskell [1935] revisited, *Journal of Geophysical Research: Solid Earth*, 101, 555-569, Doi 10.1029/95jb03208, 1996.
- Mitrovica, J. X., Tamisiea, M. E., Davis, J. L., and Milne, G. A.: Recent mass balance of polar ice sheets inferred from patterns of global sea-level change, *Nature*, 409, 1026-1029, 2001.
- Mitrovica, J. X., and Milne, G. A.: On the origin of late Holocene sea-level highstands within equatorial ocean basins, *Quat.*
- 1190 *Sci. Rev.*, 21, 2179-2190, 2002.
- Mitrovica, J. X., and Milne, G. A.: On post-glacial sea level: I. General theory, *Geophys J Int*, 154, 253-267, 2003.
- Mitrovica, J. X., and Forte, A. M.: A new inference of mantle viscosity based upon joint inversion of convection and glacial isostatic adjustment data, *Earth Planet. Sci. Lett.*, 225, 177-189, 10.1016/j.epsl.2004.06.005, 2004.
- Mitrovica, J. X., Wahr, J., Matsuyama, I., and Paulson, A.: The rotational stability of an ice-age earth, *Geophys J Int*, 161, 491-506, DOI 10.1111/j.1365-246X.2005.02609.x, 2005.
- 1195 Mitrovica, J. X., Gomez, N., and Clark, P. U.: The Sea-Level Fingerprint of West Antarctic Collapse, *Science*, 323, 753-753, 2009.
- Mitrovica, J. X., and Wahr, J.: Ice Age Earth Rotation, *Annu. Rev. Earth Planet. Sci.*, 39, 577-616, 10.1146/annurev-earth-040610-133404, 2011.
- 1200 Mitrovica, J. X., Hay, C. C., Morrow, E., Kopp, R. E., Dumberry, M., and Stanley, S.: Reconciling past changes in Earth's rotation with 20th century global sea-level rise: Resolving Munk's enigma, *Science Advances*, 1, e1500679, doi: 10.1126/sciadv.1500679, 2015.
- Moucha, R., and Ruetenik, G. A.: Interplay between dynamic topography and flexure along the US Atlantic passive margin: Insights from landscape evolution, *Global and Planetary Change*, 149, 72-78, 10.1016/j.gloplacha.2017.01.004, 2017.
- 1205 Nakada, M., Okuno, J., and Yokoyama, Y.: Total meltwater volume since the Last Glacial Maximum and viscosity structure of Earth's mantle inferred from relative sea level changes at Barbados and Bonaparte Gulf and GIA-induced J_2 , *Geophys J Int*, 204, 1237-1253, 10.1093/gji/ggv520, 2016.
- Nakiboglu, S. M., and Lambeck, K.: Deglaciation Effects on the Rotation of the Earth, *Geophysical Journal of the Royal Astronomical Society*, 62, 49-58, DOI 10.1111/j.1365-246X.1980.tb04843.x, 1980.



- 1210 Nansen, F.: The strandflat and isostasy, *Videnskapsselskapets Skrifter. 1. Matematisk-Naturhistorisk Klasse*, 313 pp. pp., 1921.
- Nield, G. A., Barletta, V. R., Bordoni, A., King, M. A., Whitehouse, P. L., Clarke, P. J., Domack, E., Scambos, T. A., and Berthier, E.: Rapid bedrock uplift in the Antarctic Peninsula explained by viscoelastic response to recent ice unloading, *Earth Planet. Sci. Lett.*, 397, 32-41, 2014.
- 1215 O'Connell, R. J.: Pleistocene Glaciation and Viscosity of Lower Mantle, *Geophysical Journal of the Royal Astronomical Society*, 23, 299-327, DOI 10.1111/j.1365-246X.1971.tb01823.x, 1971.
- Pagli, C., Sigmundsson, F., Lund, B., Sturkell, E., Geirsson, H., Einarsson, P., Arnadóttir, T., and Hreinsdóttir, S.: Glacio-isostatic deformation around the Vatnajökull ice cap, Iceland, induced by recent climate warming: GPS observations and finite element modeling, *Journal of Geophysical Research: Solid Earth*, 112, B08405, 10.1029/2006jb004421, 2007.
- 1220 Parrenin, F., Dreyfus, G., Durand, G., Fujita, S., Gagliardini, O., Gillet, F., Jouzel, J., Kawamura, K., Lhomme, N., Masson-Delmotte, V., Ritz, C., Schwander, J., Shoji, H., Uemura, R., Watanabe, O., and Yoshida, N.: 1-D-ice flow modelling at EPICA Dome C and Dome Fuji, East Antarctica, *Clim Past*, 3, 243-259, 2007.
- Patton, H., Hubbard, A., Andreassen, K., Auriac, A., Whitehouse, P. L., Stroeven, A. P., Shackleton, C., Winsborrow, M., 1225 Heyman, J., and Hall, A. M.: Deglaciation of the Eurasian ice sheet complex, *Quat. Sci. Rev.*, 169, 148-172, 2017.
- Paulson, A., Zhong, S. J., and Wahr, J.: Modelling post-glacial rebound with lateral viscosity variations, *Geophys J Int*, 163, 357-371, DOI 10.1111/j.1365-246X.2005.02645.x, 2005.
- Paulson, A., Zhong, S. J., and Wahr, J.: Inference of mantle viscosity from GRACE and relative sea level data, *Geophys J Int*, 171, 497-508, 10.1111/j.1365-246X.2007.03556.x, 2007a.
- 1230 Paulson, A., Zhong, S. J., and Wahr, J.: Limitations on the inversion for mantle viscosity from postglacial rebound, *Geophys J Int*, 168, 1195-1209, 10.1111/j.1365-246X.2006.03222.x, 2007b.
- Peltier, W. R.: Impulse Response of a Maxwell Earth, *Rev Geophys*, 12, 649-669, 10.1029/RG012i004p00649, 1974.
- Peltier, W. R., Shennan, I., Drummond, R., and Horton, B.: On the postglacial isostatic adjustment of the British Isles and the shallow viscoelastic structure of the Earth, *Geophys J Int*, 148, 443-475, DOI 10.1046/j.1365-246x.2002.01586.x, 1235 2002.
- Peltier, W. R.: Global glacial isostasy and the surface of the ice-age earth: The ICE-5G (VM2) model and GRACE, *Annu. Rev. Earth Planet. Sci.*, 32, 111-149, 2004.
- Peltier, W. R., Argus, D. F., and Drummond, R.: Space geodesy constrains ice age terminal deglaciation: The global ICE-6G_C (VM5a) model, *Journal of Geophysical Research: Solid Earth*, 120, 450-487, 10.1002/2014jb011176, 2015.
- 1240 Pico, T., Mitrovica, J. X., Ferrier, K. L., and Braun, J.: Global ice volume during MIS 3 inferred from a sea-level analysis of sedimentary core records in the Yellow River Delta, *Quat. Sci. Rev.*, 152, 72-79, 10.1016/j.quascirev.2016.09.012, 2016.
- Playfair, J.: Illustrations of the Huttonian theory of the Earth, Edinburgh, 1802.



- 1245 Pollard, D., DeConto, R. M., and Alley, R. B.: Potential Antarctic Ice Sheet retreat driven by hydrofracturing and ice cliff failure, *Earth Planet. Sci. Lett.*, 412, 112-121, [10.1016/j.epsl.2014.12.035](https://doi.org/10.1016/j.epsl.2014.12.035), 2015.
- Pollard, D., Chang, W., Haran, M., Applegate, P., and DeConto, R.: Large ensemble modeling of the last deglacial retreat of the West Antarctic Ice Sheet: comparison of simple and advanced statistical techniques, *Geosci Model Dev*, 9, 1697-1723, [10.5194/gmd-9-1697-2016](https://doi.org/10.5194/gmd-9-1697-2016), 2016.
- 1250 Pollard, D., Gomez, N., and DeConto, R. M.: Variations of the Antarctic Ice Sheet in a coupled ice sheet-Earth-sea level model: sensitivity to viscoelastic Earth properties, *Journal of Geophysical Research: Earth Surface*, 122, 2124-2138, [10.1002/2017JF004371](https://doi.org/10.1002/2017JF004371), 2017.
- Pollitz, F. F.: Transient rheology of the upper mantle beneath central Alaska inferred from the crustal velocity field following the 2002 Denali earthquake, *Journal of Geophysical Research: Solid Earth*, 110, B08407, [10.1029/2005jb003672](https://doi.org/10.1029/2005jb003672), 2005.
- 1255 Praetorius, S., Mix, A., Jensen, B., Froese, D., Milne, G., Wolhowe, M., Addison, J., and Prah, F.: Interaction between climate, volcanism, and isostatic rebound in Southeast Alaska during the last deglaciation, *Earth Planet. Sci. Lett.*, 452, 79-89, [10.1016/j.epsl.2016.07.033](https://doi.org/10.1016/j.epsl.2016.07.033), 2016.
- Purcell, A., Dehecq, A., Tregoning, P., Potter, E. K., McClusky, S. C., and Lambeck, K.: Relationship between glacial isostatic adjustment and gravity perturbations observed by GRACE, *Geophys Res Lett*, 38, L18305, [10.1029/2011gl048624](https://doi.org/10.1029/2011gl048624), 2011.
- 1260 Raymo, M. E., Mitrovica, J. X., O'Leary, M. J., DeConto, R. M., and Hearty, P. L.: Departures from eustasy in Pliocene sea-level records, *Nat Geosci*, 4, 328-332, [Doi 10.1038/Ngeo1118](https://doi.org/10.1038/Ngeo1118), 2011.
- Rietbroek, R., Brunnabend, S. E., Kusche, J., Schroter, J., and Dahle, C.: Revisiting the contemporary sea-level budget on global and regional scales, *Proc. Natl. Acad. Sci. U.S.A.*, 113, 1504-1509, [10.1073/pnas.1519132113](https://doi.org/10.1073/pnas.1519132113), 2016.
- 1265 Ritsema, J., Deuss, A., van Heijst, H. J., and Woodhouse, J. H.: S40RTS: a degree-40 shear-velocity model for the mantle from new Rayleigh wave dispersion, teleseismic traveltimes and normal-mode splitting function measurements, *Geophys J Int*, 184, 1223-1236, [10.1111/j.1365-246X.2010.04884.x](https://doi.org/10.1111/j.1365-246X.2010.04884.x), 2011.
- Riva, R. E. M., Gunter, B. C., Urban, T. J., Vermeersen, B. L. A., Lindenbergh, R. C., Helsen, M. M., Bamber, J. L., de Wal, R., van den Broeke, M. R., and Schutz, B. E.: Glacial Isostatic Adjustment over Antarctica from combined ICESat and GRACE satellite data, *Earth Planet. Sci. Lett.*, 288, 516-523, [10.1016/j.epsl.2009.10.013](https://doi.org/10.1016/j.epsl.2009.10.013), 2009.
- 1270 Riva, R. E. M., Bamber, J. L., Lavalée, D. A., and Wouters, B.: Sea-level fingerprint of continental water and ice mass change from GRACE, *Geophys Res Lett*, 37, L19605, [10.1029/2010gl044770](https://doi.org/10.1029/2010gl044770), 2010.
- Root, B. C., Tarasov, L., and van der Wal, W.: GRACE gravity observations constrain Weichselian ice thickness in the Barents Sea, *Geophys Res Lett*, 42, 3313-3320, [10.1002/2015gl063769](https://doi.org/10.1002/2015gl063769), 2015.
- 1275 Rovere, A., Raymo, M. E., Mitrovica, J. X., Hearty, P. J., O'Leary, M. J., and Inglis, J. D.: The Mid-Pliocene sea-level conundrum: Glacial isostasy, eustasy and dynamic topography, *Earth Planet. Sci. Lett.*, 387, 27-33, [DOI 10.1016/j.epsl.2013.10.030](https://doi.org/10.1016/j.epsl.2013.10.030), 2014.



- 1280 Rovere, A., Raymo, M. E., Vacchi, M., Lorscheid, T., Stocchi, P., Gómez-Pujol, L., Harris, D. L., Casella, E., O'Leary, M. J., and Hearty, P. J.: The analysis of Last Interglacial (MIS 5e) relative sea-level indicators: Reconstructing sea-level in a warmer world, *Earth-Sci Rev*, 159, 404-427, 2016.
- Rowley, D. B., Forte, A. M., Moucha, R., Mitrovica, J. X., Simmons, N. A., and Grand, S. P.: Dynamic Topography Change of the Eastern United States Since 3 Million Years Ago, *Science*, 340, 1560-1563, 10.1126/science.1229180, 2013.
- Rudzki, M. P.: Deformationen der Erde unter der Last des Inlandeises, *Bulletin international de l'Academie des sciences de Cracovie*, 169-215, 1899.
- 1285 Rugenstein, M., Stocchi, P., von der Heydt, A., Dijkstra, H., and Brinkhuis, H.: Emplacement of Antarctic ice sheet mass affects circumpolar ocean flow, *Global and Planetary Change*, 118, 16-24, 10.1016/j.gloplacha.2014.03.011, 2014.
- Sabadini, R., Yuen, D. A., and Boschi, E.: Polar Wandering and the Forced Responses of a Rotating, Multilayered, Viscoelastic Planet, *Journal of Geophysical Research*, 87, 2885-2903, DOI 10.1029/JB087iB04p02885, 1982.
- Sasgen, I., Martinec, Z., and Fleming, K.: Regional ice-mass changes and glacial-isostatic adjustment in Antarctica from 1290 GRACE, *Earth Planet. Sci. Lett.*, 264, 391-401, DOI 10.1016/j.epsl.2007.09.029, 2007.
- Sasgen, I., Konrad, H., Ivins, E. R., Van den Broeke, M. R., Bamber, J. L., Martinec, Z., and Klemann, V.: Antarctic ice-mass balance 2003 to 2012: regional reanalysis of GRACE satellite gravimetry measurements with improved estimate of glacial-isostatic adjustment based on GPS uplift rates, *Cryosphere*, 7, 1499-1512, 10.5194/tc-7-1499-2013, 2013.
- Sasgen, I., Martin-Espanol, A., Horvath, A., Klemann, V., Petrie, E. J., Wouters, B., Horvath, M., Pail, R., Bamber, J. L., 1295 Clarke, P. J., Konrad, H., and Drinkwater, M. R.: Joint inversion estimate of regional glacial isostatic adjustment in Antarctica considering a lateral varying Earth structure (ESA STSE Project REGINA), *Geophys J Int*, 211, 1534-1553, 10.1093/gji/ggx368, 2017.
- Sato, T., Larsen, C. F., Miura, S., Ohta, Y., Fujimoto, H., Sun, W. K., Motyka, R. J., and Freymueller, J. T.: Reevaluation of the viscoelastic and elastic responses to the past and present-day ice changes in Southeast Alaska, *Tectonophysics*, 511, 1300 79-88, DOI 10.1016/j.tecto.2010.05.009, 2011.
- Sato, T., Miura, S., Sun, W. K., Sugano, T., Freymueller, J. T., Larsen, C. F., Ohta, Y., Fujimoto, H., Inazu, D., and Motyka, R. J.: Gravity and uplift rates observed in southeast Alaska and their comparison with GIA model predictions, *Journal of Geophysical Research: Solid Earth*, 117, 10.1029/2011jb008485, 2012.
- Schoof, C.: Ice sheet grounding line dynamics: Steady states, stability, and hysteresis, *Journal of Geophysical Research: Earth Surface*, 112, 10.1029/2006jf000664, 2007.
- 1305 Sella, G. F., Stein, S., Dixon, T. H., Craymer, M., James, T. S., Mazzotti, S., and Dokka, R. K.: Observation of glacial isostatic adjustment in "stable" North America with GPS, *Geophys Res Lett*, 34, L02306, 2007.
- Shennan, I., Peltier, W. R., Drummond, R., and Horton, B.: Global to local scale parameters determining relative sea-level changes and the post-glacial isostatic adjustment of Great Britain, *Quat. Sci. Rev.*, 21, 397-408, Doi 10.1016/S0277- 1310 3791(01)00091-9, 2002.



- 1315 Shepherd, A., Ivins, E. R., Geruo, A., Barletta, V. R., Bentley, M. J., Bettadpur, S., Briggs, K. H., Bromwich, D. H., Forsberg, R., Galin, N., Horwath, M., Jacobs, S., Joughin, I., King, M. A., Lenaerts, J. T. M., Li, J. L., Ligtenberg, S. R. M., Luckman, A., Luthcke, S. B., McMillan, M., Meister, R., Milne, G., Mouginot, J., Muir, A., Nicolas, J. P., Paden, J., Payne, A. J., Pritchard, H., Rignot, E., Rott, H., Sorensen, L. S., Scambos, T. A., Scheuchl, B., Schrama, E. J. O., Smith, B., Sundal, A. V., van Angelen, J. H., van de Berg, W. J., van den Broeke, M. R., Vaughan, D. G., Velicogna, I., Wahr, J., Whitehouse, P. L., Wingham, D. J., Yi, D. H., Young, D., and Zwally, H. J.: A Reconciled Estimate of Ice-Sheet Mass Balance, *Science*, 338, 1183-1189, [10.1126/science.1228102](https://doi.org/10.1126/science.1228102), 2012.
- 1320 Sigmundsson, F., Pinel, V., Lund, B., Albino, F., Pagli, C., Geirsson, H., and Sturkell, E.: Climate effects on volcanism: influence on magmatic systems of loading and unloading from ice mass variations, with examples from Iceland, *Philos T R Soc A*, 368, 2519-2534, [10.1098/rsta.2010.0042](https://doi.org/10.1098/rsta.2010.0042), 2010.
- Simms, A. R., Ivins, E. R., DeWitt, R., Kouremenos, P., and Simkins, L. M.: Timing of the most recent Neoglacial advance and retreat in the South Shetland Islands, Antarctica Peninsula: insights from raised beaches and Holocene uplift rates, *Quat. Sci. Rev.*, 37, 41-55, 2012.
- 1325 Simon, K. M., James, T. S., Henton, J. A., and Dyke, A. S.: A glacial isostatic adjustment model for the central and northern Laurentide Ice Sheet based on relative sea level and GPS measurements, *Geophys J Int*, 205, 1618-1636, [10.1093/gji/ggw103](https://doi.org/10.1093/gji/ggw103), 2016.
- Simpson, M. J. R., Milne, G. A., Huybrechts, P., and Long, A. J.: Calibrating a glaciological model of the Greenland ice sheet from the Last Glacial Maximum to present-day using field observations of relative sea level and ice extent, *Quat. Sci. Rev.*, 28, 1631-1657, DOI [10.1016/j.quascirev.2009.03.004](https://doi.org/10.1016/j.quascirev.2009.03.004), 2009.
- 1330 Slangen, A. B. A., Carson, M., Katsman, C. A., van de Wal, R. S. W., Kohl, A., Vermeersen, L. L. A., and Stammer, D.: Projecting twenty-first century regional sea-level changes, *Climatic Change*, 124, 317-332, [10.1007/s10584-014-1080-9](https://doi.org/10.1007/s10584-014-1080-9), 2014.
- Spada, G., and Stocchi, P.: *The sea level equation*, Aracne, 2006.
- 1335 Spada, G., Barletta, V. R., Klemann, V., Riva, R. E. M., Martinec, Z., Gasperini, P., Lund, B., Wolf, D., Vermeersen, L. L. A., and King, M. A.: A benchmark study for glacial isostatic adjustment codes, *Geophys J Int*, 185, 106-132, [10.1111/j.1365-246X.2011.04952.x](https://doi.org/10.1111/j.1365-246X.2011.04952.x), 2011.
- Steffen, H., Kaufmann, G., and Wu, P.: Three-dimensional finite-element modeling of the glacial isostatic adjustment in Fennoscandia, *Earth Planet. Sci. Lett.*, 250, 358-375, [10.1016/j.epsl.2006.08.003](https://doi.org/10.1016/j.epsl.2006.08.003), 2006.
- 1340 Steffen, H., Gitlein, O., Denker, H., Muller, J., and Timmen, L.: Present rate of uplift in Fennoscandia from GRACE and absolute gravimetry, *Tectonophysics*, 474, 69-77, [10.1016/j.tecto.2009.01.012](https://doi.org/10.1016/j.tecto.2009.01.012), 2009.
- Steffen, H., and Wu, P.: Glacial isostatic adjustment in Fennoscandia-A review of data and modeling, *J. Geodyn.*, 52, 169-204, [10.1016/j.jog.2011.03.002](https://doi.org/10.1016/j.jog.2011.03.002), 2011.
- Steffen, H., and Wu, P.: The sensitivity of GNSS measurements in Fennoscandia to distinct three-dimensional upper-mantle structures, *Solid Earth*, 5, 557-567, [10.5194/se-5-557-2014](https://doi.org/10.5194/se-5-557-2014), 2014.



- 1345 Steffen, R., Eaton, D. W., and Wu, P.: Moment tensors, state of stress and their relation to post-glacial rebound in northeastern Canada, *Geophys J Int*, 189, 1741-1752, 10.1111/j.1365-246X.2012.05452.x, 2012.
- Steffen, R., Wu, P., Steffen, H., and Eaton, D. W.: The effect of earth rheology and ice-sheet size on fault slip and magnitude of postglacial earthquakes, *Earth Planet. Sci. Lett.*, 388, 71-80, 10.1016/j.epsl.2013.11.058, 2014.
- Sternai, P., Caricchi, L., Castellort, S., and Champagnac, J. D.: Deglaciation and glacial erosion: A joint control on magma
1350 productivity by continental unloading, *Geophys Res Lett*, 43, 1632-1641, 10.1002/2015gl067285, 2016.
- Stewart, I. S., Sauber, J., and Rose, J.: Glacio-seismotectonics: ice sheets, crustal deformation and seismicity, *Quat. Sci. Rev.*, 19, 1367-1389, Doi 10.1016/S0277-3791(00)00094-9, 2000.
- Stokes, G. G.: On the variation of gravity at the surface of the Earth, *Transactions of the Cambridge Philosophical Society*, 8, 672-695, 1849.
- 1355 Tamisiea, M. E., Mitrovica, J. X., and Davis, J. L.: GRACE gravity data constrain ancient ice geometries and continental dynamics over Laurentia, *Science*, 316, 881-883, 10.1126/science.1137157, 2007.
- Tamisiea, M. E.: Ongoing glacial isostatic contributions to observations of sea level change, *Geophys J Int*, 186, 1036-1044, DOI 10.1111/j.1365-246X.2011.05116.x, 2011.
- Tanaka, Y., Klemann, V., Martinec, Z., and Riva, R. E. M.: Spectral-finite element approach to viscoelastic relaxation in a
1360 spherical compressible Earth: application to GIA modelling, *Geophys J Int*, 184, 220-234, 10.1111/j.1365-246X.2010.04854.x, 2011.
- Tarasov, L., and Peltier, W. R.: Greenland glacial history and local geodynamic consequences, *Geophys J Int*, 150, 198-229, DOI 10.1046/j.1365-246X.2002.01702.x, 2002.
- Tarasov, L., Dyke, A. S., Neal, R. M., and Peltier, W. R.: A data-calibrated distribution of deglacial chronologies for the
1365 North American ice complex from glaciological modeling, *Earth Planet. Sci. Lett.*, 315, 30-40, 10.1016/j.epsl.2011.09.010, 2012.
- Thomas, I. D., King, M. A., Bentley, M. J., Whitehouse, P. L., Penna, N. T., Williams, S. D. P., Riva, R. E. M., Lavallee, D. A., Clarke, P. J., King, E. C., Hindmarsh, R. C. A., and Koivula, H.: Widespread low rates of Antarctic glacial isostatic adjustment revealed by GPS observations, *Geophys Res Lett*, 38, L22302, 10.1029/2011GL049277, 2011.
- 1370 van Dam, T., Francis, O., Wahr, J., Khan, S. A., Bevis, M., and van den Broeke, M. R.: Using GPS and absolute gravity observations to separate the effects of present-day and Pleistocene ice-mass changes in South East Greenland, *Earth Planet. Sci. Lett.*, 459, 127-135, 10.1016/j.epsl.2016.11.014, 2017.
- van der Wal, W., Barnhoorn, A., Stocchi, P., Gradmann, S., Wu, P., Drury, M., and Vermeersen, B.: Glacial isostatic adjustment model with composite 3-D Earth rheology for Fennoscandia, *Geophys J Int*, 194, 61-77, 10.1093/gji/ggt099, 2013.
- 1375 van der Wal, W., Whitehouse, P. L., and Schrama, E. J. O.: Effect of GIA models with 3D composite mantle viscosity on GRACE mass balance estimates for Antarctica, *Earth Planet. Sci. Lett.*, 414, 134-143, 2015.



- van der Wal, W., and IJpelaar, T.: The effect of sediment loading in Fennoscandia and the Barents Sea during the last glacial cycle on glacial isostatic adjustment observations, *Solid Earth*, 8, 955-968, <https://doi.org/10.5194/se-8-955-2017>, 2017.
- 1380 Wahr, J., Dazhong, H., and Trupin, A.: Predictions of Vertical Uplift Caused by Changing Polar Ice Volumes on a Viscoelastic Earth, *Geophys Res Lett*, 22, 977-980, Doi 10.1029/94gl02840, 1995.
- Wahr, J., Wingham, D., and Bentley, C.: A method of combining ICESat and GRACE satellite data to constrain Antarctic mass balance, *Journal of Geophysical Research*, 105, 16279-16294, 2000.
- 1385 Walcott, R. I.: Past sea levels, eustasy and deformation of the earth, *Quaternary Res*, 2, 1-14, 1972.
- Watcham, E. P., Bentley, M. J., Hodgson, D. A., Roberts, S. J., Fretwell, P. T., Lloyd, J. M., Larter, R. D., Whitehouse, P. L., Leng, M. J., Monien, P., and Moreton, S. G.: A new Holocene relative sea level curve for the South Shetland Islands, Antarctica, *Quat. Sci. Rev.*, 30, 3152-3170, 2011.
- Watts, A. B., Zhong, S. J., and Hunter, J.: The Behavior of the Lithosphere on Seismic to Geologic Timescales, *Annu. Rev. Earth Planet. Sci.*, 41, 443-468, [10.1146/annurev-earth-042711-105457](https://doi.org/10.1146/annurev-earth-042711-105457), 2013.
- 1390 Whitehouse, P., Latychev, K., Milne, G. A., Mitrovica, J. X., and Kendall, R.: Impact of 3-D Earth structure on Fennoscandian glacial isostatic adjustment: Implications for space-geodetic estimates of present-day crustal deformations, *Geophys Res Lett*, 33, L13502, 2006.
- Whitehouse, P. L., Allen, M. B., and Milne, G. A.: Glacial isostatic adjustment as a control on coastal processes: An example from the Siberian Arctic, *Geology*, 35, 747-750, 2007.
- 1395 Whitehouse, P. L., Bentley, M. J., and Le Brocq, A. M.: A deglacial model for Antarctica: geological constraints and glaciological modelling as a basis for a new model of Antarctic glacial isostatic adjustment, *Quat. Sci. Rev.*, 32, 1-24, 2012a.
- Whitehouse, P. L., Bentley, M. J., Milne, G. A., King, M. A., and Thomas, I. D.: A new glacial isostatic adjustment model for Antarctica: calibrating the deglacial model using observations of relative sea-level and present-day uplift rates, *Geophys J Int*, 190, 1464-1482, [10.1111/j.1365-246X.2012.05557.x](https://doi.org/10.1111/j.1365-246X.2012.05557.x), 2012b.
- 1400 Wickert, A. D., Mitrovica, J. X., Williams, C., and Anderson, R. S.: Gradual demise of a thin southern Laurentide ice sheet recorded by Mississippi drainage, *Nature*, 502, 668-671, [10.1038/nature12609](https://doi.org/10.1038/nature12609), 2013.
- Wilmes, S. B., Green, J. A. M., Gomez, N., Rippeth, T. P., and Lau, H.: Global Tidal Impacts of Large-Scale Ice Sheet Collapses, *J Geophys Res-Oceans*, 122, 8354-8370, [10.1002/2017jc013109](https://doi.org/10.1002/2017jc013109), 2017.
- 1405 Wolstencroft, M., Shen, Z. X., Toernqvist, T. E., Milne, G. A., and Kulp, M.: Understanding subsidence in the Mississippi Delta region due to sediment, ice, and ocean loading: Insights from geophysical modeling, *Journal of Geophysical Research: Solid Earth*, 119, 3838-3856, [10.1002/2013jb010928](https://doi.org/10.1002/2013jb010928), 2014.
- Woodward, R. S.: On the form and position of mean sea level, *United States Geological Survey Bulletin*, 48, 87-170, 1888.
- 1410 Wouters, B., Bonin, J. A., Chambers, D. P., Riva, R. E. M., Sasgen, I., and Wahr, J.: GRACE, time-varying gravity, Earth system dynamics and climate change, *Rep Prog Phys*, 77, 116801, [10.1088/0034-4885/77/11/116801](https://doi.org/10.1088/0034-4885/77/11/116801), 2014.



1415 Wu, P.: The Response of a Maxwell Earth to Applied Surface Mass Loads: Glacial Isostatic Adjustment, University of Toronto, 1978.

Wu, P., and Peltier, W. R.: Pleistocene Deglaciation and the Earth's Rotation - a New Analysis, *Geophysical Journal of the Royal Astronomical Society*, 76, 753-791, DOI 10.1111/j.1365-246X.1984.tb01920.x, 1984.

Wu, P.: Postglacial rebound modeling with power law rheology, in: *Dynamics of the Ice Age Earth: a Modern Perspective*, Trans Tech Publishers, Uetikon-Zürich, Switzerland, 365-382, 1998.

1420 Wu, P., and van der Wal, W.: Postglacial sealevels on a spherical, self-gravitating viscoelastic earth: effects of lateral viscosity variations in the upper mantle on the inference of viscosity contrasts in the lower mantle, *Earth Planet. Sci. Lett.*, 211, 57-68, 10.1016/S0012-821x(03)00199-7, 2003.


Wu, P., Wang, H., and Schotman, H.: Postglacial induced surface motions, sea-levels and geoid rates on a spherical, self-gravitating laterally heterogeneous earth, *J. Geodyn.*, 39, 127-142, 2005.

Wu, P., Steffen, H., and Wang, H. S.: Optimal locations for GPS measurements in North America and northern Europe for constraining Glacial Isostatic Adjustment, *Geophys J Int*, 181, 653-664, 10.1111/j.1365-246X.2010.04545.x, 2010.

1425 Wu, P., Wang, H. S., and Steffen, H.: The role of thermal effect on mantle seismic anomalies under Laurentia and Fennoscandia from observations of Glacial Isostatic Adjustment, *Geophys J Int*, 192, 7-17, 10.1093/gji/ggs009, 2013.

Yuen, D. A., Sabadini, R. C. A., Gasperini, P., and Boschi, E.: On Transient Rheology and Glacial Isostasy, *Journal of Geophysical Research*, 91, 1420-1438, DOI 10.1029/JB091iB11p11420, 1986.

1430 Zhong, S. J., Paulson, A., and Wahr, J.: Three-dimensional finite-element modelling of Earth's viscoelastic deformation: effects of lateral variations in lithospheric thickness, *Geophys J Int*, 155, 679-695, 10.1046/j.1365-246X.2003.02084.x, 2003.



1435

Figure 1: Components of a GIA model. Surface loading by ice sheets and the ocean, together with Earth properties, govern how the solid Earth deforms. Changes to the gravity field (determined by solving the sea-level equation) define how meltwater is redistributed across the oceans. Comparing model outputs to observations allows model inputs to be adjusted to achieve a better fit. Solid Earth deformation will affect ice-sheet evolution; this can be modelled with a coupled model (Sect. 3.1). Numbers refer to relevant sections in the text.

1440





Figure 2: Solid Earth deformation and sea-level change. a) Ice sheet losing mass results in solid Earth rebound and a decrease in sea surface height due to the decreased gravitational attraction of the ice sheet. Both processes cause near-field relative sea-level fall. Relative sea level rises in the far field due to the addition of meltwater to the ocean. b) Ongoing solid Earth relaxation after disappearance of the ice sheet. Ocean syphoning is the process whereby peripheral bulge subsidence increases the capacity of the ocean; the result is a fall in mean sea surface height. Solid lines indicate original positions, dashed lines indicate new positions. Figure adapted from Conrad (2013).

1445



1450

Figure 3: Variations in ocean area. a-b) Retreat of marine-grounded ice increases the area of the ocean over which water can be redistributed. b-c) Onlap and offlap changes the areal extent of the ocean. In the near field of a melting ice sheet, rebound results in local sea-level fall, causing the shoreline to migrate offshore (offlap). In the far field of a melting ice sheet, sea-level rise causes the shoreline to migrate onshore (onlap). t_1 , t_2 , t_3 refer to the times represented in a) b) and c). Figure adapted from Farrell and Clark (1976).

1455






Figure 4: Rotational feedback. a) Earth's rotation vector moves towards a region of mass loss, causing a change in the shape of the solid Earth and the geoid. Relative sea level rises and falls in opposing quadrants of the Earth. b) Polar ice loss results in a decrease in the oblateness of the Earth (J_2). Solid lines indicate original positions, dashed lines indicate new positions.

1460



1465

Figure 5: Non-uniqueness in GIA modelling. a) Trade-off between the timing and the magnitude of past surface load change: large ice loss at 10 ka can result in the same present-day uplift rate as smaller ice loss at 5 ka. b) Trade-off between ice load history and Earth rheology: large ice loss combined with a weak rheology can produce the same present-day uplift rate as small ice loss combined with a strong rheology.



1470

Figure 6: GIA-ice dynamic feedbacks. a) Far-field ice melt leads to local sea-level rise which causes a retreat in the position of the grounding line (the point at which an ice sheet starts to float). b) Near-field ice melt leads to solid Earth rebound which, combined with a decrease in gravitational attraction, results in local sea-level fall. This has a stabilizing effect on the ice sheet and results in grounding line advance. Solid lines indicate original positions, dashed lines indicate new positions.

1475




Figure 7: Effect of sediment redistribution on GIA. a) Sediment erosion and deposition results in solid Earth deformation and a local reduction in water depth. The net redistribution of mass (solid Earth and sediment) will also change the shape of the geoid, and hence local sea level. b) Sediment compaction over time results in a small increase in local water depth. Solid lines indicate original positions, dashed lines indicate new positions.

1480