

Response to Reviewer Comments

Response to RC1 (Anonymous Referee 1)

- 5 We thank the anonymous referee for the time and effort they spent on reading, reviewing and commenting our manuscript. We were able to improve the manuscript by following their suggestions and thinking and acting on their comments. In this communication, we highlight our responses in blue.
- 10 It does not get entirely clear which part of the methodology can be considered as original to this manuscript. It would be good to clarify that.

The individual components of the whole procedure are established methods, but the combination of methods was tailored specifically to the scientific problem at hand. We made sure to clarify this in the methods section (L136-138), and mention examples of where else this chain of processing may be used in the discussion (L677-687).

- Minor comments:
- 20 - Abstract: in the abstract the authors mention tipping points in what I consider the motivation of the research. This, in my opinion is misleading (even though the method might be used for that purpose) since the rest of the manuscript is not about tipping points. Better mention the effects on land surface processes like erosion already as motivation in the abstract (like it is done nicely in the introduction).

We thank the reviewer for noticing and highlighting this inconsistency and agree. We now omit the mention of tipping points and focus the reader on surface processes/erosion from the beginning.

- 25 - For the motivation, as well as in the discussion section it could be mentioned that this methodology might be pretty useful regarding the effects of future climate changes and the impacts on Earth surface processes.

30 This is an excellent point and may remove barriers for climate impact researchers to apply a similar method for future climate change. We make a note of that in our manuscript now (L125-127 and L677-678).

- 35 - Introduction line 70: Please do not call modern pre-industrial, since there is distinct differences and this might lead to confusions

We thank the reviewer for raising this very valid point and made sure we do not mix those two up throughout the entire manuscript.

- 40 - Introduction lines 79/80: it is not entirely clear to me why you mention uplift histories here. Could you maybe clarify that? Could that methodology help to re-evaluate uplift histories?
- We thank the reviewer for pointing this out. We originally mentioned uplift histories, because reconstructions and interpretations of erosion histories impact the refinement of uplift histories. We realise now that the mention of uplift histories is confusing at this point in the manuscript and omitted it.

- 45 - Paragraph 2.3./line 202: add "climate" to "Each variable" => "Each climate variable"
50 This has been corrected.

- Line 230: given by the models

This has been corrected.

- 55 - Line 461: impacts on physical weathering
This has been corrected.

- Discussion: please mention more often in the discussion where the different named regions are situated (e.g. mountains, lowlands etc).

- 60 We added more such descriptive terms to the discussion when we mention specific regions/clusters (e.g. L554, 576, 590).

- Conclusion: line 543: please do not only mention k, but also what it means (climate anomaly clusters?) here, since there is people around reading only the conclusions and the abstract:

- 65 This is a very good point. We instead write “assigned number of anomaly clusters” (L659), but still mention k in brackets to help those who have read the entire manuscript make that connection more easily.

- Line556: coincide with decreases in

- 70 This has been corrected.

- Conclusion: It would be good to mention future applicability of the method like using it for the investigation of other physical processes or other time slices (give examples!).

- We thank the reviewer for the suggestion. We mention at the end of the discussion that the presented methods may be applied to future climates. Additionally, we give a possible example of method transfer to a different research field and highlight the important conditions of merited use of the method. (L677-687)

80

85

90

95 Response to RC2 (Andrew Wickert)

We thank Andrew Wickert for his time and effort as a referee. We appreciate his thoroughness, comments and suggestions and think we were able to improve our manuscript by acting on those. In this communication, we highlight our responses in blue.

100

16, 21, etc. I think that there should be emdashes instead of hyphens between PI and MH, LGM, and PLIO

This has been corrected throughout the entire manuscript.

105

24. explained → described.

We use the term “explain” in a statistical sense, as is consistent with the language used to describe the results of the discriminant analysis. The LMD is an explanatory tool. However, we acknowledge that this may create some confusion in the abstract. We therefore now explicitly state in the abstract that this explanation is a statistical one.

110

48–49. Modern climate is not always assumed, though. As an example with which I am quite familiar, we (Wickert et al., 2019) actually used plausible river discharges based on glacial-stage conditions including ice-sheet melt and precipitation minus evapotranspiration (Wickert, 2016) to understand incision of the Mississippi River. I strongly agree that using the appropriate past climate is important!

115

But I am wondering if “often have to be interpreted” is a bit too strong in general – from paleo-proxy data, we generally do know something about how climate varied. But perhaps what you mean to say is that there is not a globally-spanning resource to do this and ingest it into models, which could in fact be closer to your aim (?) and a stronger way to state this. p.2 and 3 overall: Consider ways to split this 1.5-page single paragraph – perhaps between background and what you do. (Even if it is supposed to be split at the list and I just don’t see it, the p. 2 single paragraph definitely could benefit from being broken up into its sub-topics.

120

We thank Andrew Wickert for this comment. Indeed, we could have been clearer in this regard. The advantages of GCM’s are (1) having a globally-spanning resource, and (2) a way to relate observations of palaeoclimate back to processes and drivers, since GCM’s are based on our physical understanding of the climate system, (3) they are a tool suitable for sensitivity experiments, (4) they complement proxy-based reconstructions with regional averages and additional climate variables to offer a more complete picture of climate. We agree that our original phrasing is a little strong. We adjusted this and offer a clearer motivation for using GCM’s for these purposes. Additionally, we split the paragraph into three parts: (1) General motivation for the work and using GCM output, (2) previous palaeoclimate modelling work, and (3) what we do here. (L54-64)

125 79. “lies on” seems awkward to me. Why not, “We focus on 4 regions...”

Many thanks for the suggestion. We changed this in our text.

130

93, 95. Earth-surface here is a compound noun, and so should be hyphenated.
This has been corrected.

It also seems that 93-95 is an accidental self-standing one-sentence paragraph
This has been corrected.

135

99. surface-process-relevant (compound noun)
This has been corrected.

140

112. Consider updating the PMIP reference, as new PMIP projects have run since 2012 – we are now on PMIP4! The following reference is an experiment overview. But in addition to just citing, I think that their science and model outputs could be quite useful to you and your future work.
Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclaus, J. H., Otto-

Bliesner, B. L., ... Zhou, T. (2018). The PMIP4 contribution to CMIP6 – Part 1: Overview and over-arching analysis plan. Geoscientific Model Development, 11(3), 1033–1057.

150 https://doi.org/10.5194/gmd-11-1033-2018

We updated the reference in the text, but still included the Bracannot reference since our simulations were based on simulations and protocols from the third phase. We agree that PMIP4 will be very relevant to us and certainly plan to take advantage of PMIP4 in future work.

155 119. Consider starting a new paragraph at “Mutz”

We agree that this gives the text more structure and followed the suggestion.

142. “of interest to the Earth surface”: rephrase. I know what you mean, but Earth’s surface doesn’t have the mental capacity for interests.

160 We added a hyphen to clarify that we speak of the Earth-surface communities (and palaeoaltimetry communities).

143. communities (plural)

This has been corrected.

165 144-145. hyphens to emdashes (indicating range)

This has been corrected.

163-170. In your place, I would have added evapotranspiration (or P-ET) to this list, as the sum of latter across watershed is what should control river discharge. Could you explain your decision not to include this?

We refrained from using several potentially relevant variables due to their poor representation in GCM’s on the scale we are working on. GCM’s don’t represent the regional hydrology well due to the parameterisations in their land surface schemes. Routing GCM’s through more sophisticated hydrological models or regional climate models would allow the inclusion of such variables. We changed the text to be clearer about our choice of variables (in “Data and Methods”), and added those limitations (and suggestions) to the “Comments on methodical implications” section.

175 175. New paragraph at “clustering”.

This has been corrected.

178-180. I think that the writing would be clearer if you focus on this criterion before the preceding sentence, in which you give the appropriate values. So, methods ! results.

We re-arranged these sentences, so that the order of information makes more sense.

185 203. I’m a bit confused here by the mention of the number of cluster members. I had thought that the clusters themselves were not yet defined. Or by this, do you mean “grid boxes”? Or “groups selected and then tested”?

We thank Andrew Wickert for pointing out the confusing nature of this sentence. As summed up in Fig. 1, we carry out the clustering (and find the optimal cluster number k) before we conduct the LMD. However, we refer here to the number of cluster members, i.e. the number of elements or grid boxes in each cluster. We realise now that we have not used the term “cluster members” before and replaced it by “elements in each cluster” to avoid any such confusion.

195 Equation 4 seems unnecessary – perhaps you can just write that you maximized.

We acknowledge the redundancy and overlap between equation 3 and 4. However, we think equation 4 nicely summarises the whole problem in a way that you also may find it summarised in text books, thus making it easier to recognise the same thing again when reading up on LMD. We therefore decided to leave out equation 3 and keep equation 4 (now 3).

200 Equation 6 seems strange for multiple reasons. But now I realize that what looks like strange matrix multiplication is actually really several separate matrices. Please also ensure that you have reasonable

- formats, including consistent positioning and orientation of ellipses, brackets that look like brackets long brackets (this is more for typesetting, but these look like you just stretched short brackets, thus ending up with long knobs on the ends), etc. The latter comments are for style, but the former are for content.
- 205 6 (now 5 in the revised manuscript) merely expands and explains the terms of the previous equation, which is a common notation to shorten the notation for multiple regressions (cf. next comment response). It may not be necessary to include this to expand on what is already stated before, but we hope that this will look more familiar to some of the readers. We reworked the formatting.
- 210 224. Could you describe your solution technique in more detail? I do not understand why you needed to partially differentiate the equation (or with respect to what) in order to solve it.
It would involve partial differentiation with respect to discriminant coefficients, which can be seen as regression coefficients if we think of it as a multiple regression for the elements on the discrimination axis. We clarified this and inserted a reference for more details, since this is a well-established method.
- 215 224. Could you explain mathematically what you mean by “standardized”? This is the point at which I no longer have enough information to understand or reproduce your approach. In addition, I quite dislike two-letter variables: it is then quite difficult to know if they represent two variables that should be multiplied together, or a single variable.
We thank Andrew Wickert for these comments. We use the term “standardised” in a statistical sense, i.e. we put the value in relation to the standard deviation of the respective variable. We clarified this.
- 220 225. We agree that using two-letter variables can lead to confusion. We changed the variable names to one-letter variables and updated them throughout the manuscript and on our figures.
- 230 253. Please read this paragraph and split it at the point at which “Erosion rates...” is no longer a good topic sentence. I suggest at, “The anomaly cluster”.
We agree that this gives the text more structure and followed the suggestion.
- 235 262-263. The “Late Pliocene” sentence seems a bit unnecessary – if you want to add an example, consider using something with a reference to demonstrate that this would be the case.
We decided to remove this sentence as it is somewhat distracting and unnecessary.
316. Argentina, AND THE central and...
This has been corrected.
- 240 364. endashes
This has been corrected.
364. “decreases in 2m air temperature AND freeze-thaw days, and”: separate decreases and increases. In general, I suggest that you use the Oxford comma; it can disambiguate your complex lists.
245 We appreciate the suggestion and made changes accordingly.
366. These cells are covered by more than just the Scandinavian ice sheet.
We updated the text to make sure we include the British Ice Sheet.
- 250 399. PLIO!MH?
Yes, this should read “MH”. We corrected it.
423. incl. The!including the
This has been corrected.
- 255 424. temperature, and WITH (required to split lists)
This has been corrected.

- 458-459. large-scale
260 This has been corrected.
462. and→as well as to (required to split lists)
This has been corrected.
- 265** 490-494. Consider noting something about changes in evapotranspiration vs. changes in erosion rate due to the physical presence of vegetation (e.g., roots)
We have included a note on vegetation's modification of the hydrological cycle, and modifying hillslope stability by changing root characteristics.
- 270** 525. erosion-relevant
This has been corrected.
525. By “high” do you mean small spatial, short temporal, both, neither?
We mean both and clarify this now in the text.
- 275** Data availability: I suggest that you obtain a DOI for your model outputs.
As we understand, it is not possible to obtain a doi for self-hosted data, and the size of the original model output is currently still too large for hosting services.
- 280**
- 285**
- 290**
- 295**

List of Relevant Changes

300 Following the suggestions of the reviewers, we clarified many details regarding our method, emphasised the motivation of a GCM (General Circulation Model) based study, gave the manuscript more structure and clarity overall, listed further limitation and suggestions, and highlighted possibilities of method transfer. Furthermore, we updated our figures to accommodate a suggestion for change in notation.

305

310

315

320

325

Detection and Explanation of Spatiotemporal Patterns in Late Cenozoic Palaeoclimate Change Relevant to Earth Surface Processes

330 Sebastian G. Mutz¹, Todd A. Ehlers¹

¹ Department of Geosciences, University Tübingen, D-72074 Tübingen, Germany

Correspondence to: Sebastian G. Mutz (sebastian.mutz@uni-tuebingen.de)

Abstract

335 Detecting and explaining differences between palaeoclimates can provide valuable insights for Earth scientists investigating processes that are affected by climate change over geologic time. In this study, we describe and explain spatiotemporal patterns in palaeoclimate change that are relevant to Earth surface scientists. We apply a combination of multivariate cluster- and discriminant analysis techniques to a set of consistently set-up high-resolution palaeoclimate simulations conducted with the ECHAM5 climate model. A pre-industrial (PI) climate simulation serves as the control experiment, which is compared to a suite of simulations of Late Cenozoic climates, namely a Mid-Holocene (MH, ca. 6.5 ka), Last Glacial Maximum (LGM, ca. 21 ka) and Pliocene (PLIO, ca. 3 Ma) climate. For each of the study regions (Western South America, Europe, Himalaya-Tibet and South Alaska), differences in climate are subjected to geographical clustering to identify dominant modes of climate change and their spatial extent for each time slice comparison (PI-MH, PI-LGM and PI-PLIO). The selection of climate variables for the cluster analysis is made on the basis of their relevance to Earth surface processes and includes 2m air temperature, 2m air temperature amplitude, consecutive freezing days, freeze-thaw days, maximum precipitation, consecutive wet days, consecutive dry days, zonal wind speed and meridional wind speed. We then apply a two-class multivariate discriminant analysis to simulation pairs PI-MH, PI-LGM and PI-PLIO to evaluate and explain the discriminability between climates within each of the anomaly clusters. Changes in ice cover create 340 the most distinct and stable patterns of climate change, and create the best discriminability between climates in western Patagonia. The distinct nature of European palaeoclimates is statistically explained mostly by changes in 2m air temperature (MH, LGM, PLIO), consecutive freezing days (LGM) and consecutive wet days (PLIO). These factors typically contribute 30%-50%, 10%-40% and 10%-30% respectively to climate discriminability. Finally, our results identify regions particularly prone to changes in precipitation-induced erosion and temperature-dependent physical weathering.

Deleted: valuable insight into climatic tipping points and a useful framework of information

Deleted: al

Deleted: and

Deleted: -

Deleted: -

Deleted: -

Deleted: -

Deleted: -

Deleted: -

Deleted: mostly

Keywords: Cenozoic climate, climate change, ECHAM5, Last Glacial Maximum, Mid-Holocene, Pliocene, discriminant analysis, discriminability, Earth surface processes, erosion

360 1. Introduction

In the study of Earth surface processes, gaining new quantitative understanding of the atmosphere's interaction with the Earth's surface through erosional processes is limited by the difficulty of establishing reliable palaeoclimatic context for erosion rate histories. Such context is particularly useful when erosion rates are calculated using techniques such as cosmogenic radionuclides and low-temperature thermochronology [e.g. 365 Schaller et al., 2002; Bookhagen et al., 2005; Moon et al., 2011; Insel et al., 2010; Stock et al., 2009], which

integrate over timescales of 10^3 - 10^{6+} years. Despite recognition of the influence of climate on tectonic processes and landscape evolution through erosion [e.g. Whipple et al., 1999; Montgomery et al., 2001; Willett et al., 2006; Whipple, 2009; Deal et al., 2018], erosion rates calculated from geo- and thermochronological archives are often

Deleted: often have to be

380 interpreted under the assumption of modern climate due to insufficient palaeoclimate data [e.g. Starke et al., 2017]. While proxy-based palaeoclimate reconstructions, or reconstructions of climate-controlled variables such as river

discharge [e.g. Wickert, 2016], are in some cases able to provide sufficient and plausible context for specific problems, GCM's (general circulation models) offer a complementary and integrative approach to palaeoclimate reconstructions. GCM's complement proxy-based reconstructions in several ways: (1) GCM's have a global coverage [e.g. Salzmann et al., 2011; Haywood et al., 2013; Jeffrey et al., 2013] and therefore provide palaeoclimatological context in regions with sparse proxy records; (2) GCM-based palaeoclimate reconstructions allow the refinement of local proxy-based reconstructions by providing regional means and a broader climatic context; (3) GCM's are able to offer insight into atmospheric drivers of reconstructed local palaeoclimates, because they simulate atmospheric processes based on our physical understanding of the climate system. (4) GCM's allow the conduction of sensitivity experiments to investigate the relationship between climatic drivers and local observations [e.g. Takahashi and Battisti, 2007].

390 The Paleoclimate Modelling Intercomparison Project (PMIP) coordinates palaeoclimate modelling efforts [Kageyama et al. 2018] and provides experiment designs for the Mid-Holocene and the Last Interglacial [Otto-Btiesner et al., 2017], the last millennium [Jungclaus et al., 2017], the Last Glacial Maximum [Kageyama et al., 395 2017], and the Pliocene Warm Period [Haywood et al., 2016.], which is part of the Pliocene Model Intercomparison Project (PlioMIP). Despite consistent forcings, experiments carried out with different GCM's and model resolutions yield different results due to GCM specific parameterisation. [Mutz et al., 2018] conducted PMIP-style palaeoclimate experiments with the same GCM (ECHAM5) and resolution, which removes the GCM parameterisation related signal in the differences between simulated palaeoclimates. This experiment framework comprises climate simulations for the pre-industrial (PI, reference year 1850), Mid-Holocene (MH, ~6ka), Last Glacial Maximum (LGM, ~21ka) and Pliocene (PLIO, ~3Ma).

400 This study takes advantage of the Mutz et al. [2018] simulation framework and quantifies differences between simulated palaeoclimates with regard to variables relevant to Earth surface processes (e.g., rainfall characteristics, temperature derived quantities, wind speed and direction) in order to allow more refined interpretations of potential climatic drivers for changing rates in Earth surface processes. For more effective communication of our methods and results, we separate the PI control simulation from MH, LGM and PLIO in discussion by referring only to the latter three as Late Cenozoic climates. These are time periods over which reconstructed erosion rates 405 typically integrate. Understanding how different these palaeoclimates are from a pre-industrial climate with regard to variables that potentially affect erosion rates is essential in any comprehensive and merited interpretation of erosion rates, and ultimately allows for better assessment of the influence of climatic and tectonic controls on erosion. Three questions are addressed in this study:

- 410 1. What are the spatial patterns of climate change in comparisons of pre-industrial with Late Cenozoic time periods?
2. How different were Late Cenozoic palaeoclimates to the pre-industrial climate with regard to variables relevant to Earth surface processes?
3. What constitutes these quantified differences between pre-industrial and Late Cenozoic climates?

Deleted:

Deleted: part of this palaeoclimatic context, GCM's (general circulation models) offer a complementary and integrative approach to palaeoclimate reconstructions that yield a more complete picture of paleoclimate [e.g. Salzmann et al., 2011; Haywood et al., 2013; Jeffrey et al., 2013] and allow better investigation of atmosphere dynamics by conducting sensitivity experiments [e.g. Takahashi and Battisti, 2007]. This study takes advantage of the benefits of GCM's and quantifies differences between simulated palaeoclimates with regard to variables relevant to Earth surface processes (e.g., rainfall characteristics, temperature derived quantities, wind speed and direction) in order to allow more refined interpretations of potential climatic drivers for changing rates in Earth surface processes.¶

Deleted: This study is based on a suite of PMIP-style palaeoclimate experiments

Deleted: modern (

Deleted:)

We focus on 4 regions that are frequently investigated with regard to erosion histories: South Alaska, Western South America, Himalaya-Tibet and Europe. The first question is addressed by conducting a cluster analyses on the differences between pre-industrial and Late Cenozoic palaeoclimates, which subdivides the 4 study regions

Deleted: The geographical focus of this study lies on

Deleted: and uplift

440 into geographical clusters governed by a distinct character of erosion relevant climate change. Whereas Mutz et al. [2018] apply a similar cluster analysis to describe the modes of climate variability in each palaeoclimate simulation, the results of this study consist of maps showing the extent of a particular mode of climate change, and thus provides an overview of climate change over time. The resulting clusters also serve as suitable masks for values used in the discriminant analyses.

Deleted: 1

445 Questions 2 and 3 are addressed by conducting discriminant analyses on subdivisions of the 4 study areas, which are objectively pre-defined by the aforementioned cluster analyses. The results provide a quantitative assessment and explanation of differences in climate with regard to variables relevant to Earth surface processes. The third question is answered by conducting a cluster analyses on the differences between pre-industrial and Late Cenozoic palaeoclimates, which yields the extent of regions governed by a distinct character of erosion relevant climate change.

Deleted:

450 The overarching goal is to provide the Earth surface science community with an overview and quantitative assessment and explanation of how climate changed in the Late Cenozoic with regard to variables relevant to Earth surface processes. However, the same methods may also be applied to simulations of modern and future climates to detect and explain patterns of climate change that may result in shifts of Earth surface process regimes.

Deleted:

455 2. Data and Methods

This section describes the data basis, methods and processing steps. In summary, we apply a cluster analysis that identifies where surface process-relevant aspects of climate change are likely, and a discriminant analysis that quantifies and explains these changes within the regions identified by clustering. This combination of clustering

Deleted:

Deleted:

460 and discrimination of climate model simulation results yields 5 different sets of results (Fig. 1): Anomaly maps for a set of climate variables (in the supplemental material to this manuscript), multivariate anomaly cluster maps and anomaly cluster characterisations (section 2.2), discrimination scores and a measure of relative contribution by each climate variable to discriminability (section 2.3). While the individual components of the processing chain (Fig. 1) are based on well-established methods, the processing set-up and the particular combination of these methods is tailored to address this study's specific scientific problems.

465

2.1 ECHAM5 simulations

General Circulation Models (GCM's) simulate global climate based on our physical understanding of atmospheric processes and are primarily used to investigate atmospheric dynamics and contemporary climate change, but have also been applied to improve our understanding of past climates and Earth system dynamics [e.g. Kutzbach et al.,

470 1993; Ehlers and Poulsen, 2009; Maroon et al., 2015, 2016, Mutz et al., 2016, 2018]. GCM's have become well established tools in geoscience, as is reflected by the work of the Palaeoclimate Modelling Intercomparison Project (PMIP) [Kageyama et al. 2018, Bracannot et al., 2012], which adds palaeoclimate related contributions to the Coupled Model Intercomparison Project (CMIP). Palaeoclimate studies address a range of different time scales including the last millennium [e.g. Jungclaus et al., 2010], orbital [e.g. Gong et al., 2013; Lohmann et al., 2013;

475 Pfeiffer and Lohmann, 2016; Wei and Lohmann, 2012; Zhang et al., 2014] and tectonic time scales [e.g. Knorr et al., 2011; Stepanek and Lohmann, 2012]. Comparisons between palaeoclimate simulations of different time

periods are often complicated by use of different models and inconsistencies in model setup. GCM specific
485 parameterisations and differences in horizontal and vertical resolution introduce differences between simulations
that are not a result of prescribed forcings.

Mutz et al. [2018] conducted a suite of GCM simulations conducted with the same GCM (ECHAM5) and
resolution to circumvent the above-mentioned biases. This GCM simulation framework comprises palaeoclimate
490 experiments for pre-industrial times (reference year 1850), the Mid-Holocene (ca. 6.5 ka), the Last Glacial
Maximum (ca. 21 ka) and the Pliocene (ca. 3Ma) climates. The experiments were conducted at a spectral
resolution of T159 (ca. 80km x 80km), with 31 vertical levels and an output frequency of 1 day. The simulations
are based on boundary conditions from coupled AOGCM transient simulations and paleoenvironmental
reconstruction initiatives such as PMIPIII [Abe-Ouchi et al 2015], GLAMAP [Sarnthein et al. 2003], CLIMAP
[CLIMAP project members, 1981], PRISM [Haywood et al., 2010; Sohl et al., 2009; Dowsett et al., 2010] and
495 BIOME6000 [Prentice et al., 2000; Harrison et al., 2001; Bigelow et al., 2003; Pickett et al., 2004]. For detailed
description of model setups, we refer the reader to Mutz et. al [2018] and references therein. The high resolution
and model consistency across all these time slices has not been achieved previously in the palaeoclimate modelling
community. These GCM experiments therefore represent a unique, state-of-the-art simulation framework suited
for investigations of changes in climate controlled processes across the Late Cenozoic. The GCM (ECHAM5) is
500 a well-established model in the climate community, and simulated palaeoclimates are in agreement with other
modelling and proxy-based reconstruction efforts. Mutz et al. [2018] use a present-day simulation to establish
confidence in the model, and compare palaeoclimate estimates to compilations of proxy-based reconstructions for
MH and LGM precipitation over South America and Tibet, revealing an overall satisfactory performance of the
GCM. The latitudinal gradients and magnitude of difference in temperature and precipitation are in good
505 agreement with results of previous palaeoclimate modelling efforts. We refer the reader to Mutz et al. [2018] for
a detailed comparison to other palaeoclimate simulations and proxy-based reconstructions.

Deleted: In order to avoid these biases,

2.2 Clustering – Multivariate Anomaly Maps

This study's investigation of differences focusses on 4 regions, which (a) are of interest to the Earth surface and
510 palaeo-altimetry communities, and (b) are feasible to work on given the GCM's limitations [cf. Mutz et al., 2018]:

South Alaska (52° - 68° N, 125° - 165° W), Western South America (5° - 56° S, 60° - 80° W), Europe (26° - 65° N,
515 22° W- 66° E) and Hiamaya-Tibet (0° - 60° N, 40° - 120° E). Climate change within each of the investigated region

is neither spatially homogenous, nor is it the same in magnitude everywhere [Mutz et al., 2018]. In the example
of PI-LGM climate change in Europe, southern Norway experiences a strong increase in consecutive freezing

520 days and strong decrease in 2m air temperature, while continental Europe experiences only mild cooling, but
strong increases in intra-monthly 2m air temperature variability. We refer to these combinations of changing
climate attributes as modes of (climate) change in this manuscript. Each of the regions is governed by a number
of these distinct modes of climate change. These intra-regional discrepancies merit an informed and objective

subdivision of each region into geographical subdomains governed by one of these modes of change prior to the
investigation of differences in climate through time. Geographical clustering allows such subdivisions on the basis

of similarities in climate change at different locations within each region. It assesses climate change for each grid
box, calculates its similarity to climate change in other grid boxes, and groups them accordingly. More
specifically, it allows the grouping of elements (*i*), in this case climate model surface grid boxes, by the co-

Deleted:

Deleted: community

Deleted: -

variability of anomalies of selected climatic element attributes. For each region, the contained elements are subjected to agglomerative hierarchical clustering, followed by k-means clustering corrections [Mutz et al., 2016; Paeth, 2004] to address the inherent shortcomings of a pure hierarchical approach. The Mahalanobis distance [e.g. Wilks, 2011] is used as a measure of similarity (of climate change) between clusters in the entire procedure.

540 Readers are referred to Mutz et al. [2018] for a more detailed description of the procedure and aforementioned shortcomings.

The clustering is conducted on basis of the same climatic attributes as the discrimination between climates through time (section 2.3). In total, M ($= 9$) element attributes, summarised in table 1, are chosen based on (1) their relevance to Earth surface processes, (2) their adequate representation by the GCM, and (3) the feasibility of construction from GCM output: near-surface air temperature (te2m), intra-monthly near-surface air temperature amplitude (t2am), consecutive freezing days (csfd), freeze-thaw days (fthd), maximum precipitation (pmax), consecutive wet days (cswd), consecutive dry days (csdd), zonal near-surface wind speeds (u10) and meridional near-surface wind speeds (v10).

550 For calculation of consecutive freezing days and freeze-thaw days, a surface temperature of 0°C is taken as a threshold value. The maximum duration of a wet period, i.e. precipitation exceeding 1mm/day [Zolina et al., 2010; Zhang et al., 2011] constitutes the cswd attribute. Inversely, the maximum duration of a dry period, i.e. precipitation failing to exceed the 1mm/day threshold [Zin and Jemain, 2010] constitutes the csdd attribute. These attribute variables are constructed from each palaeoclimate (MH, LGM and PLIO) and reference simulation (PI) output. The climate attribute anomalies, which serve as a basis for the clustering, are then calculated for time slice comparisons PI-MH, PI-LGM and PI-PLIO.

Clustering requires an a priori decision on the number of clusters (k) or subdivisions per region. The optimal value of k is not known before clustering. Therefore, k is varied in our experiments, and the cut off point for the parameter is set once the increase in k no longer results in a cluster with distinct climatic character, but instead 560 results in a weakened or strengthened character of an already existing cluster. Since optimal k can be expected to roughly scale with region size, the parameter is varied from 3-5 for South Alaska and from 5-8 for the larger regions. The results consist of optimal geographical subdivisions (climate clusters C_1-C_k) with distinct climatic characters, which are described by mean vectors for climate attribute anomalies. Every cluster characterising vector has a length of M . For each of the time slice comparisons and clusters, the discriminability and relative 565 contribution to it by each of the M attribute variables is quantified in the procedure described in section 2.3.

2.3 Discrimination – Quantifying and Explaining Anomalies

The multivariate linear discriminant analysis (LMD) [e.g. Wilks, 2011] is a statistical tool that allows the investigation and explanation of differences between two or more groups with regard to multiple attribute 570 variables. More specifically, it quantifies the discriminability of the groups and the contribution of each of the attribute variables to this discriminability. The resulting discrimination model can be applied to objectively categorise an element with unknown group-affiliation. In this study, the time periods (PI, LGM, MH and PLIO) are used as groups, and the aforementioned climate variables relevant to Earth surface processes are chosen as group attributes. Since the focus of this study lies on the assessment of the differences between two specific time 575 periods with regard to multiple climate variables, the problem is treated as a two-group multivariate case.

The centre piece of the analysis lies in finding a discriminant function that best separates the two groups (or paleo-climate time slices). This is carried out for each comparison, i.e. each pair of paleo-climate time slices. This discriminant function can be expressed as a linear combination of the climate attribute variables:

$$Y = v_0 + v_1 X_1 + v_2 X_2 + \dots + v_m X_m + \dots + v_M X_M \quad (1)$$

where Y is the discriminant function, X_m ($m=1 \dots M$) are the climate variables used to assess the differences in paleo-climates, v_m ($m=1 \dots M$) are the discriminant coefficients associated with each variable, v_0 is a constant (the y-intercept) that is of no relevance to the goodness of separation and will therefore no longer be mentioned. In this case, $M=9$ (te2m, t2am, csfd, fthd, pmax, cswd, csdd, u10, v10). Each climate variable X_m (cf. table 1) contains elements x_{mn} ($n=1 \dots N$), where N is the number of elements in each cluster. Each element (or grid box) is associated with a discriminant value y_n described by:

$$y_n = v_1 x_{1n} + v_2 x_{2n} + \dots + v_m x_{mn} + \dots + v_M x_{Mn} \quad (2)$$

Deleted: cluster members

595 In other words, the elements y_n ($n=1 \dots N$) are projected onto the discriminant axis Y . The problem of finding a discriminant function that best separates the two groups (or paleo-climates) can therefore also be seen as the process of finding an axis on which the frequency distributions of the projected elements y_n for the two groups show the smallest overlap. Since overlap is a function of distance between groups (**D**) as well as the scatter within them (**S**), the difference between these frequency distributions is described by the distance between the two group 600 centroids (i.e. the group means of the projected elements y_n on the discriminant axis) and the scatter within the group (i.e. sum of squared deviations from the means). This distance, i.e. the discriminant criterion Γ , is maximised in order to find the best discriminant function Y and corresponding discriminant coefficients v_m ($m=1 \dots M$). The problem in the two-group multivariate case of this study can thus be summarised as:

$$\Gamma = \frac{\text{scatter between groups } (D)}{\text{scatter within groups } (S)} = \frac{(\bar{y}_{T1} - \bar{y}_{T2})^2}{\sum_{j=1}^{n_{T1}} (y_{T1j} - \bar{y}_{T1})^2 + \sum_{j=1}^{n_{T2}} (y_{T2j} - \bar{y}_{T2})^2} \rightarrow \max! \quad (3)$$

Deleted: ¶

605 Γ is the discriminant criterion, $T1$ and $T2$ are the two groups (e.g. PI and LGM in the case of time slice comparison PI-LGM) and n_{T1} and n_{T2} are the number of elements in $T1$ and $T2$. We can express the discriminant function as a matrix calculation

$$y = v_0 + v X \quad (4)$$

Deleted: -

Deleted: In order to find the best discriminant function Y and corresponding discriminant coefficients v_m ($m=1 \dots M$), Γ is maximised, so that the problem to be solved in the two-group multivariate case of this study can be summarised as ¶

where

$$y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{bmatrix} \quad v_0 = \begin{bmatrix} v_0 \\ v_0 \\ \vdots \\ v_0 \end{bmatrix} \quad v = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_M \end{bmatrix} \quad X = \begin{bmatrix} X_{11} & X_{12} & \dots & X_{1M} \\ X_{21} & X_{22} & \dots & X_{2M} \\ \vdots & \vdots & \ddots & \vdots \\ X_{N1} & X_{N2} & \dots & X_{NM} \end{bmatrix} \quad (5)$$

Formatted: Font:

Deleted: In order to solve the problem,

Deleted: we

Deleted: 5

Deleted: W

Deleted: 6

Deleted: it

Deleted: differentiation

Deleted: v

~~standard deviations of respective variables~~, to yield ρ_m . The relative contribution (ρ_m) of each of the M attribute variables to discriminability is calculated as:

$$\rho_m = \frac{\omega_m}{M} \quad (\rho_m)$$

Finally, the skill of the resulting discrimination model is evaluated. For this, the association of each element to groups T1 (PI) or T2 (MH, LGM or PLIO) is forgotten, and the elements are re-categorised according to the critical discriminant values of the models [e.g. Bahrenberg et al., 1992]. The fraction of correct classifications (the score) is calculated and used as a measure of “goodness of separation” given by the models. The described LMD procedure is applied to each time slice comparison T1-T2 (namely PI-MH, PI-LGM and PI-PLIO) and each of the k climate anomaly clusters (C_1, C_2, \dots, C_k) in the 4 study regions. Each calculation yields two variables suitable for addressing the problems treated in this study: 1) a measure for goodness of discriminability (score) and 2) a measure for the relative contribution (ρ_m) of each of the M attribute variables to discriminability. A maximum score of 1 indicates perfect separation of all values, whereas a score of 0 indicates that the discrimination model has no explanatory power at all. Attribute variables associated with a ρ_m value of 1 are solely responsible for the discrimination, whereas those associated with a ρ_m value of 0 contribute nothing to the discriminability between climates.

635 **Deleted:**
Deleted: Formatted: Font: Bold
Deleted: us
Deleted: Formatted: Font: Bold
Deleted: vr
Deleted: 7

640 **Deleted:** -
Deleted: -
Deleted: -
Deleted: vr
Deleted: vr
Deleted: vr

645 2.4 Example Problem

In summary, the clustering of anomalies (section 2.2) reveals geographical clusters (or subdivisions in each of the study regions), in which similar climate change occurs, and describes the mode of climate change in each of these clusters. The LMD (section 2.3) then quantifies the discriminability of climates in these clusters and explains it with the climatic attribute variables. The set of results for each time slice comparisons and region therefore consists of 4 components: 1) *Anomaly (Cluster) Maps* that show the spatial extent of dominant modes of climate change; 2) *Anomaly Cluster Characterisation* that consists of mean vectors of climate change within each cluster and describe the mode of climate change experienced in the grid boxes assigned to the same cluster; 3) *Discrimination Scores* that describe the goodness of discriminability of climates within each of the anomaly clusters; and 4) *Relative Variable Contribution*, which describes the contribution of each of the climatic attribute variables to the discriminability calculated for each of the anomaly clusters. How these sets of results may be used in answering questions pertaining to climate driven Earth surface processes is demonstrated in the simplified example below.

Erosion rates were calculated, for example by means of cosmogenic nuclides, for a region in a specific area of interest circled on the map (Fig. 2). Although they are taken as modern (time T1) erosion rates, the signal integrates over ten-thousand years and includes erosion rates at time T2. In order to find out if and how significantly erosion rates may actually have been different at time T2, Fig. 2 is consulted.

660 The anomaly cluster map shows the large-scale spatial patterns of changes in climatic variables relevant to Earth surface processes. Each cluster is associated with a specific mode of climate change, and all locations that fall within it experience this type of climate change. The area of interest lies in cluster C1, so all information not related to C1 and the area of interest are shaded pale grey to remove distractions. The purple-green raster plots reveal the type of climate change associated with C1 and thus with circled area of interest: T2 had a little more

rainfall, a lot more (consecutive) wet days, higher temperatures, fewer freezing days and fewer (consecutive) dry days. Does the climate of T2 have a distinct enough signature to tell it apart from T1? The score is reasonably high, which reveals that the climate of T2 does indeed have a distinct signature and consequently, it suggests a good possibility for a different erosional regime or erosive intensity. In order to assess the consequences of the mode of climate change in C1 for erosion, the circles are examined. Those explain which anomaly of which variable is responsible for the distinct signature that has been detected and described above. They indicate that ca. 60% of this “discriminability” can be explained by increases in temperature and ca. 40% by increases in consecutive wet days. Consequently, it is reasonable to assume a priori that erosive processes controlled by these parameters may be affected significantly. The specific sensitivity of conditions and processes at the Earth’s surface, e.g. vegetation or critical thresholds in landscape responses, cannot be taken into consideration in these results due to the highly variable nature of it. However, the exact magnitude of those significant (and insignificant) changes are listed in supplemental table T1 to allow the reader to assess the specifics of the impact of climate on processes of the specific area and problem at hand. The above example provides the basic concepts needed to understand the remaining figures in this manuscript.

Deleted: Conditions were warmer and wetter as would be the case for many regions in the Late Pliocene for example.

3. Results

This section contains descriptions of results from the cluster- and discriminant analyses carried out for each of the time slice comparisons and study regions. It is divided into sections for the three major study regions: 3.1 Western

695 South America, 3.2 Europe, and 3.3 Himalaya-Tibet. For brevity, the results for South Alaska are included in the supplemental material (S10). For each subsection, regional results for time slice comparisons PI-MH, PI-LGM and PI-PLIO are presented. The figures accompanying each of those consists of four elements (cf. Section 2.4):

Deleted: -
Deleted: -
Deleted: -

700 1) The first element to the figures are multivariate anomaly maps that were created by clustering the differences between climatic attribute variables in each of these comparisons (cf. Section 2.2) and show the resulting geographical subdivision into clusters governed by a specific mode of climate change. 2) The second element to the figures is (purple and green) raster plots showing the characterisation of those clusters. They describe the mode of change, i.e. the magnitude and direction of changes observed on average in each of the clusters. For visual clarity, the magnitude of change is scaled by the maximum absolute difference in each region and time slice comparison. Numerical values are listed in tabular format in the supplementary material to this manuscript (T1).

Deleted: or

705 3) The third (grey) element consists of the scores for each cluster. These are based on a discriminant analysis carried out for the cluster (cf. Section 2.3) and describe the goodness of discriminability of the palaeoclimates in comparison to the PI control simulation. 4) The fourth and final element to the figures consists of a measure for the relative contribution (α) of each of the 9 climate variables (table 1) to the overall discriminability between two time slices in each geographical cluster. This element is displayed as a layer of circles on top of element 2 and reveals how much an anomaly of a specific variable (represented by a specific shade of purple or green) contributes to the discriminability between PI and a palaeoclimate in each of the clusters. Larger circle diameters correspond to greater contributions.

Deleted: -
Deleted: -
Deleted: -

710 For brevity, results for these time slice comparisons PI-MH, PI-LGM and PI-PLIO are simply referred to as results for MH, LGM and PLIO respectively. The notation MH-C_i, LGM-C_i and PLIO-C_i is used to refer to results for the ith geographical subdivision (or cluster) in the respective set of results. Description of changes in climate are implicitly discussed in the context of the control simulation and are therefore descriptions of deviations from

the PI climate. Discussions of multivariate anomaly maps and cluster characterisation (elements 1 and 2) focus mostly on the stable and persistent patterns seen in the results, i.e. geographical clusters that are least sensitive to changes in k and keep their distinct character. Discussions of scores and relative contribution to discriminability (¶) focus primarily on clusters with good discriminability. Additional information on climate change on a sub-cluster scale are given in the form of single-variable anomaly maps in the supplementary material to this manuscript (S1-S9).

Deleted: or

3.1 Western South America

735 *Large scale patterns and modes of climate change*

In western South America, the dominant modes of change for the MH are captured in clusters MH-C₃, MH-C₄, MH-C₅ and MH-C₆ (Fig. 3 a,d). MH-C₃ covers much of central Patagonia and is characterised by decreases in consecutive freeze-thaw days and increases in consecutive freezing days and consecutive wet days. MH-C₄ is the mode of change observed in most of Argentina and the central and southern Andes. It consists of relatively small increases in consecutive wet days and freeze-thaw days. MH-C₅ covers most of the tropics in the region and is characterised by decreases in consecutive wet days and maximum precipitation, and relatively small increases in 2m air temperature and consecutive dry days. Relatively large increases in 2m air temperature and consecutive dry days, and decreases in maximum precipitation and consecutive wet days constitute the mode of change for MH-C₆, which extends over low-altitude subtropics in the region.

Deleted: ,

740 LGM-C₁ – LGM-C₃ share a number of characteristics. These modes of changes include decreases and 2m air temperature and increases in 2m air temperature amplitude and consecutive freezing days (Fig. 3 e). Furthermore, the region occupied by LGM-C₁ – LGM-C₃ is covered by ice in the LGM. Differences between these modes of changes include a large increase in maximum precipitation in LGM-C₁ and decreases in consecutive wet days in LGM-C₂. LGM-C₄, covering much of the subtropics in the region, is characterised by relatively little change in 750 all of the investigated variables. LGM-C₅ covers much of eastern Argentina and experiences increases in 2m air temperature amplitude. LGM-C₆ covers much of the tropics of the region and is characterised primarily by decreases in maximum precipitation and consecutive wet days.

The dominant modes of change in the PLIO are described by PLIO-C₁, PLIO-C₂, PLIO-C₃ and PLIO-C₅ (Fig. 3 f). Covering much of eastern Argentina and parts of the central Andes, PLIO-C₁ is characterised primarily by 755 relatively large decreases in consecutive dry days and increases in maximum precipitation and consecutive wet days. The grid boxes assigned to PLIO-C₂, extending over most of the low-altitude tropics and subtropics and the Atacama desert, experience very little change on average. For grid boxes assigned to PLIO-C₃, a decrease in maximum precipitation and consecutive wet days, and increase in consecutive dry days and meridional wind speeds can be observed. PLIO-C₅ extends over much of the central and southern Andes and is characterised by 760 decreases in freeze-thaw days and increases in 2m air temperature and meridional wind speeds. While PLIO-C₆ experiences some of the largest changes, it only covers the Northern and Southern Patagonian Ice Fields and coincides with the reduction of the ice cover in the PLIO simulation.

Discriminability

The discrimination scores (Fig. 3 d,e,f) are highest for the LGM and lowest for the MH, and changes in temperature, consecutive freezing days, maximum precipitation and consecutive dry days are factors that explain much of the discriminability overall. LGM-C₁ – LGM-C₃s have the highest scores. In all 3 clusters, decreases in

- 770 2m air temperature are one of the primary contributors to the discriminability between LGM and PI climate. It explains 40%-50% of the discriminability in LGM-C₁ and 30%-40% in LGM-C₂ and LGM-C₃. With 20%-30% explained discriminability in LGM-C₁ and 30%-40% explained discriminability in LGM-C₃, increases in consecutive freezing days are a second important factor for discriminability between the climates in western Patagonia. Discrimination with PLIO and PI simulations yields the highest scores for PLIO-C₆, which covers the
- 775 Patagonian Ice Fields, and PLIO-C₅. Increases in 2m air temperatures and decreases in consecutive wet days and consecutive dry days explain the discriminability in PLIO-C₆ in equal parts (20%-30%). An increase in temperatures and relatively small decrease in consecutive freezing days explain 20%-30% and 40%-50% of the discriminability in PLIO-C₅ respectively.

780 **3.2 Europe**

Large scale patterns and modes of climate change

MH-C₁ covers Spain and the region east of the Caspian sea (Fig. 4 a) and is associated with the least change in climate attribute variables. MH-C₂, covering areas in western Europe, western Russia and the Mediterranean

- 785 coasts (Fig. 4 a), experiences an increase in maximum precipitation (Fig. 4 d). Ukraine, Poland, much of the Baltic Sea coast and southern Scandinavia are assigned to MH-C₃ and experience decreases in freeze-thaw days. MH-C₄ and MH-C₅ primarily cover northern Africa and are characterised by increases in 2m air temperature and maximum precipitation.

LGM-C₁-LGM-C₄ (Fig. 4 b) are all partially characterised by a decrease in 2m air temperature, and freeze-thaw days, and increases in consecutive freezing days (Fig. 4 e). It should be noted that grid boxes assigned to these

- 790 clusters are covered by the [Fennoscandian and British Ice Sheets](#) in the LGM simulation. LGM-C₅ extends over much of the Mediterranean region, Spain and European North Russia and characterised by increases in 2m air temperature amplitude, consecutive dry days and relatively small increases in freeze-thaw days and decreases in maximum precipitation and consecutive wet days. Most of central Europe, western Asia and North Africa is
- 795 assigned to cluster LGM-C₆ and experiences the least change.

The dominant modes of change in the PLIO are captured in PLIO-C₃-PLIO-C₆ (Fig. 4 c). PLIO-C₃ is a mode of change mostly seen in parts of North Africa and characterised by increases in meridional and zonal wind speeds (Fig. 4 f). In the coastal regions north of it, very little change is seen in the PLIO (PLIO-C₄). PLIO-C₅ covers much of central Europe and experiences decreases in freeze-thaw days and 2m air temperature amplitude, and relatively small increases in 2m air temperature. European Russia and parts of Scandinavia are assigned to PLIO-C₆ and experience increases in freeze-thaw days and 2m air temperature, and decreases in consecutive dry days and 2m air temperature amplitude. PLIO-C₇ is mostly distributed along parts of the Mediterranean, Black Sea and Caspian Sea coasts and characterised by increases in consecutive dry days and 2m air temperature, and a decrease in freeze-thaw days. PLIO-C₈ covers southeastern Norway and the Alps and is characterised by decreases in consecutive freezing days and 2m air temperature amplitude, and increases in 2m air temperature and freeze-thaw days.

Deleted: -

Deleted: ,

Deleted: Scandinavian

810

Discriminability

In all time slice comparisons, changes in 2m air temperature explains most of the discriminability in many of the geographical clusters (Fig. 4 d,e,f). Changes in consecutive freezing days and consecutive wet days are also major contributors to discriminability in the LGM and PLIO respectively. LGM-C₁, LGM-C₂, LGM-C₅ and LGM-C₆ 815 are associated with the highest scores for the LGM. 20%-24% of the discriminability in the clusters can be explained by decreases in temperature, and a similar amount can be explained by increases in consecutive freezing days. Although all PLIO scores are high, the PLIO cluster in central Europe (PLIO-C₅) is associated with the highest value. The discriminability in the cluster can be explained by increases in 2m air temperature (30%-40%), increases in consecutive wet days (20%-30%), decreases in consecutive dry days (10-20%) and decreases in 820 temperature amplitude (10-20%). Discriminability in the high-altitude cluster (PLIO-C₈) can be explained by increases in consecutive dry days (10-20%) and decreases in consecutive wet days (20-30%), maximum precipitation (20-30%) and temperature amplitude (10-20%).

3.3 Himalaya-Tibet

825

Large scale patterns and modes of climate change

The stable patterns for the MH results include MH-C₁ covering the region south of the Himalayan orogen, MH-C₂ covering central India and Southeast Asia, MH-C₄ in the region around the Caspian sea, and MH-C₅ north of the Caspian and Aral Sea (Fig. 5 a). MH-C₁ is characterised by increases in consecutive wet days and maximum precipitation and decreases in consecutive dry days and 2m air temperature (Fig. 5 d). MH-C₂ mostly experiences changes in meridional and zonal wind speeds. MH-C₄ is characterised by increases in 2m air temperature amplitude and consecutive freezing days and decreases in freeze-thaw days. MH-C₅ grid boxes are associated with relatively large increases in freeze-thaw days and smaller increases in consecutive freezing days and 2m air temperature amplitude. 830
LGM-C₁ mostly covers the northernmost parts of the region and the Himalayan orogen (Fig. 5 b). The changes associated with it are decreases in 2m air temperature and increases in 2m air temperature amplitude and consecutive dry days (Fig. 5 e). The modes of change described by LGM-C₂ and LGM-C₃ govern large parts of the region, including the Arabian Peninsula, Iran, central Asia, the Tibetan Plateau and Tarim Basin, Mongolia and parts of China. These regions experience relatively small decreases in temperature. Central India and eastern 835 Southeast Asia (LGM-C₄) are associated with decreases in consecutive wet days, maximum precipitation and zonal wind speeds, and increases in consecutive dry days. Parts of Kazakhstan, southern Russia, China, southeast Asia and northern India are assigned to LGM-C₅, which is characterised by increases in consecutive dry days. The region covered by PLIO-C₁ includes northern India along the Himalayan orogen (Fig. 5 c) and experiences decreases in consecutive dry days and increases in consecutive wet days and maximum precipitation (Fig. 5 f). 840 PLIO-C₂ covers most of the study region and is associated with relatively little change in all climatic attributes except meridional wind speeds. Central Asia is mostly assigned to PLIO-C₃ and experiences an increase in 2m air temperature and decrease in freeze-thaw days. The north of the study region (PLIO-C₄) is characterised by decreases in 2m air temperature amplitude and consecutive freezing days and increases in 2m air temperature and freeze-thaw days. Finally, PLIO-C₅ mostly covers the high altitude locations of the Himalaya-Tibet region that

Deleted: PLIO

are close to steep topographic gradients, including the Himalayan orogen. This cluster is associated with decreases in wind speeds, consecutive freezing days, consecutive wet days and 2m air temperature, and with increases in consecutive dry days and 2m air temperature.

Deleted: .

Deleted: The

855 *Discriminability*

The significance of the climate attributes in explaining the discriminability the Himalaya-Tibet clusters (Fig. 5 d,e,f) is more variable than in Europe. While changes in 2m air temperature are important in most of the MH and LGM results, there is no clear dominant factor for PLIO clusters. In the LGM, the discriminability in the high-altitude/high latitude cluster (LGM-C₁) is mostly explained by decreases in 2m air temperature (30%-40%), mild increases in consecutive freezing days (20-30%) and mild decreases in consecutive wet days (10%-20%). LGM-C₄ has the second highest discrimination score, and the discrimination in this cluster is explained by decreases in consecutive dry days (10%-20%) and increases in consecutive wet days (10%-20%), maximum precipitation (30%-40%) and 2m air temperature (20%-30%). For PLIO, the type of climate change governing the largest cluster (PLIO-C₂) causes discriminability that is primarily explained by mild decreases in consecutive freezing days (20%-30%) and mild increases in consecutive wet days (20%-30%) and 2m air temperature amplitude (10%-20%). Discriminability in the cluster occupying the region south of the Himalayan orogen (PLIO-C₁) is explained by decreases in consecutive dry days (10%-20%) and 2m air temperature (10%-20%), and increases in consecutive wet days (10%-20%), maximum precipitation (30%-40%) and consecutive freezing days (10%-20%). Cluster PLIO-C₅ is associated with a discriminability best explained by increases in consecutive dry days (30%-40%) and decreases in maximum precipitation (10%-20%) and consecutive freezing days (20%-30%).

860 **4. Discussion**

This section describes method-related features and problems, and highlights commonly occurring patterns of change, provides possible explanations for those, and discusses these changes in context of erosional processes.

875

The role of large scale features

For many of the LGM and some of the PLIO results, changes in 2m air temperature and/or consecutive freezing days significantly contribute to the discriminability in clusters covering mid-latitudes. LGM-C₁ – LGM-C₃ in coastal and high-altitude West Patagonia (South America), LGM-C₁ and PLIO-C₄ in the Himalaya-Tibet region 880 are examples of this. Many of these high-latitude clusters are also characterised by large changes in 2m air temperature and 2m air temperature amplitude in the LGM and PLIO results. The preferential cooling in higher latitudes during the LGM and enhanced meridional temperature gradient [e.g. Otto-Bliesner et al., 2006; Bracanot et al., 2007; Mutz et al., 2018] can be expected to result in more pronounced seasonality and thus higher variation in near surface temperature amplitude. Inversely, the accentuated warming in higher latitude during the 885 (Late) Pliocene [e.g. Salzman et al., 2011; Ballantyne et al., 2010; Mutz et al., 2018] would result in the opposite. These previously studied large-scale features explain much of the characterisation of high-latitude clusters and the significant contribution of changes in temperature-related variables to regional discriminability. Associated changes in temperature variables can have decisive impacts on physical weathering due to changes in glacial and periglacial processes (see below), as well as to biotic weathering by influencing vegetation cover.

Deleted:

Deleted: and

890

895 *The role of glaciers and periglacial processes*

Changes in temperature in high altitude regions can impact physical weathering through glacial erosion [e.g. Egholm et al., 2009, Herman et al. 2013] and periglacial processes [e.g. Andersen et al., 2015, Marshall et al., 2015]. In southernmost South America, northern Europe and South Alaska (cf. supplemental information), the high discriminability and modes of change on the multivariate anomaly maps for the LGM are primarily controlled by the glaciers covering most of the region. Furthermore, many modes of change in the study regions involve consecutive freezing days and freeze-thaw days. Changes from ice-free to ice-covered conditions, such as in LGM-C₁-LGM-C₃ in [coastal and high-altitude West Patagonia](#) (South America) and LGM-C₁-LGM-C₄ in [different terrains of](#) Europe, are associated with increases in consecutive freezing days and decreases in freeze-thaw days. The opposite is the case for some modes of changes in the PLIO. For example, PLIO-C₆ in [high-altitude Patagonia](#) (South America) is associated with changes from ice-covered to ice-free conditions, as well as with an increase in consecutive freezing days. It may therefore shift from glacier to frost-cracking dominated erosional processes. These modes of change in the PLIO mark possible transition from glacier governed processes to periglacial processes and thus increased frost-cracking as the Earth's surface spends more time in the frost-cracking window [e.g. Matsuoka, 2001; Schaller et al., 2002; Andersen et al., 2015, Marshall et al., 2015], whereas many LGM modes of change suggest the opposite. Finally, glacial pre-conditioning of a landscape can modulate the effect of precipitation on landsliding [Moon et al. 2011].

The role of precipitation characteristics

Areas that have been covered by glaciers during the LGM and experienced a post-LGM increase in maximum precipitation or consecutive wet days may be particularly prone to precipitation-triggered landslides. This is the case, for example, in the regions covered by LGM-C₂ and LGM-C₃ clusters in [high-altitude](#) Patagonia and LGM-C₁ – LGM-C₃ in [northern](#) Europe. More generally, changes in storminess affect erosion through river incision and sediment transport [e.g. Whipple and Tucker, 1999; Hobley et al., 2010]. Maximum precipitation and consecutive wet days are measures of storm intensity and duration respectively, which are primary controls for runoff and relevant for erosion. In most cases, such as the results for the Himalaya-Tibet region in the LGM and PLIO, the co-variability of consecutive wet days, maximum precipitation and consecutive dry days is intuitive: changes in consecutive wet days and maximum precipitation coincide with changes in consecutive dry days in the opposite direction. Even though palaeovegetation is considered in the set-up of the GCM simulations [Mutz et al., 2018], the modulating effect of vegetation on the impact of changes in the precipitation attribute variables on erosion [e.g. Gyssels et al., 2005] cannot be taken into account here, and thus the reader is advised to do so in their evaluation of the effect of these changes on Earth surface processes. [Note also that vegetation modifies river discharge by changing evapotranspiration and infiltration, and modifies hillslope erosion by changing root characteristics](#) [e.g. Schmid et al., 2018]. In absence of significant vegetation changes, areas such as Bhutan, Nepal, Bangladesh and parts of Northern India (MH-C₁ and PLIO-C₁), which experiences strong increases in consecutive wet days and maximum precipitation in the MH and PLIO, are likely to have experienced an increase in such precipitation-induced erosion at these times.

Deleted: i

The role of winds

935 Changes in wind speed components affect aeolian erosion, transport, and deposition, as well as mean raindrop trajectories, which should also be taken into consideration in the assessment of local precipitation-induced erosion [de Lima et al., 1992]. The results of this study reveal that changes in near surface meridional and zonal wind speeds contribute little to the discriminability between climates even in regions that experience wind direction changes due to a different ice cover in Europe [e.g. Siegert and Dowdeswell, 2004], which are reproduced well
940 by the model. Wind speeds only show a significant contribution to discriminability (20%-30%) in the subtropical latitudes of South America due to slower meridional winds in the LGM in the region. The distinctiveness in the character of atmospheric dust transport during the LGM [e.g. Andersen et al., 1998] and thus aeolian erosion may be attributed more to system response and changes in vegetation, which cannot be taken into account in this study, than to a distinctiveness of LGM wind speeds.

945

Comments on methodical implications

PLIO and LGM clusters are more stable than MH clusters on the multivariate anomaly maps. This stability can be attributed to the relatively large magnitude of climate change in PLIO and LGM time slices. Lower variance of MH anomalies make element attribution to anomaly clusters in the MH more sensitive to randomisation and
950 re-categorisation procedures (section 2.2). Consequently, the nature of MH patterns can be seen as the result of climate change of lower magnitude and less distinctiveness. The most important limitation is the poor representation of precipitation amount in areas of high topography and rainfall [Meehl et al., 2007 and comparisons with ERA-interim and station-based observations not presented here]. However, the threshold for what constitutes a “wet day” or “dry day” is relatively low [Zolina et al., 2010; Zhang et al., 2011; Zin and Jemain,
955 2010], so that the typical overestimation of total precipitation amount by ECHAM5 in such regions has little or no effect on the attribute variables *consecutive wet days* and *consecutive dry days*, particularly when analysed in comparison to another simulation at a different time (as was done here) which helps reduce any systematic model bias towards high precipitation rates. The overall performance of the palaeoclimate simulations is decent, as comparison with proxy-based reconstructions showed [Li et al. 2017; Mutz et al. 2018]. Erosion-relevant
960 processes that take place on small spatial or short temporal scales, such as intra-storm variations and rainfall characteristics [e.g. Ran et al., 2012], cannot be quantified in this study due to limited model resolution, output frequency and accuracy of such estimates on that scale. The consideration of non-climatic factors, such as local topography, slope and vegetation, is beyond the scope of this study, and the reader is advised to take these into consideration in their assessment of the effect of documented climate change on Earth surface processes. Lastly,
965 since the parameterisation in ECHAM5's land surface scheme creates deficiencies in the representation of the local hydrology and partitioning of precipitation [e.g. Weiland et al., 2011], variables such as runoff are excluded in this study. Routing GCM output through more sophisticated hydrological models or nesting regional climate models in the GCM's would allow more merited exploration of the regional hydrology.

Deleted:

Deleted: high

970 **5. Conclusions**

In this study, we quantified the differences between pre-industrial and Late Cenozoic palaeoclimates with regard to variables relevant to Earth surface processes, explained these quantified differences and identified dominant patterns and modes of palaeoclimate change. The key findings of this study are:

- LGM and PLIO climate change is more distinct and more easily quantified than climate change of the MH. This is reflected in the stability of geographical regions (clusters) showing the extent of regions governed by distinct modes of climate change.

980

- Changes in ice cover result in very distinct signatures of climate change. This is reflected by 1) the creation of clusters geographically associated with ice cover changes, 2) the persistence of these clusters when the assigned number of clusters (k) is varied in the procedure, and 3) ice cover changes in South America leading to the best discriminability overall.

Deleted: k

Formatted: Font: Italic

985

- In Europe, changes in 2m air temperature explain most of the discriminability between pre-industrial and all three palaeoclimates (MH, LGM and PLIO). Changes in consecutive freezing days and consecutive wet days are also significant contributors to climate discriminability in LGM and PLIO results respectively. Consequently, these factors lend the Late Cenozoic palaeoclimates their unique signature and should be central in assessments of changes in Earth surface processes.
- Increases in freeze-thaw days and temperature often coincide with decreases in consecutive freezing days, and vice versa. Regions governed by these modes of changes, such as western Patagonia during the LGM, are prone to changes in erosional process domain from peri-glacial to glacial or vice versa.

990

- Increases in consecutive wet days and maximum precipitation often coincide with decreases in consecutive dry days. Regions governed by these modes of change, such as locations south of the Himalayan orogen in the PLIO, can be expected to be particularly prone to changes in erosion induced by precipitation and storm characteristics.

Deleted: and

995

We note that the methods presented in this study may also be applied to simulations of modern and future climates. Furthermore, the procedure can easily be modified to detect and explain spatiotemporal changes in climate attributes associated with different processes, since the link between climate and the impacted system is established solely on the basis of variable selection. For example, the procedure may be used for the investigation of spatiotemporal changes of climate attributes that are related to specific human adaptation strategies in the past or future. Such method transfer is merited if (1) the underlying statistical assumptions are still satisfied and (2) the chosen variables adequately represent the relationship between climate and the investigated processes. The procedure may also be applied more broadly to any problem that requires the detection of regions (in either geographical or any n_h -dimensional variable space) associated with specific modes of change in data, and the subsequent quantification and explanation of changes occurring within identified regions.

Formatted: Subscript

1000

1005

1010

Data availability: All data files used in the presentation of results, as well as the model simulation results this study is based on [Mutz et al., 2018], are freely available through the data download section of our work group wiki (esdynamics.geo.uni-tuebingen.de/wiki/) as set of data products tailored to the needs of different geoscientific communities.

Competing interests: The authors declare that they have no conflict of interest.

1020

Acknowledgements: European Research Council (ERC) Consolidator Grant number 615703 provided support for this study. Additional support is acknowledged from the German science foundation (DFG) project number 365266215. [We also thank the reviewers \(including Andrew Wickert\) for their constructive comments and suggestions, which helped us improve this manuscript.](#)

1025

6. References

- Abe-Ouchi, A., Saito, F., Kageyama, M., Bracannot, P., Harrison, S. P., Lambeck, K., Otto-Btiesner, B. L., Peltier, W.R., Tarasov, L., Peterschmitt, J.-Y., Takahashi, K.: Ice-sheet configuration in the CMIP5/PMIP3 Last Glacial Maximum experiments. *Geosci. Model Dev.*, 8, 3621-3637, doi:10.5194/gmd-8-3621-2015, 2015
- 1030 Andersen, K. K., Armengaud, A., and Genthon, C.: Atmospheric dust under glacial and interglacial conditions, *Geophysical Research Letters*, 25(13), 2281-2284, 1998.
- Andersen, J., Egholm, D.L., Knudsen, M.F., Jansen, J., Nielsen, S.B., The periglacial engine of mountain erosion – Part 1: Rates of frost cracking and frost creep., *Earth Surface Dynamics*, 3(4), 447-462, 2015.
- 1035 Bahreberg, G., Giese, E., and Nipper, J.: Multivariate Statistik. *Statistische Methoden in der Geographie* 2, Stuttgart, 1992.
- Ballantyne, A.P., Greenwood, D.R., Sininghe Damste, J.S., Csank, A.Z., Eberle, J.J., and Rybczynski, N.: Significantly warmer Arctic surface temperatures during the Pliocene indicated by multiple independent proxies, *Geology* 38 (7), 603–606, 2010.
- 1040 Banfield, J.F., Barker, W.W., Welch, S.A., and Taunton, A.: Biological impact on mineral dissolution: application of the lichen model to understanding mineral weathering in the rhizosphere, *Proceedings of the National Academy of Sciences*, 96, 3404-3411, 1999.
- Bendick, R., and Ehlers, T.A.: Extreme localized exhumation at syntaxes initiated by subduction geometry, *Geophys. Res. Lett.*, 41(16), 2014GL061026, doi:10.1002/2014GL061026, 2014.
- 1045 Bertrand, S., Charlet F., Charlier B., Renson V., and Fagel N.: Climate variability of southern Chile since the Last Glacial Maximum: a continuous sedimentological records from Lago Puyehue (40°S), *J. Paleolimnol.*, 39, 179-195, doi:10.1007/s10933-007-9117-y, 2008.
- Bigelow, N. H., Brubaker, L. B., Edwards, M. E., Harrison, S. P., Prentice, I. C., Anderson, P. M., Andreev, A. A., Bartlein, P. J., Christensen, T. R., Cramer, W., Kaplan, J. O., Lozhkin, A. V., Matveyeva, N. V., Murray, D. V., McGuire, A. D., Razzhivin, V. Y., Ritchie, J. C., Smith, B., Walker, D. A., Gajewski, K., Wolf, V., Holmqvist, B. H., Igarashi, Y., Kremenetskii, K., Paus, A., Pisaric, M. F. J., and Vokova, V. S.: Climate change and Arctic ecosystems I. Vegetation changes north of 55°N between the last glacial maximum, mid-Holocene and present. *Journal of Geophysical Research - Atmospheres*, 108(D19), doi: 10.1029/2002JD002558, 2003.
- 1050 Bookhagen, B., Thiede, R.C., and Strecker, M.R.: Late Quaternary intensified monsoon phases control landscape evolution in the northwest Himalaya, *Geology*, 33(2), 149-152. doi: 10.1130/G20982.1, 2005.
- Braconnot, P., Otto-Btiesner, B., Harrison, S.P., Joussaume, S., Peterschmitt, J.-Y., Abe-Ouchi, A., Crucifix, M., Driesschaert, E., Fichefet, T., Hewitt, C.D., Kagayama, M., Kitoh, A., Loutre, M.-F., Marti, O., Merkel, U.,

- 1060 Ramstein, G., Valdes, P., Weber, L., Yu, Y., and Zhao, Y.: Results of PMIP2 coupled simulations of the mid-Holocene and Last Glacial maximum, part 1: experiments and large-scale features, *Clim Past* 3:261–277, 2007.¹¹¹
- Braconnot, P., Harrison, S. P., Kageyama, M., Bartlein, P. J., Masson-Delmotte, V., Abe-Ouchi, A., Otto-Bliesner, B., and Zhao, Y.: Evaluation of climate models using palaeoclimatic data, *Nature Climate Change*, 2, 417–424, 2012.
- 1065 CLIMAP Project Members: Seasonal Reconstruction of the Earth's Surface at the Last Glacial Maximum. *Map and Chart Series*, Vol. 36, Geological Society of America, 18 pp., 1981.
- Deal, E., Braun, J., and Botter, G.: Understanding the Role of Rainfall Hydrology in Determining Fluvial Erosion Efficiency, *JGR Earth Surface*, 123(4), 744–778, 2018. doi: 10.1002/2017JF004393
- Dowsett, H.J., Robinson, M., Haywood, A., Salzmann, U., Hill, D., Sohl, L., Chandler, M., Williams, M., Foley, K., and Stoll, D.: The PRISM3D paleoenvironmental reconstruction. *Stratigraphy*, 7, 123–139, 2010.
- 1070 Ehlers, T.A., and Poulsen, C.J.: Influence of Andean uplift on climate and paleoaltimetry estimates, *Earth and Planetary Science Letters*, 281(3–4), 238–248, 2009.
- Gong, X., Knorr, G., Lohmann, G., and Zhang, X.: Dependence of abrupt Atlantic meridional ocean circulation changes on climate background states, *Geophysical Research Letters*, 40 (14), 3698–3704, doi:10.1002/grl.50701, 2013.
- 1075 Gyssels, G., Poesen, J., Bochet, E., and Li, Y.: Impact of plant roots on the resistance of soils to erosion by water: a review, *Progress in Physical Geography*, 29, 189–217. <http://doi.org/10.1191/030913305pp443ra>, 2005.
- Harrison, S. P., Yu, G., Takahara, H., and Prentice, I. C.: Palaeovegetation - Diversity of temperate plants in east Asia. *Nature* 413, 129–130, 2001.
- 1080 Haywood, A.M., Dowsett, H.J., Otto-Bliesner, B., Chandler, M.A., Dolan, A.M., Hill, D.J., Lunt, D.J., Robinson, M.M., Rosenbloom, N., Salzmann, U., and Sohl, L.E.: Pliocene Model Intercomparison Project (PlioMIP): experimental design and boundary conditions (Experiment 1), *Geoscientific Model Development* (3), 227–242, 2010.
- Haywood, A.M., Hill, D.J., Dolan, A.M., Otto-Bliesner, B., Bragg, F., Chan, W.-L., Chandler, M.A., Contoux, C., Jost, A., Kamae, Y., Lohmann, G., Lunt, D.J., Abe-Ouchi, A., Pickering, S.J., Ramstein, G., Rosenbloom, N.A., Sohl, L., Stepanek, C., Yan, Q., Ueda, H., and Zhang, Z.: Large-scale features of Pliocene climate: results from the Pliocene Model Intercomparison Project, *Clim. Past*, 9, 191–209. doi:10.5194/cp-9-191-2013, 2013.
- 1085 Haywood, A. M., Dowsett, H. J., Dolan, A. M., Rowley, D., Abe-Ouchi, A., Otto-Bliesner, B., Chandler, M. A., Hunter, S.J., Lunt, D. J., Pound, M., and Salzmann, U.: The Pliocene Model Intercomparison Project (PlioMIP) Phase 2: scientific objectives and experimental design, *Clim. Past*, 12, 663–675, <https://doi.org/10.5194/cp-12-663-2016>, 2016.
- Herman, F., Seward, D., Valla, P.G., Carter, A., Kohn, B., Willet, S. D., Ehlers, T. A.: Worldwide acceleration of mountain erosion under a cooling climate. *Nature*, 504, 423–426, 2013.
- 1090 Hobley, D.E., Sinclair, H.D., and Cowie, P.A.: Processes, rates, and time scales of fluvial response in an ancient postglacial landscape of the northwest Indian Himalaya. *Geological Society of America Bulletin*, 122, 1569–1584, 2010.

- 1100 Insel, N., Ehlers, T.A., Schaller, M., Barnes, J.B., Tawacoli, S., and Poulsen, C.J.: Spatial and Temporal Variability in Denudation across the Bolivian Andes from Multiple Geochronometers, *Geomorphology*, 122, 65-77, doi.org/10.1016/j.geomorph.2010.05.014, 2010.
- Jeffery, M.L., Ehlers, T.A., Yanites, B.J., and Poulsen, C.J.: Quantifying the role of paleoclimate and Andean Plateau uplift on river incision: PALEOCLIMATE ROLE IN RIVER INCISION, *Journal of Geophysical Research: Earth Surface*, 118(2), 852–871, doi:10.1002/jgrf.20055, 2013.
- Jungclaus, J. H., Lorenz, S. J., Timmreck, C., Reick, C. H., Brovkin, V., Six, K., Segschneider, J., Giorgetta, M.A., Crowley, T.J., Pongratz, J., Krivova, N.A., Vieira, L.E., Solanski, S.K., Klocke, D., Botzet, M., Esch, M., Gayler, V., Haak, H., Raddatz, T.J., Roeckner, E., Schnur, R., Widmann, H., Claussen, M., Stevens, B., and Marotzke, J.: Climate and carbon-cycle variability over the last millennium. *Climate of the Past*, 6, 723–737. doi:10.5194/cp-6-723-2010, 2010.
- Kageyama, M., Albani, S., Braconnot, P., Harrison, S. P., Hopcroft, P. O., Ivanovic, R. F., Lambert, F., Marti, O., Peltier, W. R., Peterschmitt, J.-Y., Roche, D. M., Tarasov, L., Zhang, X., Brady, E. C., Haywood, A. M., LeGrande, A. N., Lunt, D. J., Mahowald, N. M., Mikolajewicz, U., Nisancioglu, K. H., Otto-Bliesner, B. L., Renssen, H., Tomas, R. A., Zhang, Q., Abe-Ouchi, A., Bartlein, P. J., Cao, J., Li, Q., Lohmann, G., Ohgaito, R., Shi, X., Volodin, E., Yoshida, K., Zhang, X., and Zheng, W.: The PMIP4 contribution to CMIP6 – Part 4: Scientific objectives and experimental design of the PMIP4-CMIP6 Last Glacial Maximum experiments and PMIP4 sensitivity experiments, *Geosci. Model Dev.*, 10, 4035–4055, https://doi.org/10.5194/gmd-10-4035-2017, 2017.
- Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclaus, J., Otto-Bliesner, B. L., Peterschmitt, J.-Y., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Brierley, C., Crucifix, M., Dolan, A., Fernandez-Donado, L., Fischer, H., Hopcroft, P. O., Ivanovic, R. F., Lambert, F., Lunt, D. J., Mahowald, N. M., Peltier, W. R., Phipps, S. J., Roche, D. M., Schmidt, G. A., Tarasov, L., Valdes, P. J., Zhang, Q., and Zhou, T.: PMIP4-CMIP6: the contribution of the Paleoclimate Modelling Inter-comparison Project to CMIP6, *Geosci. Model Dev.*, https://doi.org/10.5194/gmd-11-1033-2018, 2018.
- Knorr, G., Butzin, M., Micheels, A., and Lohmann, G.: A Warm Miocene Climate at Low Atmospheric CO₂ levels. *Geophysical Research Letters*, 38, L20701, doi:10.1029/2011GL048873, 2011.
- Kutzbach, J.E., Prell, W.L., and Ruddiman, W.F.: Sensitivity of Eurasian Climate to Surface Uplift of the Tibetan Plateau, *Journal of Geology*, 101(2), 177-190, 1993.
- Li, J., Ehlers, T.A., Werner, M., Mutz, S.G., Steger, C., Paeth, H.: Late quaternary climate, precipitation δ¹⁸O, and Indian monsoon variations over the Tibetan Plateau. *Earth and Planetary Science Letters* 457, 412-422, 2017.
- Lohmann, G., Pfeiffer, M., Laepple, T., Leduc, G., and Kim, J.-H.: A model-data comparison of the Holocene global sea surface temperature evolution. *Clim. Past*, 9, 1807-1839, doi:10.5194/cp-9-1807-2013, 2013.
- Maroon, E. A., Frierson, D. M. W., and Battisti, D. S.: The tropical precipitation response to Andes topography and ocean heat fluxes in an aquaplanet model, *J. Climate*, 28, 381–398, doi:10.1175/JCLI-D-14-00188.1, 2015.
- Maroon, E. A., Frierson, D. M. W., Kang, S. M., and Scheff, J.: The precipitation response to an idealized subtropical continent, *J. Climate*, 29, 4543–4564, doi:10.1175/JCLI-D-15-0616.1, 2016.

- Marshall, J.A., Roering, J.J., Bartlein, P.J., Gavin, D.G., Granger, D.E., Rempel, A.W., Praskievicz, S.J., and Hales, T.C.: Frost for the trees: Did climate increase erosion in unglaciated landscapes during the late Pleistocene? *Science Advances*, 1, 1-10, 2015.
- 1140 Matsuoka, N.: Solifluction rates, processes and landforms: A global review. *Earth Science Reviews* 55, 107–134, 2001.
- Meehl, G.A.; Stocker, T. F.; Collins, W. D.; Friedlingstein, P.; et al. In: Solomon, S.; Qin, D.; Manning, M.; Chen, Z.; Marquis, M.; Averyt, K. B.; Tignor, M.; Miller, H. L. editors, editor/s. IPCC, 2007: Climate Change 2007: the physical science basis. contribution of Working Group I to the Fourth Assessment Report of the 1145 Intergovernmental Panel on Climate Change. Cambridge, U.K.: Cambridge University Press; 2007.
- Montgomery, D.R.: Slope distributions, threshold hillslopes, and steady-state topography, *American Journal of Science*, 301, 432-454, 2001.
- Moon, S., Chamberlain, C.P., Blisniuk, K., Levine, N., Rood, D.H., Hilley, G.E.: Climatic control of denudation in the deglaciated landscape of the Washington Cascades, *Nature Geoscience*, 4, 469-473, 2011.
- 1150 Mutz, S.G., Ehlers, T.A., Li, J., Steger, C., Peath, H., Werner, M., Poulsen, C.J.: Precipitation $\delta^{18}\text{O}$ over the South Asia Orogen from ECHAM5-wiso Simulation: Statistical Analysis of Temperature, Topography and Precipitation. *Journal of Geophysical Research-Atmospheres*, 121(16), 9278-9300, doi: 10.1002/2016JD024856, 2016.
- Mutz S.G., Ehlers T., Werner M., Lohmann G., Stepanek C., Li J.: Estimates of Late Cenozoic climate change 1155 relevant to Earth surface processes in tectonically active orogens. *Earth Surface Dynamics*. doi.org/10.5194/esurf-6-271-2018.
- Otto-Bliesner, B.L., Brady, C.B., Clauzet, G., Tomas, R., Levis, S., and Kothavala, Z.: Last Glacial Maximum and Holocene Climate in CCSM3. *Journal of Climate*, 19, 2526-2544, 2006.
- Otto-Bliesner, B. L., Braconnot, P., Harrison, S. P., Lunt, D. J., Abe-Ouchi, A., Albani, S., Bartlein, P. J., 1160 Capron, E., Carlson, A. E., Dutton, A., Fischer, H., Goelzer, H., Govin, A., Haywood, A., Joos, F., LeGrande, A. N., Lipscomb, W. H., Lohmann, G., Mahowald, N., Nehrbass-Ahles, C., Pausata, F. S. R., Peterschmitt, J.-Y., Phipps, S. J., Renssen, H., and Zhang, Q.: The PMIP4 contribution to CMIP6 – Part 2: Two interglacials, scientific objective and experimental design for Holocene and Last Interglacial simulations, *Geosci. Model Dev.*, 10, 3979–4003, <https://doi.org/10.5194/gmd-10-3979-2017>, 2017.
- 1165 Paeth, H.: Key Factors in African Climate Change Evaluated by a Regional Climate Model, *Erdkunde*, 58, 290-315, 2004.
- Pfeiffer, M., and Lohmann, G.: Greenland Ice Sheet influence on Last Interglacial climate: global sensitivity studies performed with an atmosphere-ocean general circulation model. *Climate of the Past*, 12, pp. 1313-1338, doi:10.5194/cp-12-1313-2016, 2016.
- 1170 Pickett, E.J., Harrison, S.P., Flenley, J., Grindrod, J., Haberle, S., Hassell, C., Kenyon, C., MacPhail, M., Martin, H., Martin, A.H., McKenzie, M., Newsome, J.C., Penny, D., Powell, J., Raine, J.I., Southern, W., Stevenson, J., Sutra, J.-P., Thomas, I., van der Kaars, S., Ward, J.: Pollen-based reconstructions of biome distributions for Australia, South-East Asia and the Pacific (SEAPAC region) at 0, 6000 and 18,000 14C years B.P.. *Journal of Biogeography*, 31, 1381–1444, doi: 10.1111/j.1365-2699.2004.01001.x, 2004.
- 1175 Prentice, I. C., Jolly, D., and BIOME 6000 Participants. (2000). Mid-Holocene and glacial-maximum vegetation geography of the northern continents and Africa. *Journal of Biogeography* 27, 507-519.

- Ran, Q., Su, D., Li, P., and He, Z.: Experimental study of the impact of rainfall characteristics on runoff generation and soil erosion. *Journal of Hydrology*, 424-425, 99-111, 2012.
- Salzmann, U., Williams, M., Haywood, A.M., Johnson, A.L.A., Kender, S., and Zalasiewicz, J.: Climate and environment of a Pliocene warm world. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 309 (1-8), 2011.
- Sarnthein, M., Gersonde, R., Niebler, S., Pflaumann, U., Spielhagen, R., Thiede, J., Wefer, G., Weinelt, M.: Overview of Glacial Atlantic Ocean Mapping (GLAMAP 2000). *Paleoceanography*, 18(2), doi:10.1029/2002PA000769, 2003.
- 1185 Schaller, M., von Blanckenburg, F., Veldkamp, A., Tebbens, L.A., Hovius, N., and Kubik, P.W.: A 30 000 yr record of erosion rates from cosmogenic ^{10}Be in Middle European river terraces, *Earth and Planetary Science Letters*, 204(1), 307-320, 2002.
- [Schmid, M., Ehlers, T.A., Werner, C., Hickler, T., Fuentes-Espoz, J.-P.: Effects of changing vegetation and precipitation on denudation – Part 2: Predicted landscape response to transient climate and vegetation cover over millennial to million-year timescales, *Earth Surface Dynamics*, 6, 859-881, tps://doi.org/10.5194/esurf-6-859-2018, 2018.](#)
- 190 Sohl, L.E., Chandler, M.A., Schmunk, R.B., Mankoff, K., Jonas, J.A., Foley, K.M., and Dowsett, H.J.: PRISM3/GISS topographic reconstruction, *U.S. Geological Survey Data Series*, 419, 6p., 2009.
- Siegert, M. J., and Dowdeswell, J. A.: Numerical reconstruction of the Eurasian Ice Sheet and climate during the 1195 Late Weichselian. *Quaternary Science Reviews*, 23, 1273-1283, 2004.
- Starke, J., Ehlers, T. A., and Schaller, M.: Tectonic and Climatic Controls on the Spatial Distribution of Denudation Rates in Northern Chile (18°S - 23°S) Determined From Cosmogenic Nuclides. *Journal of Geophysical Research: Earth Surface*, 122, 1949-1971. doi:10.1002/2016JF004153, 2017.
- Stepanek, C., and Lohmann, G.: Modelling mid-Pliocene climate with COSMOS , *Geosci. Model Dev.* , 5 , pp. 1200 1221-1243 . doi:10.5194/gmd-5-1221-2012, 2012.
- Stock, G.M., Frankel K.L., Ehlers, T.A., Schaller, M., Briggs, S.M., Finkel R.C.: Spatial and temporal variations in denudation of the Wasatch Mountains, Utah, USA, *Lithosphere GSA*, v. 1, pp. 34-40. doi.org/10.1130/L15.1, 2009.
- Takahashi, K., and Battisti, D.: Processes controlling the mean tropical pacific precipitation pattern. Part I: The 1205 Andes and the eastern Pacific ITCZ. *Journal of Climate*, 20(14), 3434-3451, 2007.
- Whipple, K.X., and Tucker, G.E.: Dynamics of the stream-power river incision model: Implications for height limits of mountain ranges, landscape response timescales, and research needs, *Journal of Geophysical Research-Solid Earth*, 104, 17661-17674, 1999.
- Whipple, K. X.: The influence of climate on the tectonic evolution of mountain belts, *Nat. Geosci.*, 2, 1210 doi:10.1038/ngeo413, 2009.
- Wei, W., and Lohmann, G.: Simulated Atlantic Multidecadal Oscillation during the Holocene. *Journal of Climate*, 25, 6989–7002. doi: <http://dx.doi.org/10.1175/JCLI-D-11-00667.1>, 2012.
- Weiland, F.C.S., van Beek, L.P.H., Kwadjik, J.C.J., and Bierkens, M.F.P.: [On the Suitability of GCM Runoff Fields for River Discharge Modelling: A Case Study Using Model Output from HadGEM2 and ECHAM5](#). *Journal of Hydrometeorology*, 13, 140-154, 2011.

Formatted: English (US)

Formatted: Font: Italic

Formatted: English (US)

Formatted: English (UK)

Formatted: Font: Italic

Wickert, A.D: Reconstruction of North American drainage basins and river discharge since the Last Glacial Maximum. *Earth Surface Dynamics*, 4, 831-869. doi:10.5194/esurf-4-831-2016, 2016.

Formatted: Font: Italic

Wilks, D.S.: Statistical methods in the atmospheric sciences - 3rd ed. Academic Press, Oxford, 2011.

Willett, S.D., Schluengger, F., and Picotti, V.: Messinian climate change and erosional destruction of the central
1220 European Alps. *Geology*, 34(8), 613-616, 2006.

Zhang, X., Lohmann, G., Knorr, G., and Purcell C.: Abrupt glacial climate shifts controlled by ice sheet changes.
Nature, 512 (7514), 290-294, doi:10.1038/nature13592, 2014.

Zhang, Q., Singh, V. P., Li, J., and Chen, X.: Analysis of periods of maximum consecutive wet days in China.
Journal of Geophysical Research, 116, D23106, <http://dx.doi.org/10.1029/2011JD016088>, 2011.

1225 Zin, W. Z. W., and Jemain A. A.: Statistical distributions of extreme dry spell in Peninsular Malaysia. *Theor Appl
Climatol*, 102, 253-264, 2010.

Zolina, O., Simmer, C., Gulev, S.K., and Kollet, S.: Changing structure of European precipitation: Longer wet
periods leading to more abundant rainfall. *Geophysical Research Letters*, 37, L06704,
doi:10.1029/2010GL042468, 2010.

1230

Figure Captions

1235 Fig. 1 Anomalies are created from pre-industrial (PI) and Late Cenozoic palaeoclimates (MH, LGM, PLIO). These
are subjected to geographical clustering, which results in the identification of distinct modes of climate change
(anomaly cluster characterisation) and maps showing the spatial extent of regions governed by these modes
(anomaly cluster maps). These are used as geographical masks for palaeoclimate simulations. For each of these
anomaly clusters, a discriminant analysis is conducted to quantify the discriminability in each cluster (score) and
1240 the relative contribution of each climatic variable to this discriminability.

Fig. 2 Example problem: Investigating how climatic boundary conditions for erosion processes have changed in
a specific region of interest involves consultation of part of the conceptualised results figure. The location of the
region of interest geographically coincides with the region assigned to cluster C1. Consequently, only the results
1245 related to C1 are consulted and the rest is greyed out. The clusters were calculated from the differences in the
values (delta values) of geomorphic variables between two different times (T1 and T2) in the late Cenozoic, and
thus represent a specific mode of change. The mode of change associated with the cluster of interest (C1) is
revealed by the purple-green column. The relative contributions of the delta values to the overall discriminability
1250 between T1 and T2 in cluster C1, indicated by the score, is revealed by the diameters of the circle superimposed
onto the delta values in the purple-green column.

Fig. 3 The multivariate anomaly maps for time slice comparisons PI-MH(a), PI-LGM(b) and PI-PLIO(c) show
the geographical coverage of clusters C₁-C_i in Western South America, which describe the spatial extend of
regions characterised by similar modes of change. The corresponding modes of change (d,e and f) for each cluster
1255 are expressed as relative changes in each of the 9 investigated variables (table 1): 2m air temperature (te2m), 2m
air temperature amplitude (t2am), consecutive freezing days (csfd), freeze-thaw days (fthd), maximum
precipitation (pmax), consecutive wet days (cswd), consecutive dry days (csdd), zonal near surface wind speeds
(u10) and meridional near surface wind speeds (v10). The score (d,e and f) expresses the goodness of
discriminability between the palaeoclimate pairs PI-MH(d), PI-LGM(e) and PI-PLIO(f) in each of the anomaly
1260 clusters. The size of the circles corresponds to the relative contribution of each of the 9 climatic attribute variables
to the measured discriminability in each anomaly cluster for all three time slice comparisons.

Fig. 4 The multivariate anomaly maps for time slice comparisons PI-MH(a), PI-LGM(b) and PI-PLIO(c) show
the geographical coverage of clusters C₁-C_i in Europe, which describe the spatial extent of regions characterised
1265 by similar modes of change. The corresponding modes of change (d,e and f) for each cluster are expressed as
relative changes in each of the 9 investigated variables (table 1): 2m air temperature (te2m), 2m air temperature
amplitude (t2am), consecutive freezing days (csfd), freeze-thaw days (fthd), maximum precipitation (pmax),
consecutive wet days (cswd), consecutive dry days (csdd), zonal near surface wind speeds (u10) and meridional
near surface wind speeds (v10). The score (d,e and f) expresses the goodness of discriminability between the
1270 palaeoclimate pairs PI-MH(d), PI-LGM(e) and PI-PLIO(f) in each of the anomaly clusters. The size of the circles

corresponds to the relative contribution of each of the 9 climatic attribute variables to the measured discriminability in each anomaly cluster for all three time slice comparisons.

1285

Fig. 5 The multivariate anomaly maps for time slice comparisons PI-MH(a), PI-LGM(b) and PI-PLIO(c) show the geographical coverage of clusters C₁-C_i in Himalaya-Tibet, which describe the spatial extend of regions characterised by similar mod of change. The corresponding modes of change (d,e and f) for each cluster are expressed as relative changes in each of the 9 investigated variables (table 1): 2m air temperature (te2m), 2m air 1290 temperature amplitude (t2am), consecutive freezing days (csfd), freeze-thaw days (fthd), maximum precipitation (pmax), consecutive wet days (cswd), consecutive dry days (csdd), zonal near surface wind speeds (u10) and meridional near surface wind speeds (v10). The score (d,e and f) expresses the goodness of discriminability 1295 between the palaeoclimate pairs PI-MH(d), PI-LGM(e) and PI-PLIO(f) in each of the anomaly clusters. The size of the circles corresponds to the relative contribution of each of the 9 climatic attribute variables to the measured discriminability in each anomaly cluster for all three time slice comparisons.

Deleted: -

Deleted: -

Deleted: -

Deleted: -

Deleted: -

Deleted: -

Table 1: Code, units, explanation and geomorphic relevance of each of the climate attribute variables used in the cluster- and discriminant analysis.

1300