



The destiny of orogen-parallel streams in the Eastern Alps: the Salzach-Enns drainage system

Georg Trost¹, Jörg Robl², Stefan Hergarten³, Franz Neubauer²

¹Department of Geoinformatics, Paris-Lodron University Salzburg, 5020, Austria

5 ²Department of Geography and Geology, Paris-Lodron University Salzburg, 5020, Austria

³Institute of Earth and Environmental Sciences, Albert-Ludwigs University of Freiburg, 79104, Germany

Correspondence to: Georg Trost (georg.trost@sbg.ac.at)

Abstract. The evolution of the drainage system in the Eastern Alps is inherently linked to different tectonic stages of the alpine orogeny. Crustal scale faults imposed east-directed orogen parallel flow on major rivers, whereas late orogenic surface uplift
10 increased topographic gradients between foreland and range and hence the vulnerability of such rivers to be captured. This leads to a situation where major orogen-parallel alpine rivers such as the Salzach River or the Enns River are characterized by elongated east-west oriented catchments south of the proposed capture points, whereby almost the entire drainage area is located west of the capture point. To determine the current stability of drainage divides and to predict the potential direction of divide migration, we analyzed their geometry at catchment, headwater and hillslope scale. Therefore, we employ χ mapping
15 for different base levels, generalized swath profiles along drainage divides and Gilbert metrics. Our results show that almost all drainage divides are asymmetric with steeper channels west and flatter channels east of a common drainage divide. Interpreting these results, we propose that drainage divides migrate from west towards east, so that the Inn catchment grows on expense of the Salzach catchment and the Salzach catchment consumes the westernmost tributaries of the Mur and Enns catchments. While Gilbert metrics show the same trend at hillslope scale at the Salzach–Enns and Salzach–Mur drainage
20 divide, they show no significant asymmetry at the Inn–Salzach drainage divide. As topography at the latter divide is dominated by glacial landforms such as cirques and U-shaped valleys, we interpret the missing hillslope scale asymmetry of this divide as a result of Pleistocene climate modulations, which locally obscured the large-scale signal of drainage network reorganization. We suggest that the east-directed divide migration progressively leads to symmetric catchment geometries, where eventually tributaries west and east of the capture point contribute equally to the drainage area. To test this assumption
25 we have reconstructed the proposed drainage network for different time slices. χ mapping of these reconstructed drainage networks indicates a progressive stability of the network topology in the Eastern Alps towards the present-day situation.



1 Introduction

- 30 The tectonic and climatic evolution of a collisional orogen is inherently linked to its drainage system (Beaumont et al., 1992; Bishop, 2007; Cederbom et al., 2004; Champagnac et al., 2012; Garcia-Castellanos et al., 2003; Herman et al., 2013; Montgomery et al., 2001; Robl et al., 2017a; Roe et al., 2008; Willett, 1999). In a zone of plate convergence, crustal shortening is a primary control of the horizontal and vertical metrics of the mountain range (Bartosch et al., 2017; Houseman and England, 1986; Robl et al., 2017a; Robl et al., 2008b; Robl and Stüwe, 2005b; Royden et al., 1997). Progressive shortening leads to
- 35 thickening of light, buoyant crust, which results in surface uplift and formation of high alpine topography (e.g. Molnar and Lyon-Caen, 1988). The horizontal geometry of the mountain range reflects compression in, and stretching perpendicular to the direction of plate convergence. In such a stress field, blocks of the brittle upper crust are abundantly advected along major strike slip fault zones – a process, which is commonly referred to as lateral extrusion (e.g. Ratschbacher et al., 1991; Ratschbacher et al., 1989; Robl et al., 2008b; Robl and Stüwe, 2005b; Robl and Stüwe, 2005a; Tapponnier et al., 1982).
- 40 As a consequence of the horizontal and vertical velocity field of the crust, drainage systems are also advected (Castelltort et al., 2012; Clark et al., 2004; Goren et al., 2015; Fox et al., 2014; Guerit et al., 2018; Kirby and Whipple, 2012; Miller et al., 2012; Miller and Slingerland, 2006; Stüwe et al., 2008; Yang et al., 2016). However, rivers are not just passive markers of crustal deformation, but also adjust their channel slopes to the contributing drainage area, uplift rate and bedrock properties until longitudinal channel profiles are graded and long-term erosion rates are in balance with uplift rates (Goren et al., 2014;
- 45 Kooi and Beaumont, 1996; Robl et al., 2017b; Whipple, 2001; Willett et al., 2001). Over an orogenic cycle, tectonic forcing and climatic conditioning are not steady. The signal of temporal variations is routed via mobile knickpoints in channels through the entire drainage system (Kirby and Whipple, 2012; Perron and Royden, 2013; Royden and Perron, 2013; Robl et al., 2017b; Wobus et al., 2006). Evidence for a previous tectonic phase is erased from their shapes once all knickpoints have left the drainage system at the drainage divides.
- 50 If a drainage divide becomes thereby asymmetric, it features an across divide gradient in erosion rate and starts migrating from the steep towards the less steep side (e.g. Forte and Whipple, 2018; Gilbert, 1877; Robl et al., 2017a; Robl et al., 2017b; Whipple et al., 2017). The reorganization of the drainage system due to divide migration (continuous) and river piracy events (discrete) lasts at least one order of magnitude longer than the upstream migration of knickpoints in channels (e.g. Goren, 2016; Robl et al., 2017b). Furthermore, changes in the contributing drainage area as consequence of mobile divides result in a
- 55 positive feedback, where the adaption of channel profiles to changing catchment size amplifies across divide differences in erosion rate (Willett et al., 2014). As a consequence, information on major tectonic phases associated with a large scale reorganization of the drainage pattern persists in the drainage network topology and can be revealed from analyzing the geometric properties of the drainage system and its divides, even if direct evidence from channels profiles has already vanished



(Beeson et al., 2017; Goren et al., 2015; Hergarten et al., 2016; Robl et al., 2017a; Robl et al., 2017b; Willett et al., 2014;
60 Winterberg and Willett, 2019; Yang et al., 2015).

In this study, we aim to decipher the morphological state of drainage divides in the Eastern Alps to (a) distinguish between
mobile and immobile drainage divides and (b) constrain the potential direction of divide migration by applying a set of
morphometric tools that consider divide disequilibrium at catchment, headwater and hillslope scale. Furthermore, we discuss
our results in the light of proposed changes in the drainage pattern since onset of topography formation in the Eastern Alps
65 and explore how these changes may have affected the stability of the divides compared to the present-day situation

1.1 The drainage system of the Eastern Alps

The drainage system of the Eastern Alps is characterized by two principal drainage divides (Robl et al., 2008a; Robl et al.,
2017a) (Fig. 1). One major divide follows the main ridge of the Eastern Alps including the highest peaks and separates the
Inn, Salzach and Enns catchments to the north from the Drau and Mur catchments to the south. The Danube (and eventually
70 the Black Sea) represents the common base level of all those rivers, but their confluence is located in the Pannonian Basin
hundreds of kilometers apart from the Eastern Alps. A second major drainage divide separates Alpine rivers that flow into the
Adriatic Sea (e.g. Adige River) from the Mur–Drau drainage system.

Fault tectonics controlled the configuration of the drainage systems. Major tectonic lineaments (mainly strike-slip dominated
faults, i.e. Inn Valley Fault, Salzach-Ennstal-Mariazell-Puchberg Fault (SEMP), Mur-Mürz Fault, Periadriatic Lineament (PL),
75 Möll Valley Fault) confine a corridor of lateral extrusion, where crustal blocks were actively squeezed out towards the
Pannonian Basin (e.g. Ratschbacher et al., 1989; Ratschbacher et al., 1991). Almost all major streams of the Eastern Alps
follow these major tectonic lineaments for several tens of kilometers (Bartosch et al., 2017; Robl et al., 2008a; Robl et al.,
2017a). Hence, they flow parallel to the strike of the mountain range, instead of leaving the orogen towards north and south,
following the general topographic gradient (Fig. 1).

80 The courses of the Salzach and Enns rivers in the north, and the Mur and Mürz rivers in the south are characterized by knee-
shaped bends and T-shaped river junctions, where rivers abruptly leave their tectonically preconditioned valleys and drain
towards the forelands in the north and south, respectively (Robl et al., 2008a; Robl et al., 2017a). Such sudden river course
changes in concert with the observation of wind gaps at the Salzach–Enns valleys and the Salzach–Saalach drainage divide,
as well as the provenance of sediments along the Enns valley (Dunkl et al., 2005; Neubauer, 2016) are consistent with the
85 proposed reorganization of the Salzach and Enns drainage systems (Dunkl et al., 2005; Kuhlemann et al., 2001; Robl et al.,
2008a).

South of the alpine main ridge, major valleys show also a strong tectonic control (Robl et al., 2017a). The eastern tributaries
of the Adige River and the western tributaries of the Drau River follow roughly the Periadriatic Lineament (Fig. 1). The
occurrence of a prominent wind gap between Adige and Drau rivers, as well as T-shaped river junctions at the tributaries of
90 the Adige River is discussed in terms of river piracy events and an ongoing reorganization of the drainage system (Robl et al.,
2017a).



1.2 Co-evolution of topography and drainage system of the Eastern Alps

Morphological observations (e.g. Robl et al., 2008a) and provenance analyses (e.g. Kuhlemann et al., 2001; Kuhlemann et al., 2002; Kuhlemann, 2007; Neubauer, 2016) give evidence for several large scale modifications of the Eastern Alpine drainage system. The evolution of the drainage system is inherently linked to the Late Oligocene–Early Miocene indentation (Handy et al., 2015) of the Adriatic into the European plate. At onset of indentation, the landscape of the Eastern Alps was characterized by a hilly topography (Frisch et al., 2001), which was drained by a series of northward flowing rivers (Frisch et al., 1998; Kuhlemann et al., 2006; Kuhlemann, 2007).

During Early to Middle Miocene times lateral extrusion tectonics, confined by a set of crustal scale strike-slip and associated normal faults, caused rapid exhumation of the Tauern Window. This process initiated a large scale reorganization of the drainage system, where faults imposed an east-directed orogen-parallel flow on major rivers (Frisch et al., 1998). This tectonic stage set the paleo-courses of the Enns, Mur and Drau Rivers (Dunkl et al., 2005; Kuhlemann et al., 2006; Kuhlemann, 2007). Evidence for a changing drainage pattern was recorded by the sedimentary pile deposited in the northern Molasse basin (Kuhlemann et al., 2006) and in inner alpine basins (Dunkl et al., 2005). The sedimentary record consists of characteristic rocks of surrounding Austroalpine units and later, during Middle and Late Miocene, of rocks from Penninic and Subpenninic units of the rising Tauern Window (Frisch et al., 1998).

Provenance analyses of sediments reveal the reversal of flow directions and potential stream capture events during the Late Miocene (Kuhlemann, 2007). Frisch et al. (1998) and Dunkl et al. (2005) suggested that the initially northeast directed Mur River changed its course to the current southeast directed drainage path during the Middle Miocene (Fig. 1). The detection of wind gaps (Robl et al., 2008a) and the analysis of the sedimentary composition of intra-orogenic basins (Neubauer, 2016), suggest similar changes in the Salzach and Enns drainage systems. The abrupt increase in stream power, a few kilometers upstream, but mostly downstream of the knee-shaped river bend, and a knickpoint analysis of tributaries may indicate a base level lowering of the Salzach River in the recent geological past (Robl et al., 2008a).

However, the drainage development since the early Pliocene is poorly constrained. In particular, the impact of the Pleistocene glaciations, resulting in flat valley floors of the trunk streams and hanging valleys with large knickpoints at tributaries (Robl et al., 2008a; Norton et al., 2010; Valla et al., 2010) altered the geometry of rivers and obscured the tectonic record of preceding tectonic events (Robl et al., 2017a).

2 Method

All topographic analyses are based on the EU-DEM (data funded under GMES, Global Monitoring for Environment and Security preparatory action 2009 on Reference Data Access by the European Commission) digital elevation model with a spatial resolution of approx. 25 m.



2.1 χ -mapping

In order to detect potentially mobile drainage divides due to across divide differences in erosion rate, we follow the approach of Willett et al. (2014) by employing the so called χ transform (Perron and Royden, 2013; Royden and Perron, 2013). This approach is based on the detachment-limited model for bedrock channel incision (Hergarten, 2002; Howard, 1980; Howard, 1994) of the form

$$\frac{\partial H}{\partial t} = U - KA^m \left(\frac{\partial H}{\partial x} \right)^n \quad (1)$$

where H , t and x are elevation, time and the longitudinal coordinate along the river profile, increasing in the upstream direction. U and K represent the uplift rate and the erodibility of the bedrock. The nonlinear contribution of the channel slope $\frac{\partial H}{\partial x}$ and drainage area A to river incision is represented by the exponents m and n .

In topographic steady state ($\frac{\partial H}{\partial t} = 0$), Eq. (1) simplifies to

$$\frac{\partial H}{\partial x} = \left(\frac{U}{K} \right)^{\frac{1}{n}} A^{-\frac{m}{n}} \quad (2)$$

If U and K are constant, we get

$$\frac{\partial H}{\partial x} \propto A^{-\theta} \quad (3)$$

where $\theta = m/n$ is the concavity index. Then, the product

$$k_s = \frac{\partial H}{\partial x} A^\theta \quad (4)$$

is constant and known as the steepness index k_s (e.g. Snyder et al., 2000; Wobus et al., 2006), which directly represents the ratio between U and K for $n = 1$.

The non-linear increase of contributing drainage area (and hence discharge) with downstream distance yields the curvature of the channel profile, which obscures the record of spatial or temporal changes in uplift rate or contributing drainage area in the geometry of the river channel. The χ transform eliminates the curvature of the river profile by transforming the longitudinal



coordinate x to a new coordinate χ (Perron and Royden, 2013; Royden and Perron, 2013). The contributing drainage area can be eliminated if the transformation satisfies the condition

$$\frac{dx}{d\chi} = \left(\frac{A}{A_0}\right)^\theta \quad (5)$$

155

which is achieved by

$$\chi = \int_{x_0}^x \left(\frac{A}{A_0}\right)^{-\theta} dx \quad (6)$$

160 where x_0 is an arbitrary reference point, and A_0 is an arbitrary reference catchment size ($A_0 = 1 \text{ m}^2$).

Under spatially and temporally uniform tectonic and climatic conditions, χ transformed river profiles are straight lines and their slope is constant and proportional to k_s . We calculated χ -values for all channels with a contributing drainage area $> 1 \text{ km}^2$ and $\theta = 0.5$, using the numerical approach of (Hergarten et al., 2016).

165 The analysis of across divide differences in χ exploits the fact that channels, originating at a common drainage divide (i.e. similar channel head elevation) and sharing the same base level elevation, are steep, if χ is small (Willett et al., 2014). Hence, across divide differences in χ indicate differently steep rivers on both sides of the divide, averaged from the baselevel to the channel head. Generalizing the ideas of Gilbert (1877), steeper channels result in higher erosion rates, and hence, drainage divides should migrate towards the high χ catchments.

2.2 Generalized swath profiles

170 We employ generalized swath profiles (Hergarten et al., 2014) to explore differences in headwater relief across drainage divides. The drainage divide represents the curved baseline of the swath profile. The signed minimum distance (Euclidian distance) of every data point of the digital elevation model to the base line is computed and coordinate pairs (profile coordinate, distance) are binned. Topographic maxima and minima representing the summit domain and the drainage system, respectively, as well as mean elevation and standard deviation indicating the degree of landscape dissection are represented as function of signed distance from the drainage divide. The half width of the swath profiles is 5 km.

2.3 Gilbert metrics

180 To investigate the symmetry of drainage divides and potential anomalies at the hillslope scale, we determine the so called Gilbert metrics, originally proposed by Gilbert (1877) and formalized by Whipple et al. (2017). Across divide differences in channel head elevation, hillslope gradient and local relief were computed with Divide Tools (Forte and Whipple, 2018), a collection of morphometric functions based upon TopoToolbox (Schwanghart and Scherler, 2014).



Channel heads at the transition from the hillslope to the fluvial domain are defined by a contributing drainage area threshold of 1 km². Hence, channel head elevation is the elevation at this point. The local relief is the maximum elevation (E_{\max}) within a circular window minus the elevation of the channel head. We chose the default window size with a radius of 0.5 km (Forte and Whipple, 2018), which encloses the nearby ridge lines, but does not reach far beyond. The slope gradient is the average topographic gradient between the channel head and the highest point within the analyzed window. These metrics are averaged (arithmetic averaging) at each side of the watershed. Δ -values (e.g. $\Delta_{\text{Elevation}}$) represent the difference of averaged metrics of the two sides of the drainage divide. Eventually, Δ -values are normalized to a range from -1 to 1, so that every deviation from 0 evidences for an asymmetric drainage divide. Following the nomenclature of Forte and Whipple (2018), we refer to these metrics as Gilbert metrics.

190 3 Results

By applying a set of standard morphometric analyses, we discovered several distinctly asymmetric drainage divides. We found divide asymmetry considering information from entire catchments, headwaters, and even hillslopes.

3.1 χ -mapping: across divide differences at catchment scale

As already described by Robl et al. (2017a) and Winterberg and Willett (2019), we found distinct χ anomalies at the divides of the Salzach catchment and the Inn and Adige (WS 1) catchments in the west, the Saalach (WS 2) catchment in the north, and the Enns (WS 3) and Mur (WS 4) catchments in the east (Figs. 1, 2a-d). Across divide differences in χ between the Salzach catchment and the Drau catchment in the south are small. As a clear trend, all streams at the western side of roughly north-south trending drainage divides feature significantly lower χ values than adjacent streams east of the divides. We observe this trend at WS 1, where tributaries of the eastern Salzach River show significantly higher χ values than tributaries of the western Inn River. Similar anomalies in χ occur at WS 3 and WS 4, where the tributaries of the Enns and Mur Rivers feature higher χ values than tributaries of the Salzach River. At WS 2, separating the Salzach from the Saalach catchment, higher χ values are observed north of the divide within the Saalach catchment.

A stepwise increase of the baselevel from 400 m (Fig. 2a) to 600 m (Fig. 2b) and 800 m (Fig. 2c), and tantamount a shift of the starting point of the χ computation towards the headwaters, changes the absolute χ values, but does not change the observed across divide gradients in χ . However, starting the χ integration at the very headwaters of the investigated catchments by setting a baselevel of 1000 m (Fig. 2d), several across divide χ gradients disappear or are even reverted, as observed at WS 1. There, and in contrast to lower base levels, tributaries of the Inn River feature higher χ values than tributaries of the Salzach River. The rivers on both sides of WS 2 and WS 3 show similar χ values. However, the distinct χ anomaly observed at WS 4 still remains. All tributaries of the Mur show higher χ values than tributaries across the divides to the Drau, Salzach and Enns



210 catchments. Beyond that, the analysis shows that χ gradients across the Mur and Enns drainage divides increase with increasing
baselevels.

For a baselevel of 400 m, absolute values of χ , extracted at the channel heads on both sides of the investigated drainage divide,
reflect the described across divide χ gradients quantitatively (Fig. 3). At the westernmost drainage divide of the Salzach
catchment, the distribution of χ ranges between 3190 m and 6740 m in the Inn / Adige catchment and between 5810 m and
215 8890 m in the Salzach catchment. Mean values in χ are 4887 m at the Inn / Adige side and 7525 m at the Salzach side of the
drainage divide. The χ gradient indicates that the average steepness of the channels is higher at the Inn / Adige side than at the
Salach side of the divide. At the eastern drainage divides of the Salzach catchment, the Salzach – Enns and the Salzach – Mur
divide, the χ distribution of the Salzach ranges between 2670 m and 6530 m, while channel heads at the Enns and Mur
catchment feature χ values between 4410 m and 9100 m. Mean values in χ are 4267 m and 5443 m at the Salzach catchment,
220 and 6360 m and 8093 m at the Enns and Mur catchments. The χ gradients indicate higher average channel steepness at the
Salach side of the divides. Across divide gradients at the northern Salzach – Saalach divide are distinctly smaller than those
at the western and eastern Salzach watersheds. The χ distribution ranges between 3670 m and 6220 m at the Salzach side and
between 3390 m and 8130 m at the Saalach side. Mean χ is slightly shifted towards higher values at the Saalach (5834 m)
225 catchment. relative to the Salzach side (4829 m). This, however, is caused by the long tail of the skewed right χ distribution of the Saalach
catchment.

3.2 Swath profiles: across divide differences at headwater scale

The four curved swath profiles following the watershed segments WS1 – WS4 indicate a series of distinct across divide
differences in the headwater relief (Figs. 1, 4). At first glance, WS 1 appears roughly symmetric with a steady decrease in
mean (E_{mean}) and minimum elevation (E_{min}) with increasing distance from the divide. Up to a distance of 2 km, the drop in
230 E_{mean} is larger at the Inn side of the divide. At a distance of 5 km, E_{min} is slightly lower at the Salzach side in comparison to the
Inn side. At this distance, the swath corridor already reached the trunk valley of the Salzach drainage system, but reached only
a small tributary of the Inn River at the other side of the divide. Overall, the relief ($E_{\text{max}} - E_{\text{min}}$) is larger at the Inn than at the
Salach side of the divide. The Saalach – Salzach drainage divide (Fig. 4, WS2) shows a strong asymmetry in E_{mean} and relief,
but no spatial trend in E_{min} . The latter is bound up with the fact that the drainage divide exhibits a wind gap, which connects
235 the valley floors of the Salzach and Saalach rivers without a significant drop in valley floor elevation (Fig. 1). In contrast, the
eastern divides of the Salzach catchment show a strong asymmetry (Fig. 4, WS3, 4). The drop in E_{min} and E_{mean} with increasing
distance from the divides is distinctly more pronounced at the Salzach side than at the Enns and Mur sides of the drainage
divide. In consequence, high gradients in E_{min} and E_{mean} form towards west, and gentle gradients arise towards east.



3.3 Gilbert Metrics: across divide differences at hillslopes scale

240 The Gilbert metrics suggested by Forte and Whipple (2018) comprise three measures characterizing the local differences at drainage divides (i.e. channel head elevation, mean upstream relief, mean upstream gradient). Overall, a strong divide asymmetry at hillslope scale is only observed at the Salzach – Mur drainage divide (Fig. 5).

At the westernmost drainage divide of the Salzach catchment, elevations at channel heads (Fig 5, WS 1) lie in the range between 1100 m and 2600 m. At the Salzach basin, channel head elevations show an unimodal distribution with a mean value
245 of 2044 m, while the distribution at the Inn / Adige basin is bimodal and has a mean value of 1977 m. Overall differences are small, but indicate a slight shift towards lower channel head elevations at the Inn / Adige side of the drainage divide. The upstream relief ranges between 200 m and 660 m and is uniformly distributed within the Salzach, but skewed-left distributed in the Adige catchment. Mean values of upstream relief are similar in the Salzach and Inn / Adige catchment with 386 m and
250 382 m, respectively. Analogous to the upstream relief, upstream gradient is uniformly and skewed-left distributed in the Salzach and Inn / Adige catchments, respectively. Values for upstream gradient are in the range of 0.2 and 0.8, with mean values of 0.45 for the Salzach and 0.44 for the Inn / Adige catchments. Beside outliers, upstream relief and upstream gradient appears slightly larger in the Inn / Adige catchment than in the Salzach catchment.

At the Salzach–Saalach drainage divide, differences in all Gilbert metrics are small. Elevation at channel heads (Fig 5, WS 2) ranges between 740 m to 1750 m with mean values of 1424 m and 1370 m at the Salzach and Saalach side of the divide. The
255 upstream relief and the upstream gradient range between 250 m and 760 m, and between 0.35 and 1.1 in the Salzach and Saalach catchment, respectively.

While the eastern drainage divide, separating the Salzach from the Enns and the Mur catchments, features consistently large anomalies in χ , Gilbert metrics representing the hillslope scale indicate a largely symmetric Salzach – Enns, and a distinctly asymmetric Salzach – Mur drainage divide. Channel head elevation of the Salzach – Enns divide (Fig. 5, WS 3) ranges between
260 840 m and 2200 m, with mean values of 1206 m and 1296 m at the Salzach and Enns side of the divide. The lower channel head elevation is also reflected by a slightly higher mean upstream relief and mean upstream gradient in the Salzach, in comparison to the Enns catchment. Mean values are 351 m and 0.42 for the Salzach, and 343 m and 0.4 for the Enns catchment. The divide between Salzach and Mur catchments is characterized by the largest across divide differences in all Gilbert metrics (Fig. 5, WS 4). Elevation at channel head lies between 1350 m and 2220 m. On average, channel head elevation is distinctly
265 lower in the Salzach catchment (1747 m) than in the Mur catchment (1982 m). Lower channel head elevations result in a larger mean upstream relief and higher upstream gradient for the Salzach (425 m, 0.5), in comparison to the Mur catchment (375 m, 0.42).

4 Discussion

Gilbert (1877) already recognized that cross-divide differences in erosion rate result in mobile watersheds, whereby catchments
270 featuring higher erosion rates grow on expense of adjacent catchments with lower erosion rates. Given uniform climate and



lithology, erosion rate increases with channel steepness in the drainage and topographic gradient in the hillslope domain. Hence, divide asymmetry evidences for drainage divide migration from the steep towards the less steep side (Forte and Whipple, 2018; Gilbert, 1877; Robl et al., 2017b; Willett et al., 2014). Asymmetry at drainage divides may occur at catchment, headwater and hillslope scale, but may not necessarily be observed at all these magnitudes. For example, an increase or
275 decrease in drainage area due to a river capture event may cause a χ anomaly at the drainage divide, which predicts drainage divide mobility. However, if the signal has not yet reached the divide, the divide may still be symmetric at the hillslope scale indicating divide stability at that time. Glacially controlled base level lowering with an increase in local relief at the north facing side of divides may cause a strong asymmetry at hillslope scale, but will not result in an anomaly in χ maps, as long as the drainage network topology remains unchanged.

280 4.1 Challenges and limitations interpreting drainage divide asymmetries

A direct determination of divide migration rates is unfeasible, as migration rates are in the range of millimeters per year (Goren et al., 2014) and major river capture events are rarely observed (Brocard et al., 2012; Yanites et al., 2013). In concert with sediment provenance (Frisch et al., 1998; Kuhlemann, 2007) and erosion rates based on cosmogenic nuclides (e.g. Dixon et al., 2016), topographic metrics serve as proxy for drainage divide mobility. In particular, in mid-latitude mountain ranges such
285 as the Eastern Alps, gradients in erosion rate reflect rather a transient landscape state due to glacial–interglacial periods, than across divide differences resulting from the reorganization of the drainage system. However, the interpretation of topographic metrics derived from tectonically active, mid-latitude mountain ranges such as the Eastern Alps is not unique and paved with some pitfalls. The topography of the Eastern Alps resulted from different tectonic phases and hence spatial and temporal changes in the uplift pattern (Bartosch et al., 2017; Frisch et al., 1998; Robl et al., 2008a). Furthermore, hillslopes are
290 abundantly at a critical slope angle affected by numerous landslides (Kühni and Pfiffner, 2001) and show in some regions a strong glacial imprint (e.g. Dixon et al., 2016), whereby the turnover time from glacial to fluvial landscape characteristics is controlled by lithology (Robl et al., 2015) and uplift rate (Prasicek et al., 2015).

Across divide differences in χ may indicate spatial or temporal variations of climate (e.g. precipitation) or tectonics (e.g. uplift rate) (e.g. Whipple et al., 2013), but may also reflect the long term reorganization of the drainage network accompanied by
295 changes in contributing drainage area (Goren et al., 2014; Robl et al., 2017b; Willett et al., 2014). Contrasts in bedrock erodibility may also result in differently steep channels despite uniform erosion rates. Even if channel steepness and erosion rate differ across drainage divides, divides may still be stable if differently steep channels result from spatial variations in uplift rate. Alternative formulations of the χ transform, considering spatial variation in uplift rate (Willett et al., 2014) and erosion rate (Robl et al., 2017a) have been presented, but information on the long term uplift / erosion pattern is sparse in the Eastern
300 Alps. The curvature of channel profiles removed by the χ transform is introduced by the non-linear increase of discharge in downstream direction. However, instead of discharge (or peak discharge) contributing drainage area as proxy for discharge in rivers is employed by the detachment limited version of the stream power equation, which in its original form neglects spatial



climatic variations, such as orographic precipitation. Yang et al. (2016) considered precipitation computing χ ; but again long term precipitation rates are not well constrained and the control of precipitation on the geometry of river channels is still debated (e.g. Burbank et al., 2003; Dadson et al., 2003; Wobus et al., 2003; Hodges et al., 2004; Molnar, 2003; Reiners et al., 2003). Beyond methodical issues, the vertical distance between channel head and base level divided by χ is the average steepness of the channel, but provides no explicit information on the steepness of the dividing ridge itself. Consequently, a low increase in χ at the lower channel reach may result in a steep channel on average, and small χ values even at the channel heads. This, however, means that χ anomalies at drainage divides may indicate potential divide mobility in the future, rather than currently mobile divides. Without doubt, many factors and processes may lead to an amplification or emergence of across divide gradients in χ and complicate the interpretation of χ in terms of divide stability (Whipple et al., 2017; Forte and Whipple, 2018). As a strategy to counteract some of these pitfalls, a series of χ -maps with progressively raised base levels narrows down the impact of spatial heterogeneity in tectonics and climate from catchment to headwater scale. This allows statements on the position of the disturbance within the drainage system and potential divide mobility in the far and in the near future.

The Gilbert metrics, a set of local topographic measures characterize hillslopes at both sides of the investigated divide (Forte and Whipple, 2018) and hence the (a)symmetry of the divide itself. In contrast to χ mapping, there are no far field effects and significant asymmetry of the dividing ridge should correspond in principle to across divide gradients in erosion rate and divide mobility. However, in active, mid-latitude mountain ranges, several factors and processes make the interpretation of these metrics challenging. Landslide-controlled threshold hillslopes emerge, where incision rates in the drainage system are high (Montgomery et al., 2001). Then the relationship between topographic gradient and hillslope erosion rate breaks, and dividing ridges become symmetric although they feature across divide gradients in erosion rate and migrate. For the European Alps, an average limiting slope stability angle of 25° is reported (Schmidt and Montgomery, 1995; Kühni and Pfiffner, 2001), so that most of the divides in the study area are prone to landsliding. However, in particular within the formerly glaciated realm of the Alps, many of the non-soil mantled hillslopes are distinctly steeper and still feature glacial landscape characteristics (Robl et al., 2015). There, local metrics such as relief, gradient or channel head elevation rather indicate the impact of the last glaciations on topography than long-term trends in drainage network reorganization. Glacial overprint does not primarily affect the first order drainage networks, but has a strong impact on local relief (e.g. Brocklehurst and Whipple, 2002; Norton et al., 2010; Salcher et al., 2014; van der Beek and Bourbon, 2008). Aspect-controlled differences in relief formation due to glacial erosion (e.g. north versus south facing mountain flanks) result in local, reversible compensating motions of the divides (Robl et al., 2017a) that may counteract the regional trend during the turn overtime from glacial to fluvial landscapes. Hence, such local disturbances cover large scale and long lasting changes in the drainage network topology. Generalized swath profiles and χ maps with a base level at the headwaters may bridge the gap between catchment and hillslope scale and assist detecting local peculiarities as described above.

Summarizing, the major advantage of Gilbert metrics lies in the analysis of short wavelength – high amplitude signals, e.g. the development of escarpments (e.g. Tucker and Slingerland, 1994). In contrast, the reorganization of drainage patterns forced



by tectonic processes represents a high length scale – low amplitude signal taking place in millions of years (Robl et al., 2015), which can be targeted best by the calculation of χ maps. Headwater processes and the position of the erosional signal can be addressed by varying the base level for the χ transformation and the extraction of generalized swath profiles. We hereinafter discuss the mobility of the drainage divides in consideration of the described pitfalls.

340 4.2 Mobility of Drainage Divides in the Eastern Alps

Our results imply that drainage divides of the investigated catchments are mobile and follow a general trend: at north – south running drainage divides, tributaries feature lower χ values west of the dividing ridge and hence are steeper on average than tributaries draining towards east (Fig. 2). Given uniform tectonic, climatic and lithological conditions, steeper channels are characterized by higher erosion rates (e.g. Howard, 1994), so that the observed morphological divide asymmetry at the Salzach – Enns, Salzach – Mur and Inn – Salzach watersheds may result in erosion rate gradients across the divides. However, uniform tectonic and climatic conditions and substrate properties on the scale of major drainage systems cannot be assumed in a tectonically active mountain range, such as the Eastern Alps. This means that the observed χ pattern may not necessarily indicate migrating divides, but may result from spatial variations on climate (e.g. precipitation, Yang et al., 2016) and tectonics (e.g. uplift rate, Willett et al., 2014). However, progressively increasing the base level for χ computation and shifting the observational scale from presumable tectonically and climatically heterogeneous catchments to their more homogenous headwaters shows no qualitative changes in the χ pattern. The χ anomalies across the divides remain up to a baselevel of 800 m (Fig. 2, 3) and indicate that divides are mobile. However, this trend in χ breaks down at some divides for a base level of 1000 m characterizing the very headwaters, only.

Gilbert metrics characterizing divides at hillslope scale are consistent with the χ pattern at the Salzach – Enns and Salzach – Mur drainage divide and indicate that these divides are currently mobile. However, and in contrast to the χ pattern (up to a base level of 800 m), they indicate divide stability at the Inn – Salzach drainage divide (Figs. 2, 3). In particular at the latter divide, glacial landforms such as cirques and U-shaped valleys are abundant and we interpret the missing hillslope scale asymmetry of this divide as a result of glacial erosion, which temporally stops divide migration. However, Robl et al. (2017a) showed that the impact of variable glacial erosion across divides is small and reversible. We suggest that the topographic signal of cold climate processes, primarily acting during the Pleistocene, locally obscures the large-scale signal of drainage network reorganizations in many parts of the Eastern Alps and in general limits the applicability of Gilbert metrics in glacially shaped mountain ranges.

The peculiar west – east directed migration of drainage divides is inherently linked to the plan view geometry of the Salzach and Enns catchments south of the Northern Calcareous Alps (Figs. 1, 2, 6). In this domain, the main stem of the Salzach and Enns still follows the SEMP, which is one the major tectonic lineaments of the Eastern Alps (Wang and Neubauer, 1998). It has been proposed that during the Mid-Miocene, Salzach and Enns formed a common catchment with an east-directed flow path (e.g. Neubauer, 2016), but were separated by major river piracy events due headward eroding south – north draining rivers



(Kuhlemann et al., 2001; Dunkl et al., 2005; Robl et al., 2008a). As a consequence, the major portion of the Salzach and Enns drainage areas are located west of their capture points and by reversing the flow direction only to a minor amount east of the capture points. This is consistent with the current asymmetry of the catchments, with a large western and a small eastern sub-catchment and explains the observed across divide gradients in χ . Long east-directed channel segments in concert with distinctly elongated catchments result in a slow decrease in catchment size in upstream direction. Hence, χ , which is the measure for upstream flow length after channel length profile linearization (c transform, Perron and Royden, 2013; Royden and Perron, 2013), accumulates to large χ values at the western and low χ values at the eastern drainage divides. Integrating from a common base level up to the same channel head elevation, large and small χ values on different sides of a common divide are the expression of a low and high average channel steepness of long west–east and short east–west draining channel segments, respectively. This, however, implies that observed χ anomalies at the investigated drainage divides are the consequence of the Early to Mid-Miocene lateral extrusion tectonics (Ratschbacher et al., 1989; Ratschbacher et al., 1991), where the activity of crustal scale faults imposed non-ideal flow-direction to major rivers. The indicated drainage network reorganization from orogen-parallel to orogen-perpendicular flow is a long-lasting process. While river piracy events cause a sudden large-scale modification of the drainage network, drainage divide migration and flow direction reversal is a slow continuous process at rates of few millimeters per year (Goren et al., 2014), which explains the longevity of morphological disequilibrium after changes in the tectonic forcing.

4.3 Stability of divides for different evolutionary states

Based on provenance analyses and geomorphological studies, it has been proposed that different tectonic phases have triggered a repeated reorganization of the drainage system since the onset of topography formation in the Eastern Alps (Frisch et al., 2000; Kuhlemann et al., 2001; Dunkl et al., 2005; Kuhlemann, 2007; Keil and Neubauer, 2009; Neubauer, 2016). As the position of past drainage divides is not well constrained, we test if and how different catchment geometries, roughly mimicking the catchment geometry suggested for different phases of the drainage evolution, affect the stability of drainage divides (Fig. 7). We focus on the plan view geometry of catchments only and do not consider potential topographic (e.g. uplift of the Northern Calcareous Alps) or base level changes (e.g. inversion of the northern foreland basin, the Molasse basin). For the period of lateral extrusion during the Early to Mid-Miocene, it was suggested that the Salzach and Enns formed a common drainage system (Paleo-Enns) with an orogen-parallel flow path following the SEMP fault from west to east (Fig. 7a) (Frisch et al., 1998). Compared to the present-day drainage pattern (Fig. 7d), the elongated catchment with its long main stem and numerous short tributaries contributing drainage area from south and north cause very high χ values at the eastern domain of the drainage system. In particular, the Inn – Paleo-Enns drainage divide is characterized by high across divide gradients in χ , but even the southern drainage divide between Paleo-Enns and Drau indicates a strong χ anomaly. This suggests that the Early to Mid-Miocene situation, with elongated catchments featuring hundreds of kilometers of orogen-parallel flow, were prone to river piracy events and the migration of drainage divides.



400 The timing of the following drainage network reorganization is not well constrained. However, streams originating south of
the SEMP fault and draining towards the northern foreland basin eroded headwards and captured the eastward draining Paleo-
Enns (Salzach – Enns) drainage system (Frisch et al., 1998). Currently, two rivers, the Salzach and the Enns, follow the SEMP
for more than 100 km each, but abruptly change their course in a knee-shaped bend towards north. These sudden changes in
flow direction most likely indicate capture points. In addition, a suspicious wind gap separating the Saalach from the Salzach
405 valley with a vertical drop of only a few meters may indicate that the Paleo-Enns was once captured by the Saalach River, but
redirected again potentially during the Pleistocene glaciations (Robl et al., 2008a) (Fig. 7b, c).

Although not yet constrained by provenance studies, a potential capture of the Paleo-Enns drainage system by the Saalach
River would stabilize the westernmost drainage divide (Inn – Saalach divide). This is indicated by a decrease of the across
divide χ gradient similar to the present situation (Fig. 7d). However, in this scenario the western drainage divide of the Enns
410 catchment shows a distinct across divide χ gradient with high χ values at the Paleo-Enns and low χ values at the Saalach side
of the divide, which would result in a progressive flow direction reversal of the upper Enns River.

It is still debated when the orogen-parallel Paleo-Enns River was captured by the south–north draining Salzach River (Fig. 7c).
A Pleistocene, glacial-induced activation of the northward directed drainage outflow and blocking the passage at the Saalach
– Salzach windgap is discussed (Robl et al., 2008a). However, a similar event recently happened in Yukon, Canada due to
415 climate warming and glacial retreat (Headley, 2017; Shugar et al., 2017). Assuming a Pleistocene capture event due to waxing
or waning of glaciers, the origin of the Salzach (similar to the nearby Lammer River) must have inevitable been located south
of the NCA. Beyond others, one argument for such a scenario is the small amount of flow reversal of the former Enns River
east of the capture point. However, the concurrent drainage through Saalach and Salzach valley would lead to the disappearance
of the χ gradient across the Saalach and Salzach watershed (Fig. 7c). The watershed separating Salzach from Enns basin
420 complies with the present-day location of the watershed (Fig. 7c, d) and the χ distribution shows a similar χ anomaly indicating
an eastward drainage divide migration as proposed for the present-day situation.

5 Conclusion

The tectonic evolution of the Eastern Alps caused a repeated reorganization of the drainage network since onset of topography
formation in the Late Oligocene / Early Miocene. We applied various morphometric methods to constrain the potential mobility
425 of drainage divides on catchment, headwater and hillslope scale. Based on our analysis, we came to the following conclusions.

- Almost all drainage divides of the investigated domain are asymmetric at catchment, headwater, and even hillslope
scale, which evidences for drainage divide mobility, where the steeper side of the divide migrates towards the less
steep side of the divide.
- It turned out that the western side of the considered drainage divides is in general steeper than the eastern side, so
430 that the general direction of divide migration is west towards east. This implies that the Inn catchment grows on
expense of the Salzach catchment and the Salzach catchment consumes tributaries of the Enns and Mur catchments.



- At some divides, metrics characterizing hillslopes (Gilbert metrics) are not consistent with those characterizing larger scales. We found that glacial imprint locally obscures large-scale signals of drainage network reorganization. Hence, the applicability of the classical Gilbert metrics in mid-latitude mountain ranges such as the Eastern Alps is limited.
435
- The reason for the general drainage migration trend from west towards east is caused by the geometry of catchments, which dates back to the period of lateral extrusion in the Early to Mid-Miocene. The activity of major faults north and south of the central axis of eastern Alps imposed non-ideal, orogen-parallel flow directions to major rivers. Subsequent capture events restored orogen-perpendicular flow, but relics of the lateral extrusion period remained: elongated catchments west of the capture point, with an about 100 km long east-draining main stem and short tributaries draining south–north and north–south.
440
- A rough restoration of the drainage networks from Early to Mid-Miocene towards the present situation shows that anomalies in χ at the divides decreased, indicating that divide stability increased over time. Currently, large across divide gradients in average channel steepness (and hence erosion rate) occur mostly at north – south running watersheds (i.e. Inn – Salzach, Salzach – Enns, Salzach – Mur), where tributaries with short and long flow lengths from channel heads to the base level meet at the divides. However, as divide migration is slow and major capture events at these divides are not expected, we suggest that the observed disequilibrium is long-lasting.
445

Timing of river piracy events and rates of drainage divide migration are still not well constrained. There is great need for additional provenance studies of river sediments, and dating river terraces and cave sediments.
450

Data availability

The EU-DEM (data funded under GMES, Global Monitoring for Environment and Security preparatory action 2009 on Reference Data Access by the European Commission) is available from the European Environment Agency.

455 Author contribution

GT and JR conceived the study. SH and JR developed the algorithms for χ mapping and the calculation of generalized swath profiles. GT performed the geomorphological analyses. FN contributed to the understanding of the tectonic evolution of the study area. All authors contributed to the final form of the manuscript.



Competing interests

460 The authors declare that they have no conflict of interest.

Acknowledgements

This research was funded by the Austrian Science Fund (FWF) through the Doctoral College GIScience at the University of Salzburg (DK W 1237-N23).

465

References

- Bartosch, T., Stüwe, K., and Robl, J.: Topographic evolution of the Eastern Alps: The influence of strike-slip faulting activity, *Lithosphere*, 9, 384-398, 10.1130/1594.1, 2017.
- 470 Beaumont, C., Fullsack, P., and Hamilton, J.: *Erosional control of active compressional orogens*, Thrust Tectonics, edited by: McClay, K. R., 1-18 pp., 1992.
- Beeson, H. W., McCoy, S. W., and Keen-Zebert, A.: Geometric disequilibrium of river basins produces long-lived transient landscapes, *Earth and Planetary Science Letters*, 475, 34-43, 10.1016/j.epsl.2017.07.010, 2017.
- Bishop, P.: Long-term landscape evolution: linking tectonics and surface processes, *Earth Surface Processes and Landforms*, 32, 329-365, 10.1002/esp.1493, 2007.
- 475 Brocard, G., Willenbring, J., Suski, B., Audra, P., Authemayou, C., Cosenza-Murales, B., Moran-Ical, S., Demory, F., Rochette, P., Vennemann, T., Holliger, K., and Teyssier, C.: Rate and processes of river network rearrangement during incipient faulting: The case of the Cahabon River, Guatemala, *American Journal of Science*, 312, 449-507, 10.2475/05.2012.01, 2012.
- Brocklehurst, S. H., and Whipple, K. X.: Glacial erosion and relief production in the Eastern Sierra Nevada, California, *Geomorphology*, 42, 1-24, 10.1016/S0169-555X(01)00069-1, 2002.
- 480 Burbank, D. W., Blythe, A. E., Putkonen, J., Pratt-Sitaula, B., Gabet, E., Oskin, M., Barros, A., and Ojha, T. P.: Decoupling of erosion and precipitation in the Himalayas, *Nature*, 426, 652-655, 10.1038/nature02187, 2003.
- Castelltort, S., Goren, L., Willett, S. D., Champagnac, J. D., Herman, F., and Braun, J.: River drainage patterns in the New Zealand Alps primarily controlled by plate tectonic strain, *Nature Geoscience*, 5, 744-748, 10.1038/ngeo1582, 2012.
- 485 Cederbom, C. E., Sinclair, H. D., Schlunegger, F., and Rahn, M. K.: Climate-induced rebound and exhumation of the European Alps, *Geology*, 32, 709-712, 10.1130/g20491.1, 2004.
- Champagnac, J. D., Molnar, P., Sue, C., and Herman, F.: Tectonics, climate, and mountain topography, *Journal of Geophysical Research-Solid Earth*, 117, 10.1029/2011jb008348, 2012.
- Clark, M. K., Schoenbohm, L. M., Royden, L. H., Whipple, K. X., Burchfiel, B. C., Zhang, X., Tang, W., Wang, E., and Chen, L.: Surface uplift, tectonics, and erosion of eastern Tibet from large-scale drainage patterns, *Tectonics*, 23, 1-21, 10.1029/2002TC001402, 2004.
- 490 Dadson, S. J., Hovius, N., Chen, H., Dade, W. B., Hsieh, M.-L., Willett, S. D., Hu, J.-C., Hornig, M.-J., Chen, M.-C., Stark, C. P., Lague, D., and Lin, J.-C.: Links between erosion, runoff variability and seismicity in the Taiwan orogen, *Nature*, 426, 648-651, 10.1038/nature02150, 2003.
- Dixon, J. L., von Blanckenburg, F., Stüwe, K., and Christl, M.: Glaciation's topographic control on Holocene erosion at the eastern edge of the Alps, *Earth Surface Dynamics*, 4, 895-909, 10.5194/esurf-4-895-2016, 2016.
- 495 Dunkl, I., Kuhlemann, J., Reinecker, J., and Frisch, W.: Cenozoic relief evolution of the Eastern Alps—constraints from apatite fission track age-provenance of Neogene intramontane sediments, *Mitteilungen der österreichischen geologischen Gesellschaft*, 98, 92-105, 2005.
- Forté, A. M., and Whipple, K. X.: Criteria and tools for determining drainage divide stability, *Earth and Planetary Science Letters*, 493, 102-117, 10.1016/j.epsl.2018.04.026, 2018.
- 500 Fox, M., Goren, L., May, D. A., and Willett, S. D.: Inversion of fluvial channels for paleorock uplift rates in Taiwan, *J Geophys Res-Earth*, 119, 1853-1875, 10.1002/2014jf003196, 2014.
- Frisch, W., Kuhlemann, J., Dunkl, I., and Brugel, A.: Palinspastic reconstruction and topographic evolution of the eastern Alps during late Tertiary tectonic extrusion, *Tectonophysics*, 297, 1-15, 10.1016/S0040-1951(98)00160-7, 1998.



- Frisch, W., Dunkl, I., and Kuhlemann, J.: Post-collisional orogen-parallel large-scale extension in the Eastern Alps, *Tectonophysics*, 327, 239-265, 10.1016/s0040-1951(00)00204-3, 2000.
- 505 Frisch, W., Kuhlemann, J., Dunkl, I., and Szekely, B.: The Dachstein paleosurface and the Augenstein Formation in the Northern Calcareous Alps - a mosaic stone in the geomorphological evolution of the Eastern Alps, *International Journal of Earth Sciences*, 90, 500-518, 10.1007/s005310000189, 2001.
- Garcia-Castellanos, D., Verges, J., Gaspar-Escribano, J., and Cloetingh, S.: Interplay between tectonics, climate, and fluvial transport during the Cenozoic evolution of the Ebro Basin (NE Iberia), *Journal of Geophysical Research-Solid Earth*, 108, 10.1029/2002jb002073, 2003.
- 510 Gilbert, G. K.: Report on the Geology of the Henry Mountains (Utah), United States Geological Survey, Washington D.C., 1877.
- Goren, L., Willett, S. D., Herman, F., and Braun, J.: Coupled numerical-analytical approach to landscape evolution modeling, *Earth Surface Processes and Landforms*, 39, 522-545, 10.1002/esp.3514, 2014.
- Goren, L., Castellort, S., and Klinger, Y.: Modes and rates of horizontal deformation from rotated river basins: Application to the Dead Sea fault system in Lebanon, *Geology*, 43, 843-846, 10.1130/G36841.1, 2015.
- 515 Goren, L.: A theoretical model for fluvial channel response time during time-dependent climatic and tectonic forcing and its inverse applications, *Geophysical Research Letters*, 43, 10753-10763, 10.1002/2016gl070451, 2016.
- Guerit, L., Goren, L., Dominguez, S., Malavieille, J., and Castellort, S.: Landscape 'stress' and reorganization from chi-maps: Insights from experimental drainage networks in oblique collision setting, *Earth Surface Processes and Landforms*, 43, 3152-3163, 10.1002/esp.4477, 2018.
- 520 Handy, M. R., Ustaszewski, K., and Kissling, E.: Reconstructing the Alps–Carpathians–Dinarides as a key to understanding switches in subduction polarity, slab gaps and surface motion, *International Journal of Earth Sciences*, 104, 1-26, 10.1007/s00531-014-1060-3, 2015.
- Headley, R. M.: River redirected, *Nature Geoscience*, 10, 327, 10.1038/ngeo2946, 2017.
- Hergarten, S.: Self organized criticality in earth systems, Springer, Berlin, 272 pp., 2002.
- Hergarten, S., Robl, J., and Stüwe, K.: Extracting topographic swath profiles across curved geomorphic features, *Earth Surf. Dynam.*, 2, 97-104, 10.5194/esurf-2-97-2014, 2014.
- 525 Hergarten, S., Robl, J., and Stüwe, K.: Tectonic geomorphology at small catchment sizes – extensions of the stream-power approach and the γ method, *Earth Surface Dynamics*, 4, 1-9, 10.5194/esurf-4-1-2016, 2016.
- Herman, F., Seward, D., Valla, P. G., Carter, A., Kohn, B., Willett, S. D., and Ehlers, T. A.: Worldwide acceleration of mountain erosion under a cooling climate, *Nature*, 504, 423-426, 2013.
- 530 Hodges, K. V., Wobus, C., Ruhl, K., Schildgen, T., and Whipple, K.: Quaternary deformation, river steepening, and heavy precipitation at the front of the Higher Himalayan ranges, *Earth and Planetary Science Letters*, 220, 379-389, 10.1016/s0012-821x(04)00063-9, 2004.
- Houseman, G., and England, P.: Finite strain calculations of continental deformation. 1. Method and general results for convergent zones, *Journal of Geophysical Research-Solid Earth and Planets*, 91, 3651-3663, 10.1029/JB091iB03p03651, 1986.
- Howard, A. D.: Thresholds in river regimes, *Thresholds in geomorphology*, edited by: D., J., George Allen and Unwin, London, United Kingdom (GBR), 227-258 pp., 1980.
- 535 Howard, A. D.: A detachment-limited model of drainage-basin evolution, *Water Resources Research*, 30, 2261-2285, 10.1029/94wr00757, 1994.
- Keil, M., and Neubauer, F.: Initiation and development of a fault-controlled, orogen-parallel overdeepened valley: The Upper Enns Valley, Austria, *Austrian Journal of Earth Sciences*, 102, 80-90, 10.1127/0372-8854/2012/0074, 2009.
- 540 Kirby, E., and Whipple, K. X.: Expression of active tectonics in erosional landscapes, *Journal of Structural Geology*, 44, 54-75, 10.1016/j.jsg.2012.07.009, 2012.
- Kooi, H., and Beaumont, C.: Large-scale geomorphology: Classical concepts reconciled and integrated with contemporary ideas via a surface processes model, *Journal of Geophysical Research-Solid Earth*, 101, 3361-3386, 10.1029/95jb01861, 1996.
- Kuhlemann, J., Frisch, W., Dunkl, I., Szekely, B., and Spiegel, C.: Miocene shifts of the drainage divide in the Alps and their foreland basin, *Zeitschrift Fur Geomorphologie*, 45, 239-265, 2001.
- 545 Kuhlemann, J., Frisch, W., Szekely, B., Dunkl, I., and Kazmer, M.: Post-collisional sediment budget history of the Alps: tectonic versus climatic control, *International Journal of Earth Sciences*, 91, 818-837, 10.1007/s00531-002-0266-y, 2002.
- Kuhlemann, J., Dunkl, I., Brugel, A., Spiegel, C., and Frisch, W.: From source terrains of the Eastern Alps to the Molasse Basin: Detrital record of non-steady-state exhumation, *Tectonophysics*, 413, 301-316, 10.1016/j.tecto.2005.11.007, 2006.
- 550 Kuhlemann, J.: Paleogeographic and paleotopographic evolution of the Swiss and Eastern Alps since the Oligocene, *Global and Planetary Change*, 58, 224-236, 10.1016/j.gloplacha.2007.03.007, 2007.
- Kühni, A., and Pfiffner, O. A.: The relief of the Swiss Alps and adjacent areas and its relation to lithology and structure: topographic analysis from a 250-m DEM, *Geomorphology*, 41, 285-307, https://doi.org/10.1016/S0169-555X(01)00060-5, 2001.
- 555 Miller, S. R., and Slingerland, R. L.: Topographic advection on fault-bend folds: Inheritance of valley positions and the formation of wind gaps, *Geology*, 34, 769-772, 10.1130/g22658.1, 2006.
- Miller, S. R., Baldwin, S. L., and Fitzgerald, P. G.: Transient fluvial incision and active surface uplift in the Woodlark Rift of eastern Papua New Guinea, *Lithosphere*, 4, 131-149, 10.1130/L135.1, 2012.
- Molnar, P., and Lyon-Caen, H.: Some simple physical aspects of the support, structure, and evolution of mountain belts



- Processes in Continental Lithospheric Deformation, in: Geological Society of America Special Paper, edited by: Clark, S. P., Jr., Burchfiel, B. C., and Suppe, J., Geological Society of America, 0, 1988.
- 560 Molnar, P.: Nature, nurture and landscape, *Nature*, 426, 612-613, 10.1038/426612a, 2003.
- Montgomery, D. R., Balco, G., and Willett, S. D.: Climate, tectonics, and the morphology of the Andes, *Geology*, 29, 579-582, 10.1130/0091-7613(2001)029<0579:ctatmo>2.0.co;2, 2001.
- Neubauer, F.: Formation of an intra-orogenic transtensional basin: the Neogene Wagrain basin in the Eastern Alps, *Swiss Journal of Geosciences*, 109, 37-56, 10.1007/s00015-016-0206-7, 2016.
- 565 Norton, K. P., Abbuhl, L. M., and Schlunegger, F.: Glacial conditioning as an erosional driving force in the Central Alps, *Geology*, 38, 655-658, 10.1130/g31102.1, 2010.
- Perron, J. T., and Royden, L.: An integral approach to bedrock river profile analysis, *Earth Surface Processes and Landforms*, 38, 570-576, 10.1002/esp.3302, 2013.
- 570 Prasicsek, G., Larsen, I. J., and Montgomery, D. R.: Tectonic control on the persistence of glacially sculpted topography, *Nature Communications*, 6, 8028, 10.1038/ncomms9028
<https://www.nature.com/articles/ncomms9028#supplementary-information>, 2015.
- Ratschbacher, L., Frisch, W., Neubauer, F., Schmid, S. M., and Neugebauer, J.: Extension in compressional orogenic belts: the eastern Alps, *Geology*, 17, 404-407, 1989.
- 575 Ratschbacher, L., Frisch, W., Linzer, H.-G., and Merle, O.: Lateral extrusion in the Eastern Alps, part 2: structural analysis, *Tectonics*, 10, 257-271, 1991.
- Reiners, P. W., Ehlers, T. A., Mitchell, S. G., and Montgomery, D. R.: Coupled spatial variations in precipitation and long-term erosion rates across the Washington Cascades, *Nature*, 426, 645-647, 10.1038/nature02111, 2003.
- Robl, J., and Stüwe, K.: Continental collision with finite indenter strength: 1. Concept and model formulation, *Tectonics*, 24, 10.1029/2004tc001727, 2005a.
- 580 Robl, J., and Stüwe, K.: Continental collision with finite indenter strength: 2. European Eastern Alps, *Tectonics*, 24, 1-21, 10.1029/2004TC001741, 2005b.
- Robl, J., Hergarten, S., and Stüwe, K.: Morphological analysis of the drainage system in the Eastern Alps, *Tectonophysics*, 460, 263-277, 10.1016/j.tecto.2008.08.024, 2008a.
- 585 Robl, J., Stüwe, K., Hergarten, S., and Evans, L.: Extension during continental convergence in the Eastern Alps: The influence of orogen-scale strike-slip faults, *Geology*, 36, 963-966, 2008b.
- Robl, J., Prasicsek, G., Hergarten, S., and Stüwe, K.: Alpine topography in the light of tectonic uplift and glaciation, *Global and Planetary Change*, 127, 34-49, 10.1016/j.gloplacha.2015.01.008, 2015.
- Robl, J., Heberer, B., Prasicsek, G., Neubauer, F., and Hergarten, S.: The topography of a continental indenter: The interplay between crustal deformation, erosion, and base level changes in the eastern Southern Alps, *Journal of Geophysical Research: Earth Surface*, 122, 310-334, 10.1002/2016JF003884, 2017a.
- 590 Robl, J., Hergarten, S., and Prasicsek, G.: The topographic state of fluvially conditioned mountain ranges, *Earth-Science Reviews*, 168, 190-217, 10.1016/j.earscirev.2017.03.007, 2017b.
- Roe, G. H., Whipple, K. X., and Fletcher, J. K.: Feedbacks among climate, erosion, and tectonics in a critical wedge orogen, *American Journal of Science*, 308, 815-842, 10.2475/07.2008.01, 2008.
- 595 Royden, L., and Perron, J. T.: Solutions of the stream power equation and application to the evolution of river longitudinal profiles, *Journal of Geophysical Research: Earth Surface*, 118, 497-518, 10.1002/jgrf.20031, 2013.
- Royden, L. H., Burchfiel, B. C., King, R. W., Wang, E., Chen, Z. L., Shen, F., and Liu, Y. P.: Surface deformation and lower crustal flow in eastern Tibet, *Science*, 276, 788-790, 10.1126/science.276.5313.788, 1997.
- 600 Salcher, B. C., Kober, F., Kissling, E., and Willett, S. D.: Glacial impact on short-wavelength topography and long-lasting effects on the denudation of a deglaciated mountain range, *Global and Planetary Change*, 115, 59-70, 10.1016/j.gloplacha.2014.01.002, 2014.
- Schmidt, K. M., and Montgomery, D. R.: Limits to relief, *Science*, 270, 617-620, 10.1126/science.270.5236.617, 1995.
- Schwanghart, W., and Scherler, D.: Short Communication: TopoToolbox 2-MATLAB-based software for topographic analysis and modeling in Earth surface sciences, *Earth Surface Dynamics*, 2, 1-7, 10.5194/esurf-2-1-2014, 2014.
- 605 Shugar, D. H., Clague, J. J., Best, J. L., Schoof, C., Willis, M. J., Copland, L., and Roe, G. H.: River piracy and drainage basin reorganization led by climate-driven glacier retreat, *Nature Geoscience*, 10, 370, 10.1038/ngeo2932, 2017.
- Snyder, N. P., Whipple, K. X., Tucker, G. E., and Merritts, D. J.: Landscape response to tectonic forcing: Digital elevation model analysis of stream profiles in the Mendocino triple junction region, Northern California, *Bulletin of the Geological Society of America*, 112, 1250-1263, 10.1130/0016-7606(2000)112<1250:LRTTFD>2.0.CO;2, 2000.
- 610 Stüwe, K., Robl, J., Hergarten, S., and Evans, L.: Modeling the influence of horizontal advection, deformation, and late uplift on the drainage development in the India-Asia collision zone, *Tectonics*, 27, 10.1029/2007tc002186, 2008.
- Tapponnier, P., Peltzer, G., Ledain, A. Y., Armijo, R., and Cobbold, P.: Propagating extrusion tectonics in Asia - New insights from simple experiments with plasticine, *Geology*, 10, 611-616, 10.1130/0091-7613(1982)10<611:petian>2.0.co;2, 1982.



- 615 Tucker, G. E., and Slingerland, R. L.: Erosional dynamics, flexural isostasy, and long-lived escarpments - A numerical modeling study, *Journal of Geophysical Research-Solid Earth*, 99, 12229-12243, 10.1029/94jb00320, 1994.
- Valla, P. G., van der Beek, P. A., and Lague, D.: Fluvial incision into bedrock: Insights from morphometric analysis and numerical modeling of gorges incising glacial hanging valleys (Western Alps, France), *J Geophys Res-Earth*, 115, 10.1029/2008jf001079, 2010.
- van der Beek, P., and Bourbon, P.: A quantification of the glacial imprint on relief development in the French western Alps, *Geomorphology*, 97, 52-72, 10.1016/j.geomorph.2007.02.038, 2008.
- 620 Wang, X., and Neubauer, F.: Orogen-parallel strike-slip faults bordering metamorphic core complexes: the Salzach-Enns fault zone in the eastern Alps, Austria, *Journal of Structural Geology*, 20, 799-818, 10.1016/S0191-8141(98)00013-3, 1998.
- Whipple, K. X.: Fluvial landscape response time: How plausible is steady-state denudation?, *American Journal of Science*, 301, 313-325, 10.2475/ajs.301.4-5.313, 2001.
- Whipple, K. X., DiBiase, R. A., and Crosby, B. T.: *Bedrock Rivers*, Elsevier Ltd., 550-573 pp., 2013.
- 625 Whipple, K. X., Forte, A. M., DiBiase, R. A., Gasparini, N. M., and Ouimet, W. B.: Timescales of landscape response to divide migration and drainage capture: Implications for the role of divide mobility in landscape evolution, *J Geophys Res-Earth*, 122, 248-273, 10.1002/2016jf003973, 2017.
- Willett, S. D.: Orogeny and orography: The effects of erosion on the structure of mountain belts, *Journal of Geophysical Research-Solid Earth*, 104, 28957-28981, 10.1029/1999jb900248, 1999.
- 630 Willett, S. D., Slingerland, R., and Hovius, N.: Uplift, shortening, and steady state topography in active mountain belts, *American journal of Science*, 301, 455-485, 2001.
- Willett, S. D., McCoy, S. W., Perron, J. T., Goren, L., and Chen, C. Y.: Dynamic reorganization of river basins, *Science*, 343, 1248765, 10.1126/science.1248765, 2014.
- Winterberg, S., and Willett, S. D.: Greater Alpine river network evolution, interpretations based on novel drainage analysis, *Swiss Journal of Geosciences*, 112, 3-22, 10.1007/s00015-018-0332-5, 2019.
- 635 Wobus, C., Whipple, K. X., Kirby, E., Snyder, N. P., Johnson, J., Spyropolou, K., Crosby, B., and Sheehan, D.: Tectonics from topography: Procedures, promise, and pitfalls, in: *Tectonics, Climate, and Landscape Evolution*, edited by: Willett, S. D., Hovius, N., Brandon, M. T., and Fisher, D. M., Geological Society of America Special Papers, 55-74, 2006.
- Wobus, C. W., Hodges, K. V., and Whipple, K. X.: Has focused denudation sustained active thrusting at the Himalayan topographic front?, *Geology*, 31, 861-864, 10.1130/g19730.1, 2003.
- 640 Yang, R., Willett, S. D., and Goren, L.: In situ low-relief landscape formation as a result of river network disruption, *Nature*, 520, 526-529, 10.1038/nature14354, 2015.
- Yang, R., Fellin, M. G., Herman, F. d. r., Willett, S. D., Wang, W., and Maden, C.: Spatial and temporal pattern of erosion in the Three Rivers Region, southeastern Tibet, *Earth and Planetary Science Letters*, 433, 10-20, 10.1016/j.epsl.2015.10.032, 2016.
- 645 Yanites, B. J., Ehlers, T. A., Becker, J. K., Schnellmann, M., and Heuberger, S.: High magnitude and rapid incision from river capture: Rhine River, Switzerland, *J Geophys Res-Earth*, 118, 1060-1084, 10.1002/jgrf.20056, 2013.

650

655

660





Figure 1. Topographic map of the study area and the drainage pattern of the Eastern Alps. The inset shows the position of the study within the European Alps. Blue lines indicate the drainage pattern, whereby the line width is proportional to \log_{10} of the contributing drainage area. Drainage divides are shown by thick white lines. Major faults are indicated by solid black lines and the direction of motion is shown by arrow. The red line and the grey hull indicate the course of the swath profiles shown on Figure 3.


670



675 **Figure 2.** Drainage pattern of the Eastern Alps calculated for increasing baselevels and color -coded for χ . All streams with a contributing drainage area larger than 1 km² are shown. The line width of the channels is proportional to $\log_{10}(\text{drainage area})$. White lines and annotations represent the major drainage divides. (A) Baselevel set to 400 m of elevation. (B) Baselevel set to 600 m of elevation. (C) Baselevel set to 800 m of elevation. (D) Baselevel set to 1000 m of elevation.

680

685



690


695

Figure 3. χ values measured at channel heads (Forte and Whipple, 2018) of the investigated catchments. Histograms with a black outline represent the Salzach drainage basin. Histograms with a blue filling represent the adjacent Inn/Adige (WS 1), Saalach (WS 2), Enns (WS 3) and Mur (WS 4) drainage basins. Data are divided in 10 equally-spaced bins. Error bars indicate the standard deviation and filled circles are the mean values of the dataset. n is the total number of data points.

705

710


715



720 **Figure 4.** Generalized swath profiles (Hergarten et al., 2014) along the profile lines shown in Figure 1. The profiles have a half-width
of 5 km to each side of the profile line (drainage divide). The black line indicates the mean elevation. The hull of the blue area is the
standard deviation and the hull of the grey area the extreme values within the swath segments.

725


730



735 **Figure 5.** Gilbert metric histograms (Forte and Whipple, 2018) for the investigated watersheds of the study area. Histograms with a
blue filling represent the adjacent Inn/Adige (WS 1), Saalach (WS 2), Enns (WS 3) and Mur (WS 4) drainage basins. Data are
divided in 10 equally-spaced bins. Error bars indicate the standard deviation and filled circles are the mean values of the dataset. n
is the total number of data points.

740


745



750 **Figure 6.** Proposed direction of drainage migration of the main drainage divides of the study area. (A) The major drainage basins
 are annotated. Orange arrows indicate the proposed migration direction of divides based on χ . Gilbert metrics for each investigated
 catchment are shown. Positive Δ Elevation and $\Delta\chi$ values as well as negative Δ Relief and Δ Gradient indicate migration towards the
 Salzach drainage basin. (B) Normalized Δ plot of Gilbert metrics. Negative values of Δ Relief and Δ Gradient are standardized to
 755 positive values such that all positive values indicate a migration towards the Salzach basin. Error bar and filled circles indicate 1 –
 standard deviation and mean values, respectively.

760

765



770

Figure 7. χ colored drainage pattern of reconstructed paleo drainage geometries of the Eastern Alps. All streams with a contributing drainage area larger than 1 km^2 are shown and the line width of the channels is proportional to $\log_{10}(\text{drainage area})$. The baselevel for χ computation is set to 400 m. White lines and annotations represent the major drainage divides. (A) Enns drainage path - drainage scenario assuming an elongated Enns catchment representing the Mid-Miocene situation as suggest by Frisch et al. (1998). (B) Saalach drainage path - drainage scenario assuming that the Saalach captured the westernmost part of the Paleo-Enns catchment. (C) Salzach drainage path - drainage scenario assuming that both Saalach and Salzach took over parts of the Paleo-Enns catchment as suggested by Robl et al. (2008a). (D) Present day drainage path for comparison.